

# Lactéol<sup>®</sup> fort

## capsules

### Dosage and administration

To be taken at the first sign of diarrhea.

Number of capsules	First day		Following days	
	1st dose	2nd dose	1st dose	2nd dose
Children	2	2	2	2
Adults	4	2	2	2
Contents of capsule can be mixed into water				

The 2nd dose can be taken 6-8 hours after the 1st dose.

### Usage during pregnancy

No known teratogenic effect has been reported.

### Adverse side effects

No side effect has been so far reported.

### Incompatibilities and interactions

No known incompatibilities and drug interactions have been reported.

### Stability and storage

Shelf life : 3 years

Keep in cool dry place, store below 25°C

### Non-poison drug

**Adare Pharmaceuticals SAS**  
Route de Bû - 78550 Houdan  
FRANCE

**ADARE**  
Pharmaceuticals<sup>®</sup>

# Lactéol<sup>®</sup> fort

## capsules

---

### Presentation

Strip-seal : Boxes of 12, 100 capsules.

### Composition (1 capsule)

#### • *Lactobacillus LB*

(Lyophilized killed microbial bodies):.....5 billion

• Fermented culture medium (neutralized):.....80mg

Excipients: Lactose monohydrate, calcium carbonate, silicic acid, banana-orange flavour, saccharose, q.s.....235mg

### Pharmacological properties

Antidiarrheal of microbial origin, with :

- direct bacteriostatic action and stimulation of the growth of the defensive acidogenic intestinal flora,
- non-specific immunostimulation of the mucosa (increased synthesis of IgA).
- inhibition of adherence of invasive or toxinogenic enteropathogenic organisms (*Escherichia coli*, *Campylobacter jejuni*) to epithelial cells

### Indications

Symptomatic treatment of diarrhea.

### Disposal of product

Discard the capsules after the expiry date and throw contents into the dustbin.

---

*Information contained in leaflet is limited and further information can be obtained from other sources, including the doctor and the pharmacist.*