

# Lactéol<sup>®</sup> fort

## sachets

### Dosage and administration

To be taken at the first sign of diarrhoea.

Number of sachets	First day		Following days	
	1st dose	2nd dose	1st dose	2nd dose
Infants (*)	1	1	1	1
Children	1 to 2	1	1	1
Adults	2	1	1	1
Contents of sachets can be mixed into water				

The 2nd dose can be taken 6-8 hours after the 1st dose.

### Usage during pregnancy

No known teratogenic effect has been reported.

### Adverse side effects

No side effect has been so far reported.

### Incompatibilities and interactions

No known incompatibilities and drug interactions have been reported.

### Stability and storage

Shelf life: 3 years

Keep in cool dry place, store below 25°C

### Non-poison drug

# Lactéol<sup>®</sup> fort

## sachets

---

### Presentation

Strip-seal: Boxes of 6, 100 sachets.

### Composition (1 sachet)

- Lactobacillus LB

(lyophilized inactive microbial bodies) ..... 10 billion

- Fermented culture medium (neutralized) ..... 160mg

Excipients: Lactose monohydrate, calcium carbonate, silicic acid, banana-orange flavour, saccharose, q.s ..... 800mg

### Pharmacological properties

Antidiarrheal of microbial origin, with:

- Direct bacteriostatic action and stimulation of the growth of the defensive acidogenic intestinal flora,
- Non-specific immunostimulation of the mucosa (increased synthesis of IgA),
- Inhibition of adherence of invasive or toxinogenic enteropathogenic organisms (Escherichia Coli, Campylobacter Jejuni) to epithelial cells.

### Indications

Symptomatic treatment of diarrhoea.

### Disposal of product

Discard the sachets after the expiry date and throw contents into the dustbin.

---

*Information contained in leaflet is limited and further information can be obtained from other sources, including the doctor and the pharmacist.*