

NAME OF THE MEDICINAL PRODUCT

Protopic Ointment 0.03%

Protopic Ointment 0.1%

QUALITATIVE AND QUANTITATIVE COMPOSITION

1g of Protopic 0.03% ointment contains 0.3mg of tacrolimus as tacrolimus monohydrate (0.03%).

1g of Protopic 0.1% ointment contains 1.0mg of tacrolimus as tacrolimus monohydrate (0.1%).

For excipients, see List of excipients.

PHARMACEUTICAL FORM

Ointment for external application on skin.

A white to slightly yellowish ointment.

CLINICAL PARTICULARS

Therapeutic Indications

Protopic 0.03% ointment is indicated in adults, adolescents and children from the age of 2 years.

Protopic 0.1 % ointment is indicated in adults and adolescents (16 years of age and above)

Flare treatment

Adults and adolescents (16 years of age and above)

Treatment of moderate to severe atopic dermatitis in adults and adolescents who are not adequately responsive to or are intolerant of conventional therapies such as topical corticosteroids.

Children (2 years of age and above)

Treatment of moderate to severe atopic dermatitis in children (2 years of age and above) who failed to respond adequately to conventional therapies such as topical corticosteroids.

Maintenance treatment

Treatment of moderate to severe atopic dermatitis for the prevention of flares and the prolongation of flare-free intervals in patients experiencing a high frequency of disease exacerbations (i.e. occurring 4 or more times per year) who have had an initial response to a maximum of 6 weeks treatment of twice daily tacrolimus ointment (lesions cleared, almost cleared or mildly affected).

Posology and Method of Administration

Protopic treatment should be initiated by physicians with experience in the diagnosis and treatment of atopic dermatitis.

Protopic is available in two strengths, Protopic 0.03 % and Protopic 0.1 % ointment.

Method of administration

Protopic ointment should be applied as a thin layer to affected or commonly affected areas of the skin. Protopic ointment may be used on any part of the body, including face, neck and flexure areas, except on mucous membranes. Protopic ointment should not be applied under occlusion because this method of administration has not been studied in patients (see Special warnings and special precautions for use).

Posology

Flare treatment

Protopic can be used for short-term and intermittent long-term treatment. Treatment should not be continuous on a long-term basis.

Protopic treatment should begin at the first appearance of signs and symptoms. Each affected region of the skin should be treated with Protopic until lesions are cleared, almost cleared or mildly affected. At the first signs of recurrence (flares) of the disease symptoms, treatment should be re-initiated.

Adult and adolescents (16 years of age and above)

Treatment should be started with Protopic 0.1% twice a day and treatment should be continued until clearance of the lesion. If symptoms recur, twice daily treatment with Protopic 0.1% should be restarted. An attempt should be made to reduce the frequency of application or to use the lower strength Protopic 0.03% ointment if the clinical condition allows.

Generally, improvement is seen within one week of starting treatment. If no signs of improvement are seen after two weeks of treatment, further treatment options should be considered.

Paediatric population

Children (2 years of age and above) should use the lower strength Protopic 0.03% ointment.

Treatment should be started twice a day for up to three weeks. Afterwards the frequency of application should be reduced to once a day until clearance of the lesion (see special warnings and special precautions for use).

Protopic ointment should not be used in children aged below 2 years until further data are available.

Elderly patients (65 years of age and above)

Specific studies have not been conducted in elderly patients. However, the clinical experience available in this patient population has not shown the necessity for any dosage adjustment.

Maintenance treatment

Patients who are responding to up to 6 weeks treatment using tacrolimus ointment twice daily (lesions cleared, almost cleared or mildly affected) are suitable for maintenance treatment.

Adult and adolescents (16 years of age and above)

Adult patients should use Protopic 0.1% ointment.

Protopic ointment should be applied once a day twice weekly (e.g. Monday and Thursday) to areas commonly affected by atopic dermatitis to prevent progression to flares. Between applications there should be 2-3 days without Protopic treatment.

After 12 months treatment, a review of the patient's condition should be conducted by the physician and a decision taken whether to continue maintenance treatment in the absence of safety data for maintenance treatment beyond 12 months.

If signs of a flare reoccur, twice daily treatment should be re-initiated (see Flare treatment section above).

Paediatric population

Children (2 years of age and above) should use the lower strength Protopic 0.03% ointment.

Protopic ointment should be applied once a day twice weekly (e.g. Monday and Thursday) to areas commonly affected by atopic dermatitis to prevent progression to flares. Between applications there should be 2-3 days without Protopic treatment.

The review of the child's condition after 12 months treatment should include suspension of treatment

to assess the need to continue this regimen and to evaluate the course of the disease.

Protopic ointment should not be used in children aged below 2 years until further data are available.

Elderly patients (65 years of age and above)

Specific studies have not been conducted in elderly patients (see Flare treatment section above).

Contraindications

Hypersensitivity to the active substance, macrolides in general or to any of the excipients.

Special Warnings and Special Precautions for Use

Protopic should not be used in patients with congenital or acquired immunodeficiencies or in patients on therapy that cause immunosuppression.

The effect of treatment with Protopic ointment on the developing immune system of children aged below 2 years, has not been established (see Therapeutic indications).

Exposure of the skin to sunlight should be minimised and the use of ultraviolet (UV) light from a solarium, therapy with UVB or UVA in combination with psoralens (PUVA) should be avoided during use of Protopic ointment (see Preclinical safety data). Physicians should advise patients on appropriate sun protection methods, such as minimisation of the time in the sun, use of a sunscreen product and covering of the skin with appropriate clothing. Protopic ointment should not be applied to lesions that are considered to be potentially malignant or pre-malignant.

Emollients should not be applied to the same area within 2 hours of applying Protopic ointment. Concomitant use of other topical preparations has not been assessed. There is no experience with concomitant use of systemic steroids or immunosuppressive agents.

Protopic ointment has not been evaluated for its efficacy and safety in the treatment of clinically infected atopic dermatitis. Before commencing treatment with Protopic ointment, clinical infections at treatment sites should be cleared. Patients with atopic dermatitis are predisposed to superficial skin infections. Treatment with Protopic may be associated with an increased risk of folliculitis and herpes viral infections (herpes simplex dermatitis [eczema herpeticum], herpes simplex [cold sores], Kaposi's varicelliform eruption) (see Undesirable effects). In the presence of these infections, the balance of risks and benefits associated with Protopic use should be evaluated.

The potential for local immunosuppression (possibly resulting in infections or cutaneous malignancies) in the long term (i.e. over a period of years) is unknown (see Pharmacodynamic properties).

Protopic contains the active substance tacrolimus, a calcineurin inhibitor. In transplant patients, prolonged systemic exposure to intense immunosuppression following systemic administration of calcineurin inhibitors has been associated with an increased risk of developing lymphomas and skin malignancies. In patients using tacrolimus ointment, cases of malignancies, including cutaneous and other types of lymphoma, and skin cancers have been reported (see Undesirable effects). Patients with atopic dermatitis treated with Protopic have not been found to have significant systemic tacrolimus levels.

Lymphadenopathy was uncommonly (0.8%) reported in clinical trials. The majority of these cases were related to infections (skin, respiratory tract, tooth) and resolved with appropriate antibiotic therapy. Transplant patients receiving immunosuppressive regimens (e.g. systemic tacrolimus) are at increased risk for developing lymphoma; therefore patients who receive Protopic and who develop

lymphadenopathy should be monitored to ensure that the lymphadenopathy resolves. Lymphadenopathy present at initiation of therapy should be investigated and kept under review. In case of persistent lymphadenopathy, the aetiology of the lymphadenopathy should be investigated. In the absence of a clear aetiology for the lymphadenopathy or in the presence of acute infectious mononucleosis, discontinuation of Protopic should be considered.

Care should be taken to avoid contact with eyes and mucous membranes. If accidentally applied to these areas, the ointment should be thoroughly wiped off and/or rinsed off with water. The use of Protopic ointment under occlusion has not been studied in patients. Occlusive dressings are not recommended.

As with any topical medicinal product, patients should wash their hands after application if the hands are not intended for treatment.

Tacrolimus is extensively metabolised in the liver and although blood concentrations are low following topical therapy, the ointment should be used with caution in patients with hepatic failure (see Pharmacokinetic properties).

The use of tacrolimus ointment is not recommended in patients with a skin barrier defect, such as Netherton's syndrome, lamellar ichthyosis, generalized erythroderma or cutaneous Graft Versus Host Disease. These skin conditions may increase systemic absorption of tacrolimus. Oral use of tacrolimus is also not recommended to treat these skin conditions. Post-marketing cases of increased tacrolimus blood level have been reported in these conditions.

Care should be exercised if applying Protopic to patients with extensive skin involvement over an extended period of time, especially in children (see Posology and method of administration).

The development of any new change different from previous eczema within a treated area should be reviewed by the physician.

Interactions with Other Medicinal Products and Other Forms of Interaction

Formal topical drug interaction studies with tacrolimus ointment have not been conducted.

Tacrolimus is not metabolised in human skin, indicating that there is no potential for percutaneous interactions that could affect the metabolism of tacrolimus.

Systemically available tacrolimus is metabolised via the hepatic Cytochrome P450 3A4 (CYP3A4). Systemic exposure from topical application of tacrolimus ointment is low (<1.0 ng/ml) and is unlikely to be affected by concomitant use of substances known to be inhibitors of CYP3A4. However, the possibility of interactions cannot be ruled out and the concomitant systemic administration of known CYP3A4 inhibitors (e.g. erythromycin, itraconazole, ketoconazole and diltiazem) in patients with widespread and/or erythrodermic disease should be done with caution.

Paediatric population

An interaction study with protein-conjugated vaccine against *Neisseria meningitidis* serogroup C has been investigated in children aged 2-11 years. No effect on immediate response to vaccination, the generation of immune memory, or humoral and cell-mediated immunity has been observed (see Pharmacodynamic properties).

Pregnancy and Lactation

Fertility

There are no fertility data available.

Pregnancy

There are no adequate data from the use of tacrolimus ointment in pregnant women. Studies in animals have shown reproductive toxicity following systemic administration (see Preclinical safety data). The potential risk for humans is unknown.

Protopic ointment should not be used during pregnancy unless clearly necessary.

Breastfeeding

Human data demonstrate that, after systemic administration, tacrolimus is excreted into breast milk. Although clinical data have shown that systemic exposure from application of tacrolimus ointment is low, breast-feeding during treatment with Protopic ointment is not recommended.

Effects on Ability to Drive and Use Machines

Protopic ointment is administered topically and is unlikely to have an effect on the ability to drive or use machines.

Undesirable Effects

In clinical studies approximately 50% of patients experienced some type of skin irritation adverse reaction at the site of application. Burning sensation and pruritus were very common, usually mild to moderate in severity and tended to resolve within one week of starting treatment. Erythema was a common skin irritation adverse reaction. Sensation of warmth, pain, paraesthesia and rash at the site of application were also commonly observed. Alcohol intolerance (facial flushing or skin irritation after consumption of an alcoholic beverage) was common.

Patients may be at an increased risk of folliculitis, acne and herpes viral infections.

Adverse reactions with suspected relationship to treatment are listed below by system organ class. Frequencies are defined as very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$) and uncommon ($\geq 1/1000$ to $< 1/100$). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

General disorders and administration site conditions

Very common: Application site burning, application site pruritus

Common: Application site warmth, application site erythema, application site pain, application site irritation, application site paraesthesia, application site rash

Not known (cannot be estimated from the available data): Application site oedema*

Infections and infestations

Common: Local skin infection regardless of specific etiology including but not limited to: Eczema herpeticum, Folliculitis, Herpes simplex, Herpes virus infection, Kaposi's varicelliform eruption*

Investigations

Not known (cannot be estimated from the available data): Drug level increased* (see Special warnings and special precautions for use)

Skin and subcutaneous tissue disorders

Common: Pruritus

Uncommon: Acne*

Not known (cannot be estimated from the available data): Rosacea*

Nervous system disorders

Common: Paraesthesias and dysaesthesias (hyperaesthesia, burning sensation)

Metabolism and nutrition disorders

Common: Alcohol intolerance (facial flushing or skin irritation after consumption of an alcoholic beverage)

* The adverse reaction has been reported during post-marketing experience.

Post-marketing

Cases of malignancies, including cutaneous and other types of lymphoma, and skin cancers, have been reported in patients using tacrolimus ointment (see Special warnings and special precautions for use).

Maintenance treatment

In a study of maintenance treatment (twice weekly treatment) in adults and children with moderate and severe atopic dermatitis, the following adverse events were noted to occur more frequently than in the control group: application site impetigo (7.7% in children) and application site infections (6.4% in children and 6.3% in adults).

Paediatric population

Frequency, type and severity of adverse reactions in children are similar to those reported in adults.

Overdose

Overdosage following topical administration is unlikely.

If ingested, general supportive measures may be appropriate. These may include monitoring of vital signs and observation of clinical status. Due to the nature of the ointment vehicle, induction of vomiting or gastric lavage is not recommended.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic Properties

Pharmacotherapeutic group: Other dermatologicals, ATC code: D11AH01

Mechanism of action and pharmacodynamic effects

The mechanism of action of tacrolimus in atopic dermatitis is not fully understood.

While the following have been observed, the clinical significance of these observations in atopic dermatitis is not known. Via its binding to a specific cytoplasmic immunophilin (FKBP12), tacrolimus inhibits calcium-dependent signal transduction pathways in T cells, thereby preventing the transcription and synthesis of IL-2, IL-3, IL-4, IL-5 and other cytokines such as GM-CSF, TNF- α and IFN- γ .

In vitro, in Langerhans cells isolated from normal human skin, tacrolimus reduced the stimulatory activity towards T cells. Tacrolimus has also been shown to inhibit the release of inflammatory mediators from skin mast cells, basophils and eosinophils.

In animals, tacrolimus ointment suppressed inflammatory reactions in experimental and spontaneous dermatitis models that resemble human atopic dermatitis. Tacrolimus ointment did not reduce skin thickness and did not cause skin atrophy in animals.

In patients with atopic dermatitis, improvement of skin lesions during treatment with tacrolimus ointment was associated with reduced Fc receptor expression on Langerhans cells and a reduction of their hyperstimulatory activity towards T cells. Tacrolimus ointment does not affect collagen synthesis in humans.

Clinical efficacy and safety

The efficacy and safety of Protopic was assessed in more than 13,500 patients treated with tacrolimus ointment in Phase I to Phase III clinical trials. Data from five major trials are presented here.

In a six-month multicentre double-blind randomised trial, 0.1% tacrolimus ointment was administered twice-a-day to adults with moderate to severe atopic dermatitis and compared to a topical corticosteroid based regimen (0.1% hydrocortisone butyrate on trunk and extremities, 1% hydrocortisone acetate on face and neck). The primary endpoint was the response rate at month 3 defined as the proportion of patients with at least 60% improvement in the mEASI (modified Eczema Area and Severity Index) between baseline and month 3. The response rate in the 0.1% tacrolimus group (71.6%) was significantly higher than that in the topical corticosteroid based treatment group (50.8%; $p < 0.001$; Table 1). The response rates at month 6 were comparable to the 3-month results.

Table 1 Efficacy at month 3

| | Topical corticosteroid regimen§ (N=485) | Tacrolimus 0.1% (N=487) |
|--|---|-------------------------|
| Response rate of $\geq 60\%$ improvement in mEASI (Primary Endpoint)§§ | 50.8% | 71.6% |
| Improvement $\geq 90\%$ in Physician's Global Evaluation | 28.5% | 47.7% |

§ Topical corticosteroid regimen = 0.1% hydrocortisone butyrate on trunk and extremities, 1% hydrocortisone acetate on face and neck

§§ higher values = greater improvement

The incidence and nature of most adverse events were similar in the two treatment groups. Skin burning, herpes simplex, alcohol intolerance (facial flushing or skin sensitivity after alcohol intake), skin tingling, hyperaesthesia, acne and fungal dermatitis occurred more often in the tacrolimus treatment group. There were no clinically relevant changes in the laboratory values or vital signs in either treatment group throughout the study.

In the second trial, children aged from 2 to 15 years with moderate to severe atopic dermatitis received twice daily treatment for three weeks of 0.03% tacrolimus ointment, 0.1% tacrolimus ointment or 1% hydrocortisone acetate ointment. The primary endpoint was the area-under-the-curve (AUC) of the mEASI as a percentage of baseline averaged over the treatment period. The results of this multicentre, double-blind, randomised trial showed that tacrolimus ointment, 0.03% and 0.1%, is significantly more effective ($p < 0.001$ for both) than 1% hydrocortisone acetate ointment (Table 2).

Table 2 Efficacy at week 3

| | Hydrocortisone acetate 1% (N=185) | Tacrolimus 0.03% (N=189) | Tacrolimus 0.1% (N=186) |
|---|-----------------------------------|--------------------------|-------------------------|
| Median mEASI as Percentage of Baseline mean AUC (Primary Endpoint)§ | 64.0% | 44.8% | 39.8% |
| Improvement $\geq 90\%$ in Physician's Global Evaluation | 15.7% | 38.5% | 48.4% |

§ lower values = greater improvement

The incidence of local skin burning was higher in the tacrolimus treatment groups than in the hydrocortisone group. Pruritus decreased over time in the tacrolimus groups but not in the hydrocortisone group. There were no clinically relevant changes in the laboratory values or vital signs in either treatment group throughout the clinical trial.

The purpose of the third multicentre, double-blind, randomised study was the assessment of efficacy and safety of 0.03% tacrolimus ointment applied once or twice a day relative to twice daily administration of 1% hydrocortisone acetate ointment in children with moderate to severe atopic dermatitis. Treatment duration was for up to three weeks.

Table 3 Efficacy at week 3

| | Hydrocortisone acetate 1% Twice daily (N=207) | Tacrolimus 0.03% Once daily (N=207) | Tacrolimus 0.03% Twice daily (N=210) |
|--|---|-------------------------------------|--------------------------------------|
| Median mEASI Percentage Decrease (Primary Endpoint)§ | 47.2% | 70.0% | 78.7% |
| Improvement $\geq 90\%$ in Physician's Global Evaluation | 13.6% | 27.8% | 36.7% |

§ higher values = greater improvement

The primary endpoint was defined as the percentage decrease in mEASI from the baseline to end of treatment. A statistically significant better improvement was shown for once daily and twice daily 0.03% tacrolimus ointment compared to twice daily hydrocortisone acetate ointment ($p < 0.001$ for both). Twice daily treatment with 0.03% tacrolimus ointment was more effective than once daily administration (Table 3). The incidence of local skin burning was higher in the tacrolimus treatment groups than in the hydrocortisone group. There were no clinically relevant changes in the laboratory values or vital signs in either treatment group throughout the study.

In the fourth trial, approximately 800 patients (aged ≥ 2 years) received 0.1% tacrolimus ointment intermittently or continuously in an open-label, long-term safety study for up to four years, with 300 patients receiving treatment for at least three years and 79 patients receiving treatment for a minimum of 42 months. Based on changes from baseline in EASI score and body surface area

affected, patients regardless of age had improvement in their atopic dermatitis at all subsequent time points. In addition, there was no evidence of loss of efficacy throughout the duration of the clinical trial. The overall incidence of adverse events tended to decrease as the study progressed for all patients independent of age. The three most common adverse events reported were flu-like symptoms (cold, common cold, influenza, upper respiratory infection, etc.), pruritus and skin burning. No adverse events previously unreported in shorter duration and/or previous studies were observed in this long-term study.

The efficacy and safety of tacrolimus ointment in maintenance treatment of mild to severe atopic dermatitis was assessed in 524 patients in two Phase III multicentre clinical trials of similar design, one in adult patients (≥ 16 years) and one in paediatric patients (2-15 years). In both studies, patients with active disease entered an open-label period (OLP) during which they treated affected lesions with tacrolimus ointment twice daily until improvement had reached a predefined score (Investigator's Global Assessment [IGA] ≤ 2 , i.e. clear, almost clear or mild disease) for a maximum of 6 weeks. Thereafter, patients entered a double-blind disease control period (DCP) for up to 12 months. Patients were randomised to receive either tacrolimus ointment (0.1% adults; 0.03% children) or vehicle, once a day twice weekly on Mondays and Thursdays. If a disease exacerbation occurred, patients were treated with open-label tacrolimus ointment twice daily for a maximum of 6 weeks until the IGA score returned to ≤ 2 .

The primary endpoint in both studies was the number of disease exacerbations requiring a "substantial therapeutic intervention" during the DCP, defined as an exacerbation with an IGA of 3-5 (i.e. moderate, severe and very severe disease) on the first day of the flare, and requiring more than 7 days treatment. Both studies showed significant benefit with twice weekly treatment with tacrolimus ointment with regards to the primary and key secondary endpoints over a period of 12 months in a pooled population of patients with mild to severe atopic dermatitis. In a subanalysis of a pooled population of patients with moderate to severe atopic dermatitis these differences remained statistically significant (Table 4). No adverse events not reported previously were observed in these studies.

Table 4 Efficacy (moderate to severe subpopulation)

| | Tacrolimus 0.1% Twice weekly (N=80) | Vehicle Twice weekly (N=73) | Tacrolimus 0.03% Twice weekly (N=78) | Vehicle Twice weekly (N=75) |
|---|---|--------------------------------|--|--------------------------------|
| Median number of DEs requiring substantial intervention adjusted for time at risk (% of patients without DE requiring substantial intervention) | 1.0 (48.8%) | 5.3 (17.8%) | 1.0 (46.2%) | 2.9 (21.3%) |
| Median time to first DE requiring substantial intervention | 142 days | 15 days | 217 days | 36 days |
| Median number of DEs adjusted for time at risk (% of patients without any DE periods) | 1.0 (42.5%) | 6.8 (12.3%) | 1.5 (41.0%) | 3.5 (14.7%) |
| Median time to first DE | 123 days | 14 days | 146 days | 17 days |
| Mean (SD) percentage of days of DE | 16.1 (23.6) | 39.0 (27.8) | 16.9 (22.1) | 29.9 (26.8) |

| | | | | |
|------------------------|--|--|--|--|
| exacerbation treatment | | | | |
|------------------------|--|--|--|--|

DE: disease exacerbation

P<0.001 in favour of tacrolimus ointment 0.1% (adults) and 0.03% (children) for the primary and key secondary endpoints

A seven-month, double blind, randomised parallel group study of paediatric patients (2-11 years) with moderate to severe atopic dermatitis was performed. In one arm patients received Protopic 0.03% ointment (n=121) twice a day for 3 weeks and thereafter once a day until clearance. In the comparator arm patients received 1% hydrocortisone acetate ointment (HA) for head and neck and 0.1% hydrocortisone butyrate ointment for trunk and limbs (n=111) twice a day for 2 weeks and subsequently HA twice a day to all affected areas. During this period all patients and control subjects (n=44) received a primary immunisation and a rechallenge with a protein-conjugate vaccine against *Neisseria meningitidis* serogroup C.

The primary endpoint of this study was the response rate to vaccination, defined as the percentage of patients with a serum bactericidal antibody (SBA) titre ≥ 8 at the week 5 visit. Analysis of the response rate at week 5 showed equivalence between the treatment groups (hydrocortisone 98.3%, tacrolimus ointment 95.4%; 7-11 years: 100% in both arms). The results in the control group were similar.

The primary response to vaccination was not affected.

Pharmacokinetic Properties

Clinical data have shown that tacrolimus concentrations in systemic circulation after topical administration are low and, when measurable, transient.

Absorption

Data from healthy human subjects indicate that there is little or no systemic exposure to tacrolimus following single or repeated topical application of tacrolimus ointment.

Most atopic dermatitis patients (adults and children) treated with single or repeated application of tacrolimus ointment (0.03-0.1%), and infants from age of 5 months treated with tacrolimus ointment (0.03%) had blood concentrations < 1.0 ng/ml. When observed, blood concentrations exceeding 1.0 ng/ml were transient. Systemic exposure increases with increasing treatment areas. However, both the extent and the rate of topical absorption of tacrolimus decrease as the skin heals. In both adults and children with an average of 50% body surface area treated, systemic exposure (i.e. AUC) of tacrolimus from Protopic is approximately 30-fold less than that seen with oral immunosuppressive doses in kidney and liver transplant patients. The lowest tacrolimus blood concentration at which systemic effects can be observed is not known. There was no evidence of systemic accumulation of tacrolimus in patients (adults and children) treated for prolonged periods (up to one year) with tacrolimus ointment.

Distribution

As systemic exposure is low with tacrolimus ointment, the high binding of tacrolimus (> 98.8%) to plasma proteins is considered not to be clinically relevant. Following topical application of tacrolimus ointment, tacrolimus is selectively delivered to the skin with minimal diffusion into the systemic circulation.

Metabolism

Metabolism of tacrolimus by human skin was not detectable. Systemically available tacrolimus is extensively metabolised in the liver via CYP3A4.

Elimination

When administered intravenously, tacrolimus has been shown to have a low clearance rate. The average total body clearance is approximately 2.25 l/h. The hepatic clearance of systemically available tacrolimus could be reduced in subjects with severe hepatic impairment, or in subjects who are co-treated with drugs that are potent inhibitors of CYP3A4. Following repeated topical application of the ointment the average half-life of tacrolimus was estimated to be 75 hours for adults and 65 hours for children.

Paediatric population

The pharmacokinetics of tacrolimus after topical application are similar to those reported in adults, with minimal systemic exposure and no evidence of accumulation (see above).

Preclinical Safety Data

Repeated dose toxicity and local tolerance

Repeated topical administration of tacrolimus ointment or the ointment vehicle to rats, rabbits and micropigs was associated with slight dermal changes such as erythema, oedema and papules. Long-term topical treatment of rats with tacrolimus led to systemic toxicity including alterations of kidneys, pancreas, eyes and nervous system. The changes were caused by high systemic exposure of rodents resulting from high transdermal absorption of tacrolimus. Slightly lower body weight gain in females was the only systemic change observed in micropigs at high ointment concentrations (3%). Rabbits were shown to be especially sensitive to intravenous administration of tacrolimus, reversible cardiotoxic effects being observed.

Mutagenicity

In vitro and *in vivo* tests did not indicate a genotoxic potential of tacrolimus.

Carcinogenicity

Systemic carcinogenicity studies in mice (18 months) and rats (24 months) revealed no carcinogenic potential of tacrolimus.

In a 24-month dermal carcinogenicity study performed in mice with 0.1% ointment, no skin tumours were observed. In the same study an increased incidence of lymphoma was detected in association with high systemic exposure.

In a photocarcinogenicity study, albino hairless mice were chronically treated with tacrolimus ointment and UV radiation. Animals treated with tacrolimus ointment showed a statistically significant reduction in time to skin tumour (squamous cell carcinoma) development and an increase in the number of tumours. It is unclear whether the effect of tacrolimus is due to systemic immunosuppression or a local effect. The risk for humans cannot be completely ruled out as the potential for local immunosuppression with the long-term use of tacrolimus ointment is unknown.

Reproduction toxicity

Embryo/foetal toxicity was observed in rats and rabbits, but only at doses that caused significant toxicity in maternal animals. Reduced sperm function was noted in male rats at high subcutaneous doses of tacrolimus.

PHARMACEUTICAL PARTICULARS

List of excipients

White soft paraffin

Liquid paraffin

Propylene carbonate

White beeswax
Hard paraffin
Butylhydroxytoluene (BHT)
All-rac- α -tocopherol

Incompatibilities

Not applicable

Shelf Life

Refer to Outer Carton

Special Precautions for Storage

Do not store above 25°C.

Nature and Contents of Container

Laminate tube with an inner lining of low-density-polyethylene fitted with a white polypropylene screw cap.

Package sizes: 10g, 30g

Instructions for Use and Handling and disposal

No special requirements.

LEO Pharma A/S

Industriparken 55, DK-2750 Ballerup, Denmark

DATE OF LAST REVISION OF THE TEXT

May 2019 (EU SmPC Sep 2018)