

## **1. NAME OF THE MEDICINAL PRODUCT**

Revlimid® 5 mg hard capsules  
Revlimid® 10 mg hard capsules  
Revlimid® 15 mg hard capsules  
Revlimid® 25 mg hard capsules

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each Revlimid® 5 mg hard capsule contains 5 mg of lenalidomide.  
Excipient: Each capsule contains 147 mg of anhydrous lactose.

Each Revlimid® 10 mg hard capsule contains 10 mg of lenalidomide.  
Excipient: Each capsule contains 294 mg of anhydrous lactose.

Each Revlimid® 15 mg hard capsule contains 15 mg of lenalidomide.  
Excipient: Each capsule contains 289 mg of anhydrous lactose.

Each Revlimid® 25 mg hard capsule contains 25 mg of lenalidomide.  
Excipient: Each capsule contains 200 mg of anhydrous lactose.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Hard capsule.

Revlimid® 5 mg hard capsules: White capsules, size 2, marked “REV 5 mg”.

Revlimid® 10 mg hard capsules: Blue-green/pale yellow capsules, size 0, marked “REV 10 mg”.

Revlimid® 15 mg hard capsules: Pale blue/white capsules, size 0, marked “REV 15 mg”.

Revlimid® 25 mg hard capsules: White capsules, size 0, marked “REV 25 mg”.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Revlimid® (lenalidomide) as monotherapy is indicated for the maintenance treatment of adult patients with newly diagnosed multiple myeloma who have undergone autologous stem cell transplantation.

Revlimid® (lenalidomide) in combination with dexamethasone is indicated for the treatment of previously untreated multiple myeloma patients who are not eligible for transplant.

Revlimid® (lenalidomide) in combination with dexamethasone is indicated for the treatment of multiple myeloma patients who have received at least one prior therapy.

### **4.2 Posology and method of administration**

Revlimid® (lenalidomide) should only be prescribed by Specialist Physician experienced in the management of malignancies, who have undergone the Revlimid® educational programme on Pregnancy Prevention Programme.

Treatment must be initiated and monitored under the supervision of physicians experienced in the management of multiple myeloma (MM).

For all indications described below:

- Dose is modified based upon clinical and laboratory findings (see section 4.4).
- Dose adjustments, during treatment and restart of treatment, are recommended to manage grade 3 or 4 thrombocytopenia, neutropenia, or other grade 3 or 4 toxicity judged to be related to lenalidomide.
- In case of neutropenia, the use of growth factors in patient management should be considered.
- If less than 12 hours has elapsed since missing a dose, the patient can take the dose. If more than 12 hours has elapsed since missing a dose at the normal time, the patient should not take the dose, but take the next dose at the normal time on the following day.

#### Posology

##### Newly diagnosed multiple myeloma (NDMM)

- Lenalidomide maintenance in patients who have undergone autologous stem cell transplantation (ASCT)

Lenalidomide maintenance should be initiated after adequate haematologic recovery following ASCT in patients without evidence of progression. Lenalidomide must not be started if the Absolute Neutrophil Count (ANC) is  $< 1.0 \times 10^9/L$ , and/or platelet counts are  $< 75 \times 10^9/L$ .

#### *Recommended dose*

The recommended starting dose is lenalidomide 10 mg orally once daily continuously (on days 1 to 28 of repeated 28-day cycles) given until disease progression or intolerance. After 3 cycles of lenalidomide maintenance, the dose can be increased to 15 mg orally once daily if tolerated.

- *Dose reduction steps*

	Starting dose (10 mg)	If dose increased (15 mg) <sup>a</sup>
Dose level -1	5 mg	10 mg
Dose level -2	5 mg (days 1-21 every 28 days)	5 mg
Dose level -3	Not applicable	5 mg (days 1-21 every 28 days)
	Do not dose below 5 mg (days 1-21 every 28 days)	

<sup>a</sup> After 3 cycles of lenalidomide maintenance, the dose can be increased to 15 mg orally once daily if tolerated.

- *Thrombocytopenia*

When platelets	Recommended course
Falls to $< 30 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 30 \times 10^9/L$	Resume lenalidomide at dose level -1 once daily
For each subsequent drop below $30 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 30 \times 10^9/L$	Resume lenalidomide at next lower dose level once daily

- *Absolute neutrophil count (ANC) - neutropenia*

When ANC	Recommended course <sup>a</sup>
Falls to $< 0.5 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 0.5 \times 10^9/L$	Resume lenalidomide at dose level -1 once daily
For each subsequent drop below $< 0.5 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 0.5 \times 10^9/L$	Resume lenalidomide at next lower dose level once daily

<sup>a</sup> At the physician's discretion, if neutropenia is the only toxicity at any dose level, add granulocyte colony stimulating factor (G-CSF) and maintain the dose level of lenalidomide.

- Lenalidomide in combination with dexamethasone until disease progression in patients who are not eligible for transplant

Lenalidomide treatment must not be started if the ANC is  $< 1.0 \times 10^9/L$ , and/or platelet counts are  $< 50 \times 10^9/L$ .

*Recommended dose*

The recommended starting dose of lenalidomide is 25 mg orally once daily on days 1 to 21 of repeated 28-day cycles.

The recommended dose of dexamethasone is 40 mg orally once daily on days 1, 8, 15 and 22 of repeated 28-day cycles. Patients may continue lenalidomide and dexamethasone therapy until disease progression or intolerance.

- *Dose reduction steps*

	Lenalidomide <sup>a</sup>	Dexamethasone <sup>a</sup>
Starting dose	25 mg	40 mg
Dose level-1	20 mg	20 mg
Dose level-2	15 mg	12 mg
Dose level-3	10 mg	8 mg
Dose level-4	5 mg	4 mg
Dose level-5	5mg every other day	Not applicable

<sup>a</sup> Dose reduction for both products can be managed independently

- *Thrombocytopenia*

When platelets	Recommended course
Falls to $< 25 \times 10^9/L$	Stop lenalidomide dosing for remainder of cycle <sup>a</sup>
Returns to $\geq 50 \times 10^9/L$	Decrease by one dose level when dosing resumed at next cycle

<sup>a</sup> If Dose limiting toxicity (DLT) occurs on  $> \text{day}15$  of a cycle, lenalidomide dosing will be interrupted for at least the remainder of the current 28-day cycle.

- *Absolute neutrophil count (ANC) - neutropenia*

When ANC	Recommended course <sup>a</sup>
First falls to $< 0.5 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 1 \times 10^9/L$ when neutropenia is the only observed toxicity	Resume lenalidomide at starting dose once daily
Returns to $\geq 0.5 \times 10^9/L$ when dose-dependent haematological toxicities other than neutropenia are observed	Resume lenalidomide at dose level -1 once daily

For each subsequent drop below $< 0.5 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 0.5 \times 10^9/L$	Resume lenalidomide at next lower dose level once daily.

<sup>a</sup> At the physician's discretion, if neutropenia is the only toxicity at any dose level, add granulocyte colony stimulating factor (G-CSF) and maintain the dose level of lenalidomide.

For hematologic toxicity the dose of lenalidomide may be re-introduced to the next higher dose level (up to the starting dose) upon improvement in bone marrow function (no hematologic toxicity for at least 2 consecutive cycles: ANC  $\geq 1.5 \times 10^9/L$  with a platelet count  $\geq 100 \times 10^9/L$  at the beginning of a new cycle).

*Multiple myeloma with at least one prior therapy*

Lenalidomide treatment must not be started if the ANC  $< 1.0 \times 10^9/L$ , and/or platelet counts  $< 75 \times 10^9/L$  or, dependent on bone marrow infiltration by plasma cells, platelet counts  $< 30 \times 10^9/L$ .

*Recommended dose*

The recommended starting dose of lenalidomide is 25 mg orally once daily on days 1 to 21 of repeated 28-day cycles. The recommended dose of dexamethasone is 40 mg orally once daily on days 1 to 4, 9 to 12, and 17 to 20 of each 28-day cycle for the first 4 cycles of therapy and then 40 mg once daily on days 1 to 4 every 28 days.

Prescribing physicians should carefully evaluate which dose of dexamethasone to use, taking into account the condition and disease status of the patient.

- Dose reduction steps*

Starting dose:	25 mg
Dose level -1:	15 mg
Dose level -2:	10 mg
Dose level -3:	5 mg

- Thrombocytopenia*

When platelets	Recommended course
First falls to $< 30 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 30 \times 10^9/L$	Resume lenalidomide at dose level -1
For each subsequent drop below $30 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 30 \times 10^9/L$	Resume lenalidomide at next lower dose level (dose level -2 or -3) once daily. Do not dose below 5 mg once daily.

- Absolute neutrophil count (ANC) - neutropenia*

When ANC	Recommended course <sup>a</sup>
First falls to $< 1.0 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 1.0 \times 10^9/L$ when neutropenia is	Resume lenalidomide at starting dose

When ANC	Recommended course
the only observed toxicity	once daily
Returns to $\geq 1.0 \times 10^9/\text{L}$ when dose-dependent haematological toxicities other than neutropenia are observed	Resume lenalidomide at dose level -1 once daily

For each subsequent drop below $< 1.0 \times 10^9/L$	Interrupt lenalidomide treatment
Returns to $\geq 1.0 \times 10^9/L$	Resume lenalidomide at next lower dose level (dose level -1, -2 or -3) once daily.
	Do not dose below 5 mg once daily.

<sup>a</sup> At the physician's discretion, if neutropenia is the only toxicity at any dose level, add granulocyte colony stimulating factor (G-CSF)

and maintain the dose level of lenalidomide.

#### All indications

For other Grade 3 or 4 toxicities judged to be related to lenalidomide, treatment should be stopped and only restarted at next lower dose level when toxicity has resolved to  $\leq$  Grade 2 depending on the physician's discretion.

Lenalidomide interruption or discontinuation should be considered for Grade 2 or 3 skin rash. Lenalidomide must be discontinued for angioedema, anaphylactic reaction, Grade 4 rash, exfoliative or bullous rash, or if Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) or Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) is suspected and should not be resumed following discontinuation from these reactions.

#### Special populations

- Paediatric population

Revlimid® should not be used in children and adolescents from birth to less than 18 years because of safety concerns (see section 5.1).

- Elderly

Currently available pharmacokinetic data are described in section 5.2. Lenalidomide has been used in clinical trials in multiple myeloma patients up to 91 years of age (see section 5.1).

Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection and it would be prudent to monitor renal function.

#### *Newly diagnosed multiple myeloma: patients who are not eligible for transplant*

Patients with newly diagnosed multiple myeloma aged 75 years and older should be carefully assessed before treatment is considered (see section 4.4).

For patients older than 75 years of age treated with lenalidomide in combination with dexamethasone, the starting dose of dexamethasone is 20 mg once daily on days 1, 8, 15 and 22 of each 28-day treatment cycle.

In patients with newly diagnosed multiple myeloma aged 75 years and older who received lenalidomide, there was a higher incidence of serious adverse reactions and adverse reactions that led to treatment discontinuation.

Lenalidomide combined therapy was less tolerated in newly diagnosed multiple myeloma patients older than 75 years of age compared to the younger population. These patients discontinued at a higher rate due to intolerance (Grade 3 or 4 adverse events and serious adverse events), when compared to patients  $< 75$  years.

#### *Multiple myeloma: patients with at least one prior therapy*

The percentage of multiple myeloma patients aged 65 or over was not significantly different between the lenalidomide/dexamethasone and placebo/dexamethasone groups. No overall difference in safety or efficacy was observed between these patients and younger patients, but greater pre-disposition of older individuals cannot be ruled out.

- Patients with renal impairment

Lenalidomide is primarily excreted by the kidney; patients with greater degrees of renal

impairment can have impaired treatment tolerance (see section 4.4). Care should be taken in dose selection and monitoring of renal function is advised.

No dose adjustments are required for patients with mild renal impairment.

The following dose adjustments are recommended at the start of therapy and throughout treatment for patients with moderate or severe impaired renal function or end stage renal disease.

There are no Phase 3 trial experiences with End Stage Renal Disease (ESRD) (CLCr < 30 mL/min, requiring dialysis).

#### *Multiple myeloma*

<b>Renal function (CLCr)</b>	<b>Dose adjustment</b> (days 1 to 21 of repeated 28- day cycles)
Moderate renal impairment ( $30 \leq \text{CLCr} < 50 \text{ mL/min}$ )	10 mg once daily <sup>1</sup>
Severe renal impairment (CLCr < 30 mL/min, not requiring dialysis)	15 mg every other day
End Stage Renal Disease (ESRD) (CLCr < 30 mL/min, requiring dialysis)	5 mg once daily. On dialysis days, the dose should be administered following dialysis.

<sup>1</sup> The dose may be escalated to 15 mg once daily after 2 cycles if patient is not responding to treatment and is tolerating the treatment.

After initiation of lenalidomide therapy, subsequent lenalidomide dose modification in renally impaired patients should be based on individual patient treatment tolerance, as described above.

- Patients with hepatic impairment

Lenalidomide has not formally been studied in patients with impaired hepatic function and there are no specific dose recommendations.

#### Method of administration

Oral use.

Revlimid® capsules should be taken orally at about the same time on the scheduled days. The capsules should not be opened, broken or chewed. The capsules should be swallowed whole, preferably with water, either with or without food.

It is recommended to press only on one end of the capsule to remove it from the blister thereby reducing the risk of capsule deformation or breakage.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Women who are pregnant.
- Women of childbearing potential unless all of the conditions of the Pregnancy Prevention Programme are met (see sections 4.4 and 4.6).

### **4.4 Special warnings and precautions for use**

#### Pregnancy warning

Lenalidomide is structurally related to thalidomide. Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects. Lenalidomide induced in monkeys' malformations similar to those described with thalidomide (see sections 4.6 and 5.3). If lenalidomide is taken during pregnancy, a teratogenic effect of lenalidomide in humans is expected.

The conditions of the Pregnancy Prevention Programme must be fulfilled for all patients unless there is reliable evidence that the patient does not have childbearing potential.

#### Criteria for women of non-childbearing potential

A female patient or a female partner of a male patient is considered to have childbearing potential unless she meets at least one of the following criteria:

- Age  $\geq$  50 years and naturally amenorrhoeic for  $\geq$  1 year (Amenorrhoea following cancer therapy or during breast-feeding does not rule out childbearing potential).
- Premature ovarian failure confirmed by a specialist gynaecologist
- Previous bilateral salpingo-oophorectomy, or hysterectomy
- XY genotype, Turner syndrome, uterine agenesis.

### Counselling

For women of childbearing potential, lenalidomide is contraindicated unless all of the following are met:

- She understands the expected teratogenic risk to the unborn child
- She understands the need for effective contraception, without interruption, at least 4 weeks before starting treatment, throughout the entire duration of treatment, and at least 4 weeks after the end of treatment
- Even if a woman of childbearing potential has amenorrhea she must follow all the advice on effective contraception
- She should be capable of complying with effective contraceptive measures
- She is informed and understands the potential consequences of pregnancy and the need to rapidly consult if there is a risk of pregnancy
- She understands the need to commence the treatment as soon as lenalidomide is dispensed following a negative pregnancy test
- She understands the need and accepts to undergo pregnancy testing at least every 4 weeks except in case of confirmed tubal sterilisation
- She acknowledges that she understands the hazards and necessary precautions associated with the use of lenalidomide.

For male patients taking lenalidomide, pharmacokinetic data has demonstrated that lenalidomide is present in human semen at extremely low levels during treatment and is undetectable in human semen 3 days after stopping the substance in the healthy subject (see section 5.2). As a precaution and taking into account special populations with prolonged elimination time such as renal impairment, all male patients taking lenalidomide must meet the following conditions:

- Understand the expected teratogenic risk if engaged in sexual activity with a pregnant woman or a woman of childbearing potential
- Understand the need for the use of a condom if engaged in sexual activity with a pregnant woman or a woman of childbearing potential not using effective contraception (even if the man has had a vasectomy), throughout treatment duration, during dose interruption and for at least 7 days after dose interruptions and/or cessation of treatment.
- Understand that if his female partner becomes pregnant whilst he is taking Revlimid® or within 7 days after he has stopped taking Revlimid®, he should inform his treating physician immediately and that it is recommended to refer the female partner to a physician specialised or experienced in teratology for evaluation and advice.

The prescriber must ensure that for women of childbearing potential:

- The patient complies with the conditions of the Pregnancy Prevention Programme, including confirmation that she has an adequate level of understanding.
- The patient has acknowledged the aforementioned conditions.

### Contraception

Women of childbearing potential must use at least one effective method of contraception for at least 4 weeks before therapy, during therapy, and until at least 4 weeks after lenalidomide therapy and even in case of dose interruption unless the patient commits to absolute and continuous abstinence confirmed on a monthly basis. If not established on effective contraception, the patient must be referred to an appropriately trained health care professional for contraceptive advice in order that contraception can be initiated.

The following can be considered to be examples of suitable methods of contraception:

- Implant



- Levonorgestrel-releasing intrauterine system (IUS)
- Medroxyprogesterone acetate depot
- Tubal sterilisation
- Sexual intercourse with a vasectomised male partner only; vasectomy must be confirmed by two negative semen analyses
- Ovulation inhibitory progesterone-only pills (i.e. desogestrel)

Because of the increased risk of venous thromboembolism in patients with multiple myeloma taking lenalidomide and dexamethasone, and to a lesser extent in patients with multiple myeloma taking lenalidomide monotherapy, combined oral contraceptive pills are not recommended (see also section 4.5). If a patient is currently using combined oral contraception the patient should switch to one of the effective methods listed above. The risk of venous thromboembolism continues for 4–6 weeks after discontinuing combined oral contraception. The efficacy of contraceptive steroids may be reduced during co-treatment with dexamethasone (see section 4.5).

Implants and levonorgestrel-releasing intrauterine systems are associated with an increased risk of infection at the time of insertion and irregular vaginal bleeding. Prophylactic antibiotics should be considered particularly in patients with neutropenia.

Copper-releasing intrauterine devices are generally not recommended due to the potential risks of infection at the time of insertion and menstrual blood loss which may compromise patients with neutropenia or thrombocytopenia.

#### Pregnancy testing

According to local practice, medically supervised pregnancy tests with a minimum sensitivity of 25 mIU/mL must be performed for women of childbearing potential as outlined below. This requirement includes women of childbearing potential who practice absolute and continuous abstinence. Ideally, pregnancy testing, issuing a prescription and dispensing should occur on the same day. Dispensing of lenalidomide to women of childbearing potential should occur within 7 days of the prescription.

#### Prior to starting treatment

A medically supervised pregnancy test should be performed during the consultation, when lenalidomide is prescribed, or in the 3 days prior to the visit to the prescriber once the patient had been using effective contraception for at least 4 weeks. The test should ensure the patient is not pregnant when she starts treatment with lenalidomide.

#### Follow-up and end of treatment

A medically supervised pregnancy test should be repeated at least every 4 weeks, including at least 4 weeks after the end of treatment, except in the case of confirmed tubal sterilisation. These pregnancy tests should be performed on the day of the prescribing visit or in the 3 days prior to the visit to the prescriber.

#### Men

Lenalidomide is present in human semen during treatment. As a precaution, and taking into account special populations with potentially prolonged elimination time such as renal impairment, all male patients taking lenalidomide, including those who have had a vasectomy, should use condoms throughout treatment duration, during dose interruption and for at least 7 days after cessation of treatment if their partner is pregnant or of childbearing potential and has no contraception.

Male patients should not donate semen or sperm during treatment (including during dose interruptions) and for at least 7 days following discontinuation of lenalidomide.

#### Additional precautions

Patients should be instructed never to give this medicinal product to another person and to return any unused capsules to their pharmacist at the end of treatment for safe disposal.

Patients should not donate blood, semen or sperm during treatment (including during dose interruptions) and for at least 7 days following discontinuation of lenalidomide.

Healthcare professionals and caregivers should wear disposable gloves when handling the blister or capsule. Women who are pregnant or suspect they may be pregnant should not handle the blister or capsule (see section 6.5).

#### Educational materials, prescribing and dispensing restrictions

In order to assist patients in avoiding foetal exposure to lenalidomide, the marketing authorization holder will provide educational material to health care professionals to reinforce the warnings about the expected teratogenicity of lenalidomide, to provide advice on contraception before therapy is started, and to provide guidance on the need for pregnancy testing. The prescriber must inform male and female patients about the expected teratogenic risk and the strict pregnancy prevention measures as specified in the Pregnancy Prevention Programme and provide patients with appropriate patient educational brochure, patient card and/or equivalent tool. Ideally, pregnancy testing, issuing a prescription and dispensing should occur on the same day. Dispensing of lenalidomide to women of childbearing potential should occur within 7 days of the prescription and following a medically supervised negative pregnancy test result. Prescriptions for women of childbearing potential can be for a maximum duration of treatment of 4 weeks according to the approved indications dosing regimens (see section 4.2), and prescriptions for all other patients can be for a maximum duration of treatment of 12 weeks.

#### Other special warnings and precautions for use

##### *Myocardial infarction*

Myocardial infarction has been reported in patients receiving lenalidomide, particularly in those with known risk factors and within the first 12 months when used in combination with dexamethasone. Patients with known risk factors – including prior thrombosis – should be closely monitored, and action should be taken to try to minimize all modifiable risk factors (e.g. smoking, hypertension, and hyperlipidaemia).

##### *Venous and arterial thromboembolic events*

In patients with multiple myeloma, the combination of lenalidomide with dexamethasone is associated with an increased risk of venous thromboembolism (predominantly deep vein thrombosis and pulmonary embolism).

In patients with multiple myeloma, treatment with lenalidomide monotherapy was associated with a lower risk of venous thromboembolism (predominantly deep vein thrombosis and pulmonary embolism) than in patients with multiple myeloma treated with lenalidomide in combination therapy (see sections 4.5 and 4.8).

In patients with multiple myeloma, the combination of lenalidomide with dexamethasone is associated with an increased risk of arterial thromboembolism (predominantly myocardial infarction and cerebrovascular event). The risk of arterial thromboembolism is lower in patients with multiple myeloma treated with lenalidomide monotherapy than in patients with multiple myeloma treated with lenalidomide in combination therapy.

Consequently, patients with known risk factors for thromboembolism – including prior thrombosis – should be closely monitored. Action should be taken to try to minimize all modifiable risk factors (e.g. smoking, hypertension, and hyperlipidaemia). Concomitant administration of erythropoietic agents or previous history of thromboembolic events may also increase thrombotic risk in these patients. Therefore, erythropoietic agents, or other agents that may increase the risk of thrombosis, such as hormone replacement therapy, should be used with caution in multiple myeloma patients receiving lenalidomide with dexamethasone. A haemoglobin concentration above 12 g/dl should lead to discontinuation of erythropoietic agents.

Patients and physicians are advised to be observant for the signs and symptoms of thromboembolism. Patients should be instructed to seek medical care if they develop symptoms such as shortness of breath, chest pain, arm or leg swelling. Prophylactic antithrombotic medicines should be recommended, especially in patients with additional thrombotic risk factors. The decision to take antithrombotic prophylactic measures should be made after careful assessment of an individual

patient's underlying risk factors.

If the patient experiences any thromboembolic events, treatment must be discontinued and standard anticoagulation therapy started. Once the patient has been stabilised on the anticoagulation treatment and any complications of the thromboembolic event have been managed, the lenalidomide treatment may be restarted at the original dose dependent upon a benefit risk assessment. The patient should continue anticoagulation therapy during the course of lenalidomide treatment.

### Pulmonary hypertension

Cases of pulmonary hypertension, some fatal, have been reported in patients treated with lenalidomide. Patients should be evaluated for signs and symptoms of underlying cardiopulmonary disease prior to initiating and during lenalidomide therapy.

### Neutropenia and thrombocytopenia

The major dose limiting toxicities of lenalidomide include neutropenia and thrombocytopenia. A complete blood cell count, including white blood cell count with differential count, platelet count, haemoglobin, and haematocrit should be performed at baseline, every week for the first 8 weeks of lenalidomide treatment and monthly thereafter to monitor for cytopenias. A dose interruption and/or a dose reduction may be required (see section 4.2).

In case of neutropenia, the physician should consider the use of growth factors in patient management. Patients should be advised to promptly report febrile episodes.

Patients and physicians are advised to be observant for signs and symptoms of bleeding, including petechiae and epistaxis, especially in patients receiving concomitant medicinal products susceptible to induce bleeding (see section 4.8, Haemorrhagic disorders).

Co-administration of lenalidomide with other myelosuppressive agents should be undertaken with caution.

- Newly diagnosed multiple myeloma: patients who have undergone ASCT treated with lenalidomide maintenance

The adverse reactions from CALGB 100104 included events reported post-high dose melphalan and ASCT (HDM/ASCT) as well as events from the maintenance treatment period. A second analysis identified events that occurred after the start of maintenance treatment. In IFM 2005-02, the adverse reactions were from the maintenance treatment period only.

Overall, Grade 4 neutropenia was observed at a higher frequency in the lenalidomide maintenance arms compared to the placebo maintenance arms in the 2 studies evaluating lenalidomide maintenance in NDMM patients who have undergone ASCT (32.1% vs 26.7% [16.1% vs 1.8% after the start of maintenance treatment] in CALGB 100104 and 16.4% vs 0.7% in IFM 2005-02, respectively). Treatment-emergent AEs of neutropenia leading to lenalidomide discontinuation were reported in 2.2% of patients in CALGB 100104 and 2.4% of patients in IFM 2005-02, respectively. Grade 4 febrile neutropenia was reported at similar frequencies in the lenalidomide maintenance arms compared to placebo maintenance arms in both studies (0.4% vs 0.5% [0.4% vs 0.5% after the start of maintenance treatment] in CALGB 100104 and 0.3% vs 0% in IFM 2005-02, respectively). Patients should be advised to promptly report febrile episodes, a treatment interruption and/or dose reductions may be required (see section 4.2).

Grade 3 or 4 thrombocytopenia was observed at a higher frequency in the lenalidomide maintenance arms compared to the placebo maintenance arms in studies evaluating lenalidomide maintenance in NDMM patients who have undergone ASCT (37.5% vs 30.3% [17.9% vs 4.1% after the start of maintenance treatment] in CALGB 100104 and 13.0% vs 2.9% in IFM 2005-02, respectively). Patients and physicians are advised to be observant for signs and symptoms of bleeding, including petechiae and epistaxes, especially in patients receiving concomitant medicinal products susceptible to induce bleeding (see section 4.8, Haemorrhagic disorders).

- Newly diagnosed multiple myeloma: patients who are not eligible for transplant treated with lenalidomide in combination with low dose dexamethasone

Grade 4 neutropenia was observed in the lenalidomide arms in combination with low dose dexamethasone to a lesser extent than in the comparator arm (8.5% in the Rd [continuous treatment] and Rd18 [treatment for 18 four-week cycles] compared with 15% in the melphalan/ prednisone/ thalidomide arm, see section 4.8). Grade 4 febrile neutropenia episodes were consistent with the comparator arm (0.6% in the Rd and Rd18 lenalidomide/dexamethasone-treated patients compared with 0.7% in the melphalan/prednisone/thalidomide arm, see section 4.8).

Grade 3 or 4 thrombocytopenia was observed to a lesser extent in the Rd and Rd18 arms than in the comparator arm (8.1% vs 11.1%, respectively).

- Multiple myeloma: patients with at least one prior therapy

The combination of lenalidomide with dexamethasone in multiple myeloma patients with at least one prior therapy is associated with a higher incidence of Grade 4 neutropenia (5.1% in lenalidomide/dexamethasone-treated patients compared with 0.6% in placebo/dexamethasone-treated patients; see section 4.8). Grade 4 febrile neutropenia episodes were observed infrequently (0.6% in lenalidomide/dexamethasone-treated patients compared to 0.0% in placebo/dexamethasone-treated patients; see section 4.8).

The combination of lenalidomide with dexamethasone in multiple myeloma patients is associated with a higher incidence of Grade 3 and Grade 4 thrombocytopenia (9.9% and 1.4%, respectively, in lenalidomide/dexamethasone-treated patients compared to 2.3% and 0.0% in placebo/dexamethasone-treated patients; see section 4.8).

#### Thyroid disorders

Cases of hypothyroidism and cases of hyperthyroidism have been reported. Optimal control of co-morbid conditions influencing thyroid function is recommended before start of treatment. Baseline and ongoing monitoring of thyroid function is recommended.

#### Peripheral neuropathy

Lenalidomide is structurally related to thalidomide, which is known to induce severe peripheral neuropathy. There was no increase in peripheral neuropathy observed with lenalidomide in combination with dexamethasone or lenalidomide monotherapy or with long term use of lenalidomide for the treatment of newly diagnosed multiple myeloma.

#### Tumour flare reaction and tumour lysis syndrome

Because lenalidomide has anti-neoplastic activity the complications of tumour lysis syndrome (TLS) may occur. Cases of TLS and tumour flare reaction (TFR), including fatal cases, have been reported (see section 4.8). The patients at risk of TLS and TFR are those with high tumour burden prior to treatment. Caution should be practiced when introducing these patients to lenalidomide. These patients should be monitored closely, especially during the first cycle or dose-escalation, and appropriate precautions taken.

#### Allergic reactions and severe skin reactions

Cases of allergic reactions including angioedema, anaphylactic reaction and severe cutaneous reactions including SJS, TEN and DRESS have been reported in patients treated with lenalidomide (see section 4.8). Patients should be advised of the signs and symptoms of these reactions by their prescribers and should be told to seek medical attention immediately if they develop these symptoms. Lenalidomide must be discontinued for angioedema, anaphylactic reaction, exfoliative or bullous rash, or if SJS, TEN or DRESS is suspected, and should not be resumed following discontinuation for these reactions. Interruption or discontinuation of lenalidomide should be considered for other forms of skin reaction depending on severity. Patients who had previous allergic reactions while treated with thalidomide should be monitored closely, as a possible cross-reaction between lenalidomide and thalidomide has been reported in the literature. Patients with a history of severe rash associated with thalidomide treatment should not receive lenalidomide.

#### Lactose intolerance

Revlimid® capsules contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

### Second primary malignancies

An increase of second primary malignancies (SPM) has been observed in clinical trials in previously treated myeloma patients receiving lenalidomide/dexamethasone (3.98 per 100 person-years) compared to controls (1.38 per 100 person-years). Non-invasive SPM comprise basal cell or squamous cell skin cancers. Most of the invasive SPMs were solid tumour malignancies.

In clinical trials of newly diagnosed multiple myeloma patients not eligible for transplant, a 4.9-fold increase in incidence rate of hematologic SPM (cases of AML, MDS) has been observed in patients receiving lenalidomide in combination with melphalan and prednisone until progression (1.75 per 100 person-years) compared with melphalan in combination with prednisone (0.36 per 100 person-years).

A 2.12-fold increase in incidence rate of solid tumour SPM has been observed in patients receiving lenalidomide (9 cycles) in combination with melphalan and prednisone (1.57 per 100 person-years) compared with melphalan in combination with prednisone (0.74 per 100 person-years).

In patients receiving lenalidomide in combination with dexamethasone until progression or for 18 months, the hematologic SPM incidence rate (0.16 per 100 person-years) was not increased as compared to thalidomide in combination with melphalan and prednisone (0.79 per 100 person-years).

A 1.3-fold increase in incidence rate of solid tumour SPM has been observed in patients receiving lenalidomide in combination with dexamethasone until progression or for 18 months (1.58 per 100 person-years) compared to thalidomide in combination with melphalan and prednisone (1.19 per 100 person-years).

The increased risk of secondary primary malignancies associated with lenalidomide is relevant also in the context of NDMM after stem cell transplantation. Though this risk is not yet fully characterized, it should be kept in mind when considering and using Revlimid® in this setting.

The incidence rate of hematologic malignancies, most notably AML, MDS and B-cell malignancies (including Hodgkin's lymphoma), was 1.31 per 100 person-years for the lenalidomide arms and 0.58 per 100 person-years for the placebo arms (1.02 per 100 person-years for patients exposed to lenalidomide after ASCT and 0.60 per 100 person-years for patients not-exposed to lenalidomide after ASCT). The incidence rate of solid tumour SPMs was 1.36 per 100 person-years for the lenalidomide arms and 1.05 per 100 person-years for the placebo arms (1.26 per 100 person-years for patients exposed to lenalidomide after ASCT and 0.60 per 100 person-years for patients not-exposed to lenalidomide after ASCT).

The risk of occurrence of hematologic SPM must be taken into account before initiating treatment with lenalidomide. Physicians should carefully evaluate patients before and during treatment using standard cancer screening for occurrence of SPM and institute treatment as indicated.

### Hepatic disorders

Hepatic failure, including fatal cases, has been reported in patients treated with lenalidomide in combination therapy: acute hepatic failure, toxic hepatitis, cytolytic hepatitis, cholestatic hepatitis, and mixed cytolytic/cholestatic hepatitis have been reported. The mechanisms of severe drug-induced hepatotoxicity remain unknown although, in some cases, pre-existing viral liver disease, elevated baseline liver enzymes, and possibly treatment with antibiotics might be risk factors.

Abnormal liver function tests were commonly reported and were generally asymptomatic and reversible upon dosing interruption. Once parameters have returned to baseline, treatment at a lower dose may be considered.

Lenalidomide is excreted by the kidneys. It is important to dose adjust patients with renal impairment in order to avoid plasma levels which may increase the risk for higher haematological adverse reactions

or hepatotoxicity. Monitoring of liver function is recommended, particularly when there is a history of or concurrent viral liver infection or when lenalidomide is combined with medicinal products known to be associated with liver dysfunction.

#### Infection with or without neutropenia

Patients with multiple myeloma are prone to develop infections including pneumonia. A higher rate of infections was observed with lenalidomide in combination with dexamethasone than with MPT in patients with NDMM who are not eligible for transplant, and with lenalidomide maintenance compared to placebo in patients with NDMM who had undergone ASCT. Grade  $\geq 3$  infections occurred within the context of neutropenia in less than one-third of the patients. Patients with known risk factors for infections should be closely monitored. All patients should be advised to seek medical attention promptly at the first sign of infection (eg, cough, fever, etc) thereby allowing for early management to reduce severity.

#### Viral reactivation

Cases of viral reactivation have been reported in patients receiving lenalidomide, including serious cases of herpes zoster or hepatitis B virus (HBV) reactivation.

Some of the cases of viral reactivation had a fatal outcome.

Some of the cases of herpes zoster reactivation resulted in disseminated herpes zoster, meningitis herpes zoster or ophthalmic herpes zoster requiring a temporary hold or permanent discontinuation of the treatment with lenalidomide and adequate antiviral treatment.

Reactivation of hepatitis B has been reported rarely in patients receiving lenalidomide who have previously been infected with the hepatitis B virus. Some of these cases have progressed to acute hepatic failure resulting in discontinuation of lenalidomide and adequate antiviral treatment. Hepatitis B virus status should be established before initiating treatment with lenalidomide. For patients who test positive for HBV infection, consultation with a physician with expertise in the treatment of hepatitis B is recommended. Caution should be exercised when lenalidomide is used in patients previously infected with HBV, including patients who are anti-HBc positive but HBsAg negative. These patients should be closely monitored for signs and symptoms of active HBV infection throughout therapy.

#### Progressive multifocal leukoencephalopathy

Cases of progressive multifocal leukoencephalopathy (PML), including fatal cases, have been reported with lenalidomide. PML was reported several months to several years after starting the treatment with lenalidomide. Cases have generally been reported in patients taking concomitant dexamethasone or prior treatment with other immunosuppressive chemotherapy. Physicians should monitor patients at regular intervals and should consider PML in the differential diagnosis in patients with new or worsening neurological symptoms, cognitive or behavioural signs or symptoms. Patients should also be advised to inform their partner or caregivers about their treatment, since they may notice symptoms that the patient is not aware of.

The evaluation for PML should be based on neurological examination, magnetic resonance imaging of the brain, and cerebrospinal fluid analysis for JC virus (JCV) DNA by polymerase chain reaction (PCR) or a brain biopsy with testing for JCV. A negative JCV PCR does not exclude PML. Additional follow-up and evaluation may be warranted if no alternative diagnosis can be established.

If PML is suspected, further dosing must be suspended until PML has been excluded. If PML is confirmed, lenalidomide must be permanently discontinued.

#### Newly diagnosed multiple myeloma patients

There was a higher rate of intolerance (Grade 3 or 4 adverse events, serious adverse events, discontinuation) in patients with age  $> 75$  years, ISS stage III, ECOG PS  $\geq 2$  or CLcr  $< 60$  mL/min when lenalidomide is given in combination. Patients should be carefully assessed for their ability to tolerate lenalidomide in combination, with consideration to age, ISS stage III, ECOG PS  $\geq 2$  or CLcr  $< 60$  mL/min (see sections 4.2 and 4.8).

#### Cataract

Cataract has been reported with a higher frequency in patients receiving lenalidomide in combination with dexamethasone particularly when used for a prolonged time. Regular monitoring of visual ability is recommended.

#### Solid Organ Transplant Rejection

Cases of solid organ transplant (SOT) rejection have been reported in the post-market setting with the use of Revlimid® and, in some cases, have resulted in a fatal outcome. Onset may be acute, occurring within 1 to 3 cycles of Revlimid® treatment. Potential contributing factors for SOT rejection in the reported cases include underlying disease (e.g., amyloidosis), concurrent infections and recent discontinuation or reduction of immunosuppressive therapy. The incidence rate of SOT rejection cannot be reliably estimated due to the limitation of post-marketing safety data and that patients with SOT were generally excluded from Celgene-sponsored Revlimid® clinical trials. The benefit of treatment with Revlimid® versus the risk of possible SOT rejection should be considered in patients with a history of SOT before initiating Revlimid® therapy. Clinical and laboratory signs of SOT rejection should be closely monitored and Revlimid® therapy should be discontinued in the event of SOT rejection.

### **4.5 Interaction with other medicinal products and other forms of interaction**

Erythropoietic agents, or other agents that may increase the risk of thrombosis, such as hormone replacement therapy, should be used with caution in multiple myeloma patients receiving lenalidomide with dexamethasone (see sections 4.4 and 4.8).

#### Oral contraceptives

No interaction study has been performed with oral contraceptives. Lenalidomide is not an enzyme inducer. In an *in vitro* study with human hepatocytes, lenalidomide, at various concentrations tested did not induce CYP1A2, CYP2B6, CYP2C9, CYP2C19 and CYP3A4/5. Therefore, induction leading to reduced efficacy of medicinal products, including hormonal contraceptives, is not expected if lenalidomide is administered alone. However, dexamethasone is known to be a weak to moderate inducer of CYP3A4 and is likely to also affect other enzymes as well as transporters. It may not be excluded that the efficacy of oral contraceptives may be reduced during treatment. Effective measures to avoid pregnancy must be taken (see sections 4.4 and 4.6).

#### Warfarin

Co-administration of multiple 10 mg doses of lenalidomide had no effect on the single dose pharmacokinetics of R- and S- warfarin. Co-administration of a single 25 mg dose of warfarin had no effect on the pharmacokinetics of lenalidomide. However, it is not known whether there is an interaction during clinical use (concomitant treatment with dexamethasone). Dexamethasone is a weak to moderate enzyme inducer and its effect on warfarin is unknown. Close monitoring of warfarin concentration is advised during the treatment.

#### Digoxin

Concomitant administration with lenalidomide 10 mg once daily increased the plasma exposure of digoxin (0.5 mg, single dose) by 14% with a 90% CI (confidence interval) [0.52%-28.2%]. It is not known whether the effect will be different in the clinical use (higher lenalidomide doses and concomitant treatment with dexamethasone). Therefore, monitoring of the digoxin concentration is advised during lenalidomide treatment.

#### Statins

There is an increased risk of rhabdomyolysis when statins are administered with lenalidomide, which may be simply additive. Enhanced clinical and laboratory monitoring is warranted notably during the first weeks of treatment.

#### Dexamethasone

Co-administration of single or multiple doses of dexamethasone (40 mg once daily) has no clinically relevant effect on the multiple dose pharmacokinetics of lenalidomide (25 mg once daily).

#### Interactions with P-glycoprotein (P-gp) inhibitors

*In vitro*, lenalidomide is a substrate of P-gp, but is not a P-gp inhibitor. Co-administration of multiple doses of the strong P-gp inhibitor quinidine (600 mg, twice daily) or the moderate P-gp inhibitor/substrate temsirolimus (25 mg) has no clinically relevant effect on the pharmacokinetics of lenalidomide (25 mg). Co-administration of lenalidomide does not alter the pharmacokinetics of temsirolimus.

#### **4.6 Fertility, pregnancy and lactation**

Due to the teratogenic potential, lenalidomide must be prescribed under a Pregnancy Prevention Programme (see section 4.4) unless there is reliable evidence that the patient does not have childbearing potential.

##### Women of childbearing potential / Contraception in males and females

Women of childbearing potential should use at least one effective method of contraception. If pregnancy occurs in a woman treated with lenalidomide, treatment must be stopped and the patient should be referred to a physician specialised or experienced in teratology for evaluation and advice. If pregnancy occurs in a partner of a male patient taking lenalidomide, it is recommended to refer the female partner to a physician specialised or experienced in teratology for evaluation and advice.

Lenalidomide is present in human semen at extremely low levels during treatment and is undetectable in human semen 3 days after stopping the substance in the healthy subject (see section 5.2). As a precaution and taking into account special populations with prolonged elimination time such as renal impairment, all male patients taking lenalidomide should use condoms throughout treatment duration, during dose interruption and for at least 7 days after cessation of treatment if their partner is pregnant or of childbearing potential and has no contraception.

##### Pregnancy

Lenalidomide is structurally related to thalidomide. Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects.

Lenalidomide induced malformation in monkeys similar to those described with thalidomide (see section 5.3). Therefore, a teratogenic effect of lenalidomide is expected and lenalidomide is contraindicated during pregnancy (see section 4.3).

##### Breast-feeding

It is not known whether lenalidomide is excreted in human milk. Therefore, breast-feeding should be discontinued during therapy with lenalidomide.

##### Fertility

A fertility study in rats with lenalidomide doses up to 500 mg/kg (approximately 200 to 500 times the human doses of 25 mg and 10 mg, respectively, based on body surface area) produced no adverse effects on fertility and no parental toxicity.

#### **4.7 Effects on ability to drive and use machines**

Lenalidomide has minor or moderate influence on the ability to drive and use machines. Fatigue, dizziness, somnolence, vertigo and blurred vision have been reported with the use of lenalidomide. Therefore, caution is recommended when driving or operating machines.

#### **4.8 Undesirable effects**

##### Summary of the safety profile

##### *Newly diagnosed multiple myeloma: patients who have undergone ASCT treated with lenalidomide maintenance*

A conservative approach was applied to determine the adverse reactions from CALGB 100104. The adverse reactions described in Table 1 included events reported post-HDM/ASCT as well as events from the maintenance treatment period. A second analysis that identified events that occurred after the start of maintenance treatment suggests that the frequencies described in Table 1 may be higher than actually



observed during the maintenance treatment period. In IFM 2005-02, the adverse reactions were from the maintenance treatment period only.

The serious adverse reactions observed more frequently ( $\geq 5\%$ ) with lenalidomide maintenance than placebo were:

- Pneumonia (10.6%; combined term) from IFM 2005-02
- Lung infection (9.4% [9.4% after the start of maintenance treatment]) from CALGB 100104

In the IFM 2005-02 study, the adverse reactions observed more frequently with lenalidomide maintenance than placebo were neutropenia (60.8%), bronchitis (47.4%), diarrhoea (38.9%), nasopharyngitis (34.8%), muscle spasms (33.4%), leucopenia (31.7%), asthenia (29.7%), cough (27.3%), thrombocytopenia (23.5%), gastroenteritis (22.5%) and pyrexia (20.5%).

In the CALGB 100104 study, the adverse reactions observed more frequently with lenalidomide maintenance than placebo were neutropenia (79.0% [71.9% after the start of maintenance treatment]), thrombocytopenia (72.3% [61.6%]), diarrhoea (54.5% [46.4%]), rash (31.7% [25.0%]), upper respiratory tract infection (26.8% [26.8%]), fatigue (22.8% [17.9%]), leucopenia (22.8% [18.8%]) and anaemia (21.0% [13.8%]).

*Newly diagnosed multiple myeloma: patients who are not eligible for transplant treated with lenalidomide in combination with low dose dexamethasone*

The serious adverse reactions observed more frequently ( $\geq 5\%$ ) with lenalidomide in combination with low dose dexamethasone (Rd and Rd18) than with melphalan, prednisone and thalidomide (MPT) were:

- Pneumonia (9.8%)
- Renal failure (including acute) (6.3%)

The adverse reactions observed more frequently with Rd or Rd18 than MPT were: diarrhoea (45.5%), fatigue (32.8%), back pain (32.0%), asthenia (28.2%), insomnia (27.6%), rash (24.3%), decreased appetite (23.1%), cough (22.7%), pyrexia (21.4%), and muscle spasms (20.5%).

*Multiple myeloma: patients with at least one prior therapy*

In two Phase 3 placebo-controlled studies, 353 patients with multiple myeloma were exposed to the lenalidomide/dexamethasone combination and 351 to the placebo/dexamethasone combination.

The most serious adverse reactions observed more frequently in lenalidomide/ dexamethasone than placebo/ dexamethasone combination were:

- Venous thromboembolism (deep vein thrombosis, pulmonary embolism) (see section 4.4)
- Grade 4 neutropenia (see section 4.4).

The observed adverse reactions which occurred more frequently with lenalidomide and dexamethasone than placebo and dexamethasone in pooled multiple myeloma clinical trials (MM-009 and MM-010) were fatigue (43.9%), neutropenia (42.2%), constipation (40.5%), diarrhoea (38.5%), muscle cramp (33.4%), anaemia (31.4%), thrombocytopenia (21.5%), and rash (21.2%).

*Tabulated list of adverse reactions*

The adverse reactions observed in patients treated with lenalidomide are listed below by system organ class and frequency. Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. Frequencies are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data).

Adverse reactions have been included under the appropriate category in the table below according to the highest frequency observed in any of the main clinical trials.

*Tabulated summary for monotherapy in MM*

The following table is derived from data gathered during NDMM studies in patients who have undergone ASCT treated with lenalidomide maintenance. The data were not adjusted according to the longer duration

of treatment in the lenalidomide-containing arms continued until disease progression versus the placebo arms in the pivotal multiple myeloma studies (see section 5.1).

**Table 1. ADRs reported in clinical trials in patients with multiple myeloma treated with lenalidomide maintenance therapy**

<b>System Organ Class/Preferred Term</b>	<b>All ADRs/Frequency</b>	<b>Grade 3-4 ADRs/Frequency</b>
<b>Infections and Infestations</b>	<p><u>Very Common</u> Pneumonia<sup>◊, a</sup>, Upper respiratory tract infection, Neutropenic infection, Bronchitis<sup>◊</sup>, Influenza<sup>◊</sup>, Gastroenteritis<sup>◊</sup>, Sinusitis, Nasopharyngitis, Rhinitis</p> <p><u>Common</u> Infection<sup>◊</sup>, Urinary tract infection<sup>◊, *</sup>, Lower respiratory tract infection, Lung infection<sup>◊</sup></p>	<p><u>Very Common</u> Pneumonias<sup>◊, a</sup>, Neutropenic infection</p> <p><u>Common</u> Sepsis<sup>◊, b</sup>, Bacteraemia, Lung infection<sup>◊</sup>, Lower respiratory tract infection bacterial, Bronchitis<sup>◊</sup>, Influenza<sup>◊</sup>, Gastroenteritis<sup>◊</sup>, Herpes zoster<sup>◊</sup>, Infection<sup>◊</sup></p>
<b>Neoplasms Benign, Malignant and Unspecified (incl cysts and polyps)</b>	<p><u>Common</u> Myelodysplastic syndrome<sup>◊, *</sup></p>	
<b>Blood and Lymphatic System Disorders</b>	<p><u>Very Common</u> Neutropenia<sup>^, ◊</sup>, Febrile neutropenia<sup>^, ◊</sup>, Thrombocytopenia<sup>^, ◊</sup>, Anaemia, Leucopenia<sup>◊</sup>, Lymphopenia</p>	<p><u>Very Common</u> Neutropenia<sup>^, ◊</sup>, Febrile neutropenia<sup>^, ◊</sup>, Thrombocytopenia<sup>^, ◊</sup>, Anaemia, Leucopenia<sup>◊</sup>, Lymphopenia</p> <p><u>Common</u> Pancytopenia<sup>◊</sup></p>
<b>Metabolism and Nutrition Disorders</b>	<p><u>Very Common</u> Hypokalaemia</p>	<p><u>Common</u> Hypokalaemia, Dehydration</p>
<b>Nervous System Disorders</b>	<p><u>Very Common</u> Paraesthesia</p> <p><u>Common</u> Peripheral neuropathy<sup>c</sup></p>	<p><u>Common</u> Headache</p>
<b>Vascular Disorders</b>	<p><u>Common</u> Pulmonary embolism<sup>◊, *</sup></p>	<p><u>Common</u> Deep vein thrombosis<sup>^, ◊, d</sup></p>
<b>Respiratory, Thoracic and Mediastinal Disorders</b>	<p><u>Very Common</u> Cough</p> <p><u>Common</u> Dyspnoea<sup>◊</sup>, Rhinorrhoea</p>	<p><u>Common</u> Dyspnoea<sup>◊</sup></p>
<b>Gastrointestinal Disorders</b>	<p><u>Very Common</u> Diarrhoea, Constipation, Abdominal pain, Nausea</p> <p><u>Common</u> Vomiting, Abdominal pain upper</p>	<p><u>Common</u> Diarrhoea, Vomiting, Nausea</p>
<b>Hepatobiliary Disorders</b>	<p><u>Very Common</u> Abnormal liver function tests</p>	<p><u>Common</u> Abnormal liver function tests</p>

System Organ Class/Preferred Term	All ADRs/Frequency	Grade 3-4 ADRs/Frequency
<b>Skin and Subcutaneous Tissue Disorders</b>	<u>Very Common</u> Rash, Dry skin	<u>Common</u> Rash, Pruritus
<b>Musculoskeletal and Connective Tissue Disorders</b>	<u>Very Common</u> Muscle spasms  <u>Common</u> Myalgia, Musculoskeletal pain	
<b>General Disorders and Administration Site Conditions</b>	<u>Very Common</u> Fatigue, Asthenia, Pyrexia	<u>Common</u> Fatigue, Asthenia

<sup>◇</sup> Adverse reactions reported as serious in clinical trials in patients with NDMM who had undergone ASCT

<sup>\*</sup> Applies to serious adverse drug reactions only

<sup>^</sup> See section 4.8 description of selected adverse reactions

<sup>a</sup> "Pneumonia" combined AE term includes the following PTs: Bronchopneumonia, Lobar pneumonia, Pneumocystis jiroveci pneumonia, Pneumonia, Pneumonia klebsiella, Pneumonia legionella, Pneumonia mycoplasma, Pneumonia pneumococcal, Pneumonia streptococcal, Pneumonia viral, Lung disorder, Pneumonitis

<sup>b</sup> "Sepsis" combined AE term includes the following PTs: Bacterial sepsis, Pneumococcal sepsis, Septic shock, Staphylococcal sepsis

<sup>c</sup> "Peripheral neuropathy" combined AE term includes the following preferred terms (PTs): Neuropathy peripheral, Peripheral sensory neuropathy, Polyneuropathy

<sup>d</sup> "Deep vein thrombosis" combined AE term includes the following PTs: Deep vein thrombosis, Thrombosis, Venous thrombosis

#### Tabulated summary for combination therapy in MM

The following table is derived from data gathered during the multiple myeloma studies with combination therapy. The data were not adjusted according to the longer duration of treatment in the lenalidomide-containing arms continued until disease progression versus the comparator arms in the pivotal multiple myeloma studies (See section 5.1).

**Table 2: Overall reported adverse drug reactions reported in pivotal clinical studies MM-020, MM-009 and MM-010 and post-marketing data in patients with multiple myeloma treated with Revlimid/Dexamethasone**

System Organ Class / Preferred Term	All ADRs/Frequency	Grade 3-4 ADRs/Frequency
<b>Infections and Infestations</b>	<u>Very Common</u> Pneumonia <sup>◇</sup> , Upper respiratory tract infection <sup>◇</sup> , Bacterial, viral and fungal infections (including opportunistic infections) <sup>◇</sup> , Nasopharyngitis, Pharyngitis, Bronchitis <sup>◇</sup>  <u>Common</u> Sepsis <sup>◇</sup> , Sinusitis <sup>◇</sup>  <u>Not Known</u> <sup>†</sup> Viral infections, including herpes zoster and hepatitis B virus reactivation <sup>†</sup>	<u>Common</u> Pneumonia <sup>◇</sup> , Bacterial, viral and fungal infections (including opportunistic infections) <sup>◇</sup> , Cellulitis <sup>◇</sup> , Sepsis <sup>◇</sup> , Bronchitis <sup>◇</sup>  <u>Not Known</u> <sup>†</sup> Viral infections, including herpes zoster and hepatitis B virus reactivation <sup>†</sup>

<b>Neoplasms Benign, Malignant and Unspecified (incl cysts and polyps)</b>	<u>Uncommon</u> Basal cell carcinoma <sup>^,◇</sup> , Squamous skin cancer <sup>^,◇,*</sup>	<u>Common</u> Acute myeloid leukaemia <sup>◇</sup> , Myelodysplastic syndrome <sup>◇</sup> , Squamous cell carcinoma of skin <sup>^,◇,**</sup>  <u>Uncommon</u> T-cell type acute leukaemia <sup>◇</sup> , Basal cell carcinoma <sup>^,◇</sup> , Tumour lysis syndrome  <u>Rare†</u> Tumour lysis syndrome†
<b>Blood and Lymphatic System Disorders</b>	<u>Very Common</u> Thrombocytopenia <sup>^,◇</sup> , Neutropenia <sup>^,◇</sup> , Anemia <sup>◇</sup> , Haemorrhagic disorder <sup>^</sup> , Leucopenias  <u>Common</u> Febrile neutropenia <sup>^,◇</sup> , Pancytopenia <sup>◇</sup>  <u>Uncommon</u> Haemolysis, Autoimmune haemolytic anemia, Haemolytic anemia  <u>Not Known†</u> Acquired haemophilia†	<u>Very Common</u> Thrombocytopenia <sup>^,◇</sup> , Neutropenia <sup>^,◇</sup> , Anemia <sup>◇</sup> , Leucopenias  <u>Common</u> Febrile neutropenia <sup>^,◇</sup> , Pancytopenia <sup>◇</sup> , Haemolytic anemia  <u>Uncommon</u> Hypercoagulation, Coagulopathy
<b>Immune System Disorders</b>	<u>Uncommon</u> Hypersensitivity <sup>^</sup>  <u>Rare†</u> Anaphylactic reaction <sup>^†</sup>  <u>Not Known†</u> Solid organ transplant rejection†	<u>Rare†</u> Anaphylactic reaction <sup>^†</sup>
<b>Endocrine Disorders</b>	<u>Common</u> Hypothyroidism, Hyperthyroidism†	
<b>Metabolism and Nutrition Disorders</b>	<u>Very Common</u> Hypokalaemia <sup>◇</sup> , Hyperglycaemia, Hypocalcaemia <sup>◇</sup> , Decreased appetite, Weight decreased  <u>Common</u> Hypomagnesaemia, Hyperuricaemia, Dehydration <sup>◇</sup> , Hypercalcaemia <sup>+</sup>	<u>Common</u> Hypokalaemia <sup>◇</sup> , Hyperglycaemia, Hypocalcaemia <sup>◇</sup> , Diabetes mellitus <sup>◇</sup> , Hypophosphataemia, Hyponatraemia <sup>◇</sup> , Hyperuricaemia, Gout, Decreased appetite, Weight decreased
<b>Psychiatric Disorders</b>	<u>Very Common</u> Depression, Insomnia  <u>Uncommon</u> Loss of libido	<u>Common</u> Depression, Insomnia

<b>Nervous System Disorders</b>	<u>Very Common</u> Peripheral neuropathies (excluding motor neuropathy), Dizziness, Tremor, Dysgeusia, Headache  <u>Common</u> Ataxia, Balance impaired	<u>Common</u> Cerebrovascular accident <sup>◊</sup> , Dizziness, Syncope  <u>Uncommon</u> Intracranial haemorrhage <sup>^</sup> , Transient ischaemic attack, Cerebral ischaemia
<b>Eye Disorders</b>	<u>Very Common</u> Cataracts, Blurred vision  <u>Common</u> Reduced visual acuity	<u>Common</u> Cataract  <u>Uncommon</u> Blindness
<b>Ear and Labyrinth Disorders</b>	<u>Common</u> Deafness (Including Hypoacusis), Tinnitus	
<b>Cardiac Disorders</b>	<u>Common</u> Atrial fibrillation <sup>◊</sup> , Bradycardia  <u>Uncommon</u> Arrhythmia, QT prolongation, Atrial flutter, Ventricular extrasystoles	<u>Common</u> Myocardial infarction (including acute) <sup>^,◊</sup> , Atrial fibrillation <sup>◊</sup> , Congestive cardiac failure <sup>◊</sup> , Tachycardia, Cardiac failure <sup>◊</sup> , Myocardial ischaemia <sup>◊</sup>
<b>Vascular Disorders</b>	<u>Very Common</u> Venous thromboembolic events, predominantly deep vein thrombosis and pulmonary embolism <sup>^,◊</sup>  <u>Common</u> Hypotension <sup>◊</sup> , Hypertension, Ecchymosis <sup>^</sup>	<u>Very Common</u> Venous thromboembolic events, predominantly deep vein thrombosis and pulmonary embolism <sup>^,◊</sup>  <u>Common</u> Vasculitis  <u>Uncommon</u> Ischemia, Peripheral ischemia, Intracranial venous sinus thrombosis
<b>Respiratory, Thoracic and Mediastinal Disorders</b>	<u>Very Common</u> Dyspnoea <sup>◊</sup> , Epistaxis <sup>^</sup>  <u>Uncommon</u> <sup>†</sup> Pulmonary hypertension <sup>†</sup>	<u>Common</u> Respiratory distress <sup>◊</sup> , Dyspnoea <sup>◊</sup>  <u>Rare</u> <sup>†</sup> Pulmonary hypertension <sup>†</sup>  <u>Not Known</u> <sup>†</sup> Interstitial pneumonitis <sup>†</sup>

<b>Gastrointestinal Disorders</b>	<p><u>Very Common</u> Constipation<sup>◊</sup>, Diarrhoea<sup>◊</sup>, Nausea, Vomiting, Abdominal pain<sup>◊</sup>, Dyspepsia</p> <p><u>Common</u> Gastrointestinal haemorrhage (including rectal haemorrhage, haemorrhoidal haemorrhage, peptic ulcer haemorrhage and gingival bleeding)<sup>^</sup>, Dry mouth, Stomatitis, Dysphagia</p> <p><u>Uncommon</u> Colitis, Caecitis</p>	<p><u>Common</u> Constipation<sup>◊</sup>, Diarrhoea<sup>◊</sup>, Abdominal pain<sup>◊</sup>, Nausea, Vomiting</p> <p><u>Not Known</u><sup>†</sup> Pancreatitis<sup>†</sup>, Gastrointestinal perforation (including diverticular, intestinal and large intestine perforations)<sup>^</sup> <sup>†</sup></p>
<b>Hepatobiliary Disorders</b>	<p><u>Common</u> Abnormal liver function tests<sup>◊</sup></p> <p><u>Uncommon</u> Hepatic failure<sup>^</sup></p> <p><u>Not Known</u><sup>†</sup> Acute hepatic failure<sup>^</sup><sup>†</sup>, Hepatitis toxic<sup>^</sup><sup>†</sup>, Cytolytic hepatitis<sup>^</sup><sup>†</sup>, Cholestatic hepatitis<sup>^</sup><sup>†</sup>, Mixed cytolytic/cholestatic hepatitis<sup>^</sup><sup>†</sup></p>	<p><u>Common</u> Cholestasis<sup>◊</sup>, Abnormal liver function tests<sup>◊</sup></p> <p><u>Uncommon</u> Hepatic failure<sup>^</sup></p> <p><u>Not Known</u><sup>†</sup> Acute hepatic failure<sup>^</sup><sup>†</sup>, Hepatitis toxic<sup>^</sup><sup>†</sup></p>
<b>Skin and Subcutaneous Tissue Disorders</b>	<p><u>Very Common</u> Rashes, Pruritus</p> <p><u>Common</u> Urticaria, Hyperhidrosis, Dry skin, Skin hyperpigmentation, Eczema, Erythema</p> <p><u>Uncommon</u> Skin discolouration, Photosensitivity reaction</p>	<p><u>Common</u> Rashes</p> <p><u>Uncommon</u><sup>†</sup> Angioedema<sup>†</sup></p> <p><u>Rare</u><sup>†</sup> Stevens-Johnson Syndrome<sup>^</sup><sup>†</sup>, Toxic epidermal necrolysis<sup>^</sup><sup>†</sup></p> <p><u>Not Known</u><sup>†</sup> Leukocytoclastic vasculitis<sup>†</sup>, Drug Reaction with Eosinophilia and Systemic Symptoms<sup>^</sup><sup>†</sup></p>
<b>Musculoskeletal and Connective Tissue Disorders</b>	<p><u>Very Common</u> Muscle spasms, Bone pain<sup>◊</sup>, Musculoskeletal and connective tissue pain and discomfort (including back pain<sup>◊</sup>), Arthralgia<sup>◊</sup></p> <p><u>Common</u> Joint swelling, Muscular weakness, Myalgia</p>	<p><u>Common</u> Muscular weakness, Bone pain<sup>◊</sup>, Musculoskeletal and connective tissue pain and discomfort (including back pain<sup>◊</sup>)</p> <p><u>Uncommon</u> Joint swelling</p>

<b>Renal and Urinary Disorders</b>	<u>Very Common</u> Renal failure (including acute) <sup>◇</sup>  <u>Common</u> Haematuria <sup>^</sup> , Urinary retention, Urinary incontinence  <u>Uncommon</u> Acquired Fanconi syndrome	<u>Uncommon</u> Renal tubular necrosis
<b>Reproductive System and Breast Disorders</b>	<u>Common</u> Erectile dysfunction	
<b>General Disorders and Administration Site Conditions</b>	<u>Very Common</u> Fatigue <sup>◇</sup> , Oedema (including peripheral oedema), Pyrexia <sup>◇</sup> , Influenza like illness syndrome (including pyrexia, cough, myalgia, musculoskeletal pain, headache and rigors), Asthenia  <u>Common</u> Chest pain, Lethargy	<u>Common</u> Fatigue <sup>◇</sup> , Pyrexia <sup>◇</sup> , Asthenia
<b>Investigations</b>	<u>Common</u> C-reactive protein increased	
<b>Injury, Poisoning and Procedural Complications</b>	<u>Common</u> Fall, Contusion <sup>^</sup>	

<sup>^</sup>see section 4.8 description of selected adverse reactions

<sup>†</sup>reports from post-marketing data

<sup>◇</sup> Adverse reactions reported as serious in clinical trials in patients with multiple myeloma treated with lenalidomide in combination with dexamethasone, or with melphalan and prednisone

<sup>+</sup>Applies to serious adverse drug reactions only

\*Squamous skin cancer was reported in clinical trials in previously treated myeloma patients with lenalidomide/dexamethasone compared to controls

\*\*Squamous cell carcinoma of skin was reported in a clinical trial in newly diagnosed multiple myeloma patients with lenalidomide/dexamethasone compared to controls

### Description of selected adverse reactions

#### Teratogenicity

Lenalidomide is structurally related to thalidomide. Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects. In monkeys, lenalidomide induced malformations similar to those described with thalidomide (see sections 4.6 and 5.3). If lenalidomide is taken during pregnancy, a teratogenic effect of lenalidomide in humans is expected.

#### Neutropenia and thrombocytopenia

- Newly diagnosed multiple myeloma: patients who have undergone ASCT treated with lenalidomide maintenance

Lenalidomide maintenance after ASCT is associated with a higher frequency of Grade 4 neutropenia compared to placebo maintenance (32.1% vs 26.7% [16.1% vs 1.8% after the start of maintenance treatment] in CALGB 100104 and 16.4% vs 0.7% in IFM 2005-02, respectively). Treatment-emergent AEs of neutropenia leading to lenalidomide discontinuation were reported in 2.2% of patients in CALGB 100104 and 2.4% of patients in IFM 2005-02, respectively. Grade 4 febrile neutropenia was reported at similar frequencies in the lenalidomide maintenance arms compared to placebo maintenance arms in both

studies (0.4% vs 0.5% [0.4% vs 0.5% after the start of maintenance treatment] in CALGB 100104 and 0.3% vs 0% in IFM 2005-02, respectively).

Lenalidomide maintenance after ASCT is associated with a higher frequency of Grade 3 or 4 thrombocytopenia compared to placebo maintenance (37.5% vs 30.3% [17.9% vs 4.1% after the start of maintenance treatment] in CALGB 100104 and 13.0% vs 2.9% in IFM 2005-02, respectively).

- Newly diagnosed multiple myeloma: patients who are not eligible for transplant treated with lenalidomide in combination with low dose dexamethasone

The combination of lenalidomide with low dose dexamethasone in newly diagnosed multiple myeloma patients is associated with a lower frequency of Grade 4 neutropenia (8.5% in Rd and Rd18, compared with MPT (15%). Grade 4 febrile neutropenia was observed infrequently (0.6% in Rd and Rd18 compared with 0.7% in MPT).

The combination of lenalidomide with low dose dexamethasone in newly diagnosed multiple myeloma patients is associated with a lower frequency of Grade 3 and 4 thrombocytopenia (8.1% in Rd and Rd18) compared with MPT (11%).

- Multiple myeloma: patients with at least one prior therapy

The combination of lenalidomide with dexamethasone in multiple myeloma patients is associated with a higher incidence of Grade 4 neutropenia (5.1% in lenalidomide/dexamethasone-treated patients compared with 0.6% in placebo/dexamethasone-treated patients). Grade 4 febrile neutropenia episodes were observed infrequently (0.6% in lenalidomide/dexamethasone-treated patients compared to 0.0% in placebo/dexamethasone treated patients).

The combination of lenalidomide with dexamethasone in multiple myeloma patients is associated with a higher incidence of Grade 3 and Grade 4 thrombocytopenia (9.9% and 1.4%, respectively, in lenalidomide/dexamethasone-treated patients compared to 2.3% and 0.0% in placebo/dexamethasone-treated patients).

#### Venous thromboembolism

An increased risk of DVT and PE is associated with the use of the combination of lenalidomide with dexamethasone in patients with multiple myeloma, and to a lesser extent in patients with multiple myeloma treated with lenalidomide monotherapy (see section 4.5).

Concomitant administration of erythropoietic agents or previous history of DVT may also increase thrombotic risk in these patients.

#### Myocardial infarction

Myocardial infarction has been reported in patients receiving lenalidomide, particularly in those with known risk factors.

#### Haemorrhagic disorders

Haemorrhagic disorders are listed under several system organ classes: Blood and lymphatic system disorders; nervous system disorders (intracranial haemorrhage); respiratory, thoracic and mediastinal disorders (epistaxis); gastrointestinal disorders (gingival bleeding, haemorrhoidal haemorrhage, rectal haemorrhage); renal and urinary disorders (haematuria); injury, poisoning and procedural complications (contusion) and vascular disorders (ecchymosis).

#### Allergic reactions and severe skin reactions

Cases of allergic reactions including angioedema, anaphylactic reaction and severe cutaneous reactions including SJS, TEN and DRESS have been reported with the use of lenalidomide. A possible cross-reaction between lenalidomide and thalidomide has been reported in the literature. Patients with a history of severe rash associated with thalidomide treatment should not receive lenalidomide (see section 4.4).

#### Second primary malignancies

In clinical trials in previously treated myeloma patients with lenalidomide/dexamethasone compared to controls, mainly comprising of basal cell or squamous cell skin cancers.



#### Acute myeloid leukaemia

- **Multiple myeloma**

Cases of AML have been observed in clinical trials of newly diagnosed multiple myeloma in patients taking lenalidomide treatment in combination with melphalan or immediately following HDM/ASCT (see section 4.4). This increase was not observed in clinical trials of newly diagnosed multiple myeloma in patients taking lenalidomide in combination with low dose dexamethasone compared to thalidomide in combination with melphalan and prednisone.

#### Hepatic disorders

The following post-marketing adverse reactions have been reported (frequency unknown): acute hepatic failure and cholestasis (both potentially fatal), toxic hepatitis, cytolytic hepatitis, mixed cytolytic/cholestatic hepatitis.

#### Rhabdomyolysis

Rare cases of rhabdomyolysis have been observed, some of them when lenalidomide is administered with a statin.

#### Thyroid disorders

Cases of hypothyroidism and cases of hyperthyroidism have been reported (see section 4.4 Thyroid disorders).

#### Gastrointestinal disorders

Gastrointestinal perforations have been reported during treatment with lenalidomide. Gastrointestinal perforations may lead to septic complications and may be associated with fatal outcome.

#### Acute Graft Versus Host Disease

In the literature and post-marketing setting, acute graft-versus-host disease has been reported with lenalidomide therapy following allogeneic hematopoietic transplant.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

### **4.9 Overdose**

There is no specific experience in the management of lenalidomide overdose in multiple myeloma patients, although in dose-ranging studies some patients were exposed to up to 150 mg, and in single-dose studies, some patients were exposed to up to 400 mg. The dose limiting toxicity in these studies was essentially haematological. In the event of overdose, supportive care is advised.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Other immunosuppressants. ATC code: L04AX04.

#### Mechanism of action

Lenalidomide binds directly to cereblon, a component of a cullin ring E3 ubiquitin ligase enzyme complex that includes deoxyribonucleic acid (DNA) damage-binding protein 1 (DDB1), cullin 4 (CUL4), and regulator of cullins 1 (Roc1). In haematopoietic cells, lenalidomide binding to cereblon recruits substrate proteins Aiolos and Ikaros, lymphoid transcriptional factors, leading to their ubiquitination and subsequent degradation resulting in direct cytotoxic and immunomodulatory effects.

Specifically, lenalidomide inhibits proliferation and enhances apoptosis of certain haematopoietic tumour cells (including MM plasma tumour cells, follicular lymphoma tumour cells and those with deletions of chromosome 5), enhances T cell- and Natural Killer (NK) cell-mediated immunity and increases the number of NK, T and NK T cells.

The lenalidomide mechanism of action also includes additional activities such as anti-angiogenic and pro-erythropoietic properties. Lenalidomide inhibits angiogenesis by blocking the migration and adhesion of endothelial cells and the formation of microvessels, augments foetal haemoglobin production by CD34+ haematopoietic stem cells, and inhibits production of pro-inflammatory cytokines (e.g., TNF- $\alpha$  and IL-6) by monocytes.

#### Clinical efficacy and safety

Lenalidomide efficacy and safety have been evaluated in five phase 3 studies in newly diagnosed multiple myeloma and two phase 3 studies in relapsed refractory multiple myeloma as described below.

#### Newly diagnosed multiple myeloma

- Lenalidomide maintenance in patients who have undergone ASCT

The efficacy and safety of lenalidomide maintenance was assessed in two phase 3 multicentre, randomised, double-blind 2-arm, parallel group, placebo-controlled studies: CALGB 100104 and IFM 2005-02

#### *CALGB 100104*

Patients between 18 and 70 years of age with active MM requiring treatment and without prior progression after initial therapy were eligible.

Patients were randomised 1:1 within 90-100 days after ASCT to receive either lenalidomide or placebo maintenance. The maintenance dose was 10 mg once daily on days 1-28 of repeated 28-day cycles (increased up to 15 mg once daily after 3 months in the absence of dose-limiting toxicity), and treatment was continued until disease progression.

The primary efficacy endpoint in the study was progression free survival (PFS) from randomisation to the date of progression or death, whichever occurred first; the study was not powered for the overall survival endpoint. In total 460 patients were randomised: 231 patients to Lenalidomide and 229 patients to placebo. The demographic and disease-related characteristics were balanced across both arms.

The study was unblinded upon the recommendations of the data monitoring committee after surpassing the threshold for a preplanned interim analysis of PFS. After unblinding, patients in the placebo arm were allowed to cross over to receive lenalidomide before disease progression.

The results of PFS at unblinding, following a preplanned interim analysis, using a cut-off of 17 December 2009 (15.5 months follow up) showed a 62% reduction in risk of disease progression or death favouring lenalidomide (HR = 0.38; 95% CI 0.27, 0.54; p <0.001). The median overall PFS was 33.9 months (95% CI NE, NE) in the lenalidomide arm versus 19.0 months (95% CI 16.2, 25.6) in the placebo arm.

The PFS benefit was observed both in the subgroup of patients with CR and in the subgroup of patients who had not achieved a CR.

The results for the study, using a cut-off of 1 February 2016, are presented in Table 3

**Table 3: Summary of overall efficacy data**

	Lenalidomide (N = 231)	Placebo (N = 229)
<b>Investigator-assessed PFS</b>		
Median <sup>a</sup> PFS time, months (95% CI) <sup>b</sup>	<b>56.9</b> (41.9, 71.7)	<b>29.4</b> (20.7, 35.5)
HR [95% CI] <sup>c</sup> ; p-value <sup>d</sup>	<b>0.61</b> (0.48, 0.76); <0.001	
<b>PFS2<sup>e</sup></b>		
Median <sup>a</sup> PFS2 time, months (95% CI) <sup>b</sup>	<b>80.2</b> (63.3, 101.8)	<b>52.8</b> (41.3, 64.0)
HR [95% CI] <sup>c</sup> ; p-value <sup>d</sup>	<b>0.61</b> (0.48, 0.78); <0.001	
<b>Overall survival</b>		

	Lenalidomide (N = 231)	Placebo (N = 229)
Median <sup>a</sup> OS time, months (95% CI) <sup>b</sup>	<b>111.0</b> (101.8, NE)	<b>84.2</b> (71.0, 102.7)
8-year survival rate, % (SE)	60.9 (3.78)	44.6 (3.98)
HR [95% CI] <sup>c</sup> ; p-value <sup>d</sup>	<b>0.61</b> (0.46, 0.81); <0.001	
<b>Follow-up</b>		
Median <sup>f</sup> (min, max), months: all surviving patients	<b>81.9</b> (0.0, 119.8)	<b>81.0</b> (4.1, 119.5)

CI = confidence interval; HR = hazard ratio; max = maximum; min = minimum; NE = not estimable; OS = overall survival;

PFS = progression-free survival

<sup>a</sup> The median is based on the Kaplan-Meier estimate.

<sup>b</sup> The 95% CI about the median.

<sup>c</sup> Based on Cox proportional hazards model comparing the hazard functions associated with the indicated treatment arms.

<sup>d</sup> The p-value is based on the unstratified log-rank test of Kaplan-Meier curve differences between the indicated treatment arms.

<sup>e</sup> Exploratory endpoint (PFS2). Lenalidomide received by subjects in the placebo arm who crossed over prior to PD upon study unblinding was not considered as a second-line therapy.

<sup>f</sup> Median follow-up post-ASCT for all surviving subjects.

**Data cuts:** 17 Dec 2009 and 01 Feb 2016

### *IFM 2005-02*

Patients aged < 65 years at diagnosis who had undergone ASCT and had achieved at least a stable disease response at the time of hematologic recovery were eligible. Patients were randomised 1:1 to receive either lenalidomide or placebo maintenance (10 mg once daily on days 1-28 of repeated 28-day cycles increased up to 15 mg once daily after 3 months in the absence of dose-limiting toxicity) following 2 courses of lenalidomide consolidation (25 mg/day, days 1-21 of a 28-day cycle). Treatment was to be continued until disease progression.

The primary endpoint was PFS defined from randomisation to the date of progression or death, whichever occurred first; the study was not powered for the overall survival endpoint. In total 614 patients were randomised: 307 patients to lenalidomide and 307 patients to placebo.

The study was unblinded upon the recommendations of the data monitoring committee after surpassing the threshold for a preplanned interim analysis of PFS. After unblinding, patients receiving placebo were not crossed over to lenalidomide therapy prior to progressive disease. The lenalidomide arm was discontinued, as a proactive safety measure, after observing an imbalance of SPMs (see Section 4.4).

The results of PFS at unblinding, following a preplanned interim analysis, using a cut-off of 7 July 2010 (31.4 months follow up) showed a 48% reduction in risk of disease progression or death favouring lenalidomide (HR = 0.52; 95% CI 0.41, 0.66; p < 0.001). The median overall PFS was 40.1 months (95% CI 35.7, 42.4) in the lenalidomide arm versus 22.8 months (95% CI 20.7, 27.4) in the placebo arm.

The PFS benefit was less in the subgroup of patients with CR than in the subgroup of patients who had not achieved a CR.

The updated PFS, using a cut-off of 1 February 2016 (96.7 months follow up) continues to show a PFS advantage: HR = 0.57 (95% CI 0.47, 0.68; p < 0.001). The median overall PFS was 44.4 months (39.6, 52.0) in the lenalidomide arm versus 23.8 months (95% CI 21.2, 27.3) in the placebo arm. For PFS2, the observed HR was 0.80 (95% CI 0.66, 0.98; p = 0.026) for lenalidomide versus placebo. The median overall PFS2 was 69.9 months (95% CI 58.1, 80.0) in the lenalidomide arm versus 58.4 months (95% CI 51.1, 65.0) in the placebo arm. For OS, the observed HR was 0.90: (95% CI 0.72, 1.13; p = 0.355) for lenalidomide versus placebo. The median overall survival time was 105.9 months (95% CI 88.8, NE) in the lenalidomide arm versus 88.1 months (95% CI 80.7, 108.4) in the placebo arm.

- Lenalidomide in combination with dexamethasone in patients who are not eligible for stem cell transplantation

The safety and efficacy of lenalidomide was assessed in a Phase 3, multicentre, randomised, open-label, 3-arm study (MM-020) of patients who were at least 65 years of age or older or, if younger than 65 years of age, were not candidates for stem cell transplantation because they declined to undergo stem cell transplantation or stem cell transplantation is not available to the patient due to cost or other reason. The study (MM-020) compared lenalidomide and dexamethasone (Rd) given for 2 different durations

of time (i.e., until progressive disease [Arm Rd] or for up to eighteen 28-day cycles [72 weeks, Arm Rd18]) to melphalan, prednisone and thalidomide (MPT) for a maximum of twelve 42-day cycles (72 weeks). Patients were randomised (1:1:1) to 1 of 3 treatment arms. Patients were stratified at randomisation by age ( $\leq 75$  versus  $> 75$  years), stage (ISS Stages I and II versus Stage III), and country.

Patients in the Rd and Rd18 arms took lenalidomide 25 mg once daily on days 1 to 21 of 28-day cycles according to protocol arm. Dexamethasone 40 mg was dosed once daily on days 1, 8, 15, and 22 of each 28-day cycle. Initial dose and regimen for Rd and Rd18 were adjusted according to age and renal function (see section 4.2). Patients  $> 75$  years received a dexamethasone dose of 20 mg once daily on days 1, 8, 15, and 22 of each 28-day cycle. All patients received prophylactic anticoagulation (low molecular weight heparin, warfarin, heparin, low-dose aspirin) during the study.

The primary efficacy endpoint in the study was progression free survival (PFS). In total 1623 patients were enrolled into the study, with 535 patients randomised to Rd, 541 patients randomised to Rd18 and 547 patients randomised to MPT. The demographics and disease-related baseline characteristics of the patients were well balanced in all 3 arms. In general, study subjects had advanced-stage disease: of the total study population, 41% had ISS stage III, 9% had severe renal insufficiency (creatinine clearance [CLCr]  $< 30$  mL/min). The median age was 73 in the 3 arms.

In an updated analysis of PFS, PFS2 and OS using a cut off of 3 March 2014 where the median follow-up time for all surviving subjects was 45.5 months, the results of the study are presented in Table 4:

**Table 4: Summary of overall efficacy data**

	<b>Rd (N = 535)</b>	<b>Rd18 (N = 541)</b>	<b>MPT (N = 547)</b>
<b>Investigator-assessed PFS (months)</b>			
Median <sup>a</sup> PFS time, months (95% CI) <sup>b</sup>	26.0 (20.7, 29.7)	21.0 (19.7, 22.4)	21.9 (19.8, 23.9)
HR [95% CI] <sup>c</sup> ; p-value <sup>d</sup>			
Rd vs MPT	0.69 (0.59, 0.80); <0.001		
Rd vs Rd18	0.71 (0.61, 0.83); <0.001		
Rd18 vs MPT	0.99 (0.86, 1.14); 0.866		
<b>PFS2<sup>e</sup> (months)</b>			
Median <sup>a</sup> PFS2 time, months (95% CI) <sup>b</sup>	42.9 (38.1, 47.4)	40.0 (36.2, 44.2)	35.0 (30.4, 37.8)
HR [95% CI] <sup>c</sup> ; p-value <sup>d</sup>			
Rd vs MPT	0.74 (0.63, 0.86); <0.001		
Rd vs Rd18	0.92 (0.78, 1.08); 0.316		
Rd18 vs MPT	0.80 (0.69, 0.93); 0.004		
<b>Overall survival (months)</b>			
Median <sup>a</sup> OS time, months (95% CI) <sup>b</sup>	58.9 (56.0, NE)	56.7 (50.1, NE)	48.5 (44.2, 52.0)
HR [95% CI] <sup>c</sup> ; p-value <sup>d</sup>			
Rd vs MPT	0.75 (0.62, 0.90); 0.002		
Rd vs Rd18	0.91 (0.75, 1.09); 0.305		
Rd18 vs MPT	0.83 (0.69, 0.99); 0.034		
<b>Follow-up (months)</b>			
Median <sup>f</sup> (min, max): all patients	40.8 (0.0, 65.9)	40.1 (0.4, 65.7)	38.7 (0.0, 64.2)
<b>Myeloma response<sup>g</sup> n (%)</b>			
CR	81 (15.1)	77 (14.2)	51 (9.3)
VGPR	152 (28.4)	154 (28.5)	103 (18.8)
PR	169 (31.6)	166 (30.7)	187 (34.2)
Overall response: CR, VGPR, or PR	402 (75.1)	397 (73.4)	341 (62.3)
<b>Duration of response (months)<sup>h</sup></b>			
Median <sup>a</sup> (95% CI) <sup>b</sup>	35.0 (27.9, 43.4)	22.1 (20.3, 24.0)	22.3 (20.2, 24.9)

AMT = antimyeloma therapy; CI = confidence interval; CR = complete response; d = low-dose dexamethasone; HR = hazard ratio; IMWG = International Myeloma Working Group; IRAC = Independent Response Adjudication Committee; M = melphalan; max = maximum; min = minimum; NE = not estimable; OS = overall survival; P = prednisone; PFS = progression-free survival; PR = partial response; R = lenalidomide; Rd = Rd given until documentation of progressive disease; Rd18 = Rd given for  $\leq 18$  cycles; SE = standard error; T = thalidomide; VGPR = very good partial response; vs = versus.

<sup>a</sup> The median is based on the Kaplan-Meier estimate.

- <sup>b</sup> The 95% CI about the median.
- <sup>c</sup> Based on Cox proportional hazards model comparing the hazard functions associated with the indicated treatment arms.
- <sup>d</sup> The p-value is based on the unstratified log-rank test of Kaplan-Meier curve differences between the indicated treatment arms.
- <sup>e</sup> Exploratory endpoint (PFS2)
- <sup>f</sup> The median is the univariate statistic without adjusting for censoring.
- <sup>g</sup> Best assessment of adjudicated response during the treatment phase of the study (for definitions of each response category, Data cutoff date = 24 May 2013).
- <sup>h</sup> data cut 24 May 2013

### *Supportive newly diagnosed multiple myeloma studies*

An open-label, randomised, multicentre, Phase 3 study (ECOG E4A03) was conducted in 445 patients with newly diagnosed multiple myeloma; 222 patients were randomised to the lenalidomide/low dose dexamethasone arm, and 223 were randomised to the lenalidomide/standard dose dexamethasone arm. Patients randomised to the lenalidomide/standard dose dexamethasone arm received lenalidomide 25 mg/day, days 1 to 21 every 28 days plus dexamethasone 40 mg/day on days 1 to 4, 9 to 12, and 17 to 20 every 28 days for the first four cycles. Patients randomised to the lenalidomide/low dose dexamethasone arm received lenalidomide 25 mg/day, days 1 to 21 every 28 days plus low dose dexamethasone – 40 mg/day on days 1, 8, 15, and 22 every 28 days. In the lenalidomide/low dose dexamethasone group, 20 patients (9.1%) underwent at least one dose interruption compared to 65 patients (29.3%) in the lenalidomide/standard dose dexamethasone arm.

In a post-hoc analysis, lower mortality was observed in the lenalidomide/low dose dexamethasone arm 6.8% (15/220) compared to the lenalidomide/standard dose dexamethasone arm 19.3% (43/223), in the newly diagnosed multiple myeloma patient population, with a median follow up of 72.3 weeks.

However, with a longer follow-up, the difference in overall survival in favour of lenalidomide/low dose dexamethasone tends to decrease.

### *Multiple myeloma with at least one prior therapy*

The efficacy and safety of lenalidomide were evaluated in two Phase 3 multi-centre, randomised, double-blind, placebo-controlled, parallel-group controlled studies (MM-009 and MM-010) of lenalidomide plus dexamethasone therapy versus dexamethasone alone in previously treated patients with multiple myeloma. Out of 353 patients in the MM-009 and MM-010 studies who received lenalidomide/dexamethasone, 45.6% were aged 65 or over. Of the 704 patients evaluated in the MM-009 and MM-010 studies, 44.6% were aged 65 or over.

In both studies, patients in the lenalidomide/dexamethasone (len/dex) group took 25 mg of lenalidomide orally once daily on days 1 to 21 and a matching placebo capsule once daily on days 22 to 28 of each 28-day cycle. Patients in the placebo/dexamethasone (placebo/dex) group took 1 placebo capsule on days 1 to 28 of each 28-day cycle. Patients in both treatment groups took 40 mg of dexamethasone orally once daily on days 1 to 4, 9 to 12, and 17 to 20 of each 28-day cycle for the first 4 cycles of therapy. The dose of dexamethasone was reduced to 40 mg orally once daily on days 1 to 4 of each 28-day cycle after the first 4 cycles of therapy. In both studies, treatment was to continue until disease progression. In both studies, dose adjustments were allowed based on clinical and laboratory finding.

The primary efficacy endpoint in both studies was time to progression (TTP). In total, 353 patients were evaluated in the MM-009 study; 177 in the len/dex group and 176 in the placebo/dex group and, in total, 351 patients were evaluated in the MM-010 study; 176 in the len/dex group and 175 in the placebo/dex group.

In both studies, the baseline demographic and disease-related characteristics were comparable between the len/dex and placebo/dex groups. Both patient populations presented a median age of 63 years, with a comparable male to female ratio. The ECOG performance status was comparable between both groups, as was the number and type of prior therapies.

Pre-planned interim analyses of both studies showed that len/dex was statistically significantly superior ( $p < 0.00001$ ) to dexamethasone alone for the primary efficacy endpoint, TTP (median follow-up duration of 98.0 weeks). Complete response and overall response rates in the len/dex arm were also significantly higher than the placebo/dex arm in both studies. Results of these analyses

subsequently led to an unblinding in both studies, in order to allow patients in the placebo/dex group to receive treatment with the len/dex combination.

An extended follow-up efficacy analysis was conducted with a median follow-up of 130.7 weeks. Table 5 summarises the results of the follow-up efficacy analyses – pooled studies MM-009 and MM-010.

In this pooled extended follow-up analysis, the median TTP was 60.1 weeks (95% CI: 44.3, 73.1) in patients treated with len/dex (N = 353) versus 20.1 weeks (95% CI: 17.7, 20.3) in patients treated with placebo/dex (N = 351). The median progression free survival was 48.1 weeks (95% CI: 36.4, 62.1) in patients treated with len/dex versus 20.0 weeks (95% CI: 16.1, 20.1) in patients treated with placebo/dex. The median duration of treatment was 44.0 weeks (min: 0.1, max: 254.9) for len/dex and 23.1 weeks (min: 0.3, max: 238.1) for placebo/dex. Complete response (CR), partial response (PR) and overall response (CR+PR) rates in the len/dex arm remain significantly higher than in the placebo/dex arm in both studies. The median overall survival in the extended follow-up analysis of the pooled studies is 164.3 weeks (95% CI: 145.1, 192.6) in patients treated with len/dex versus 136.4 weeks (95% CI: 113.1, 161.7) in patients treated with placebo/dex. Despite the fact that 170 out of the 351 patients randomised to placebo/dex received lenalidomide after disease progression or after the studies were unblinded, the pooled analysis of overall survival demonstrated a statistically significant survival advantage for len/dex relative to placebo/dex (HR = 0.833, 95% CI = [0.687, 1.009], p=0.045).

**Table 5: Summary of results of efficacy analyses as of cut-off date for extended follow-up — pooled studies MM-009 and MM-010 (cut-offs 23 July 2008 and 2 March 2008, respectively)**

Endpoint	len/dex (N=353)	placebo/dex (N=351)	
Time to event			HR [95% CI], p-value <sup>a</sup>
Time to progression Median [95% CI], weeks	60.1 [44.3, 73.1]	20.1 [17.7, 20.3]	0.350 [0.287, 0.426], p < 0.001
Progression free survival Median [95% CI], weeks	48.1 [36.4, 62.1]	20.0 [16.1, 20.1]	0.393 [0.326, 0.473] p < 0.001
Overall survival Median [95% CI], weeks 1-year Overall Survival rate	164.3 [145.1, 192.6] 82%	136.4 [113.1, 161.7] 75%	0.833 [0.687, 1.009] p = 0.045
Response rate			Odds ratio [95% CI], p-value <sup>b</sup>
Overall response [n, %] Complete response [n, %]	212 (60.1) 58 (16.4)	75 (21.4) 11 (3.1)	5.53 [3.97, 7.71], p < 0.001 6.08 [3.13, 11.80], p < 0.001

a: Two-tailed log rank test comparing survival curves between treatment groups.

b: Two-tailed continuity-corrected chi-square test.

## 5.2 Pharmacokinetic properties

Lenalidomide has an asymmetric carbon atom and can therefore exist as the optically active forms S(-) and R(+). Lenalidomide is produced as a racemic mixture. Lenalidomide is generally more soluble in organic solvents but exhibits the greatest solubility in 0.1N HCl buffer.

### Absorption

Lenalidomide is rapidly absorbed following oral administration in healthy volunteers, under fasting conditions, with maximum plasma concentrations occurring between 0.5 and 2 hours post-dose. In patients, as well as in healthy volunteers, the maximum concentration (C<sub>max</sub>) and area-under-the-concentration time curve (AUC) increase proportionally with increases in dose. Multiple dosing does not cause marked medicinal product accumulation. In plasma, the relative exposures of the S- and R-enantiomers of lenalidomide are approximately 56% and 44%, respectively.

Co-administration with a high-fat and high-calorie meal in healthy volunteers reduces the extent of absorption, resulting in an approximately 20% decrease in area under the concentration versus time

curve (AUC) and 50% decrease in  $C_{max}$  in plasma. However, in the main multiple myeloma registration trials where the efficacy and safety were established for lenalidomide, the medicinal product was administered without regard to food intake. Thus, lenalidomide can be administered with or without food.

#### Distribution

*In vitro* ( $^{14}C$ )-lenalidomide binding to plasma proteins was low with mean plasma protein binding at 23% and 29% in multiple myeloma patients and healthy volunteers, respectively.

Lenalidomide is present in human semen (< 0.01% of the dose) after administration of 25 mg/day and the medicinal product is undetectable in semen of a healthy subject 3 days after stopping the substance (see section 4.4).

#### Biotransformation and elimination

Results from human *in vitro* metabolism studies indicate that lenalidomide is not metabolised by cytochrome P450 enzymes suggesting that administration of lenalidomide with medicinal products that inhibit cytochrome P450 enzymes is not likely to result in metabolic medicinal product interactions in humans. *In vitro* studies indicate that lenalidomide has no inhibitory effect on CYP1A2, CYP2C9, CYP2C19, CYP2D6, CYP2E1, CYP3A, or UGT1A1. Therefore, lenalidomide is unlikely to cause any clinically relevant medicinal product interactions when co-administered with substrates of these enzymes.

*In vitro* studies indicate that lenalidomide is not a substrate of human breast cancer resistance protein (BCRP), multidrug resistance protein (MRP) transporters MRP1, MRP2, or MRP3, organic anion transporters (OAT) OAT1 and OAT3, organic anion transporting polypeptide 1B1 (OATP1B1), organic cation transporters (OCT) OCT1 and OCT2, multidrug and toxin extrusion protein (MATE) MATE1, and organic cation transporters novel (OCTN) OCTN1 and OCTN2.

*In vitro* studies indicate that lenalidomide has no inhibitory effect on human bile salt export pump (BSEP), BCRP, MRP2, OAT1, OAT3, OATP1B1, OATP1B3, and OCT2.

A majority of lenalidomide is eliminated through urinary excretion. The contribution of renal excretion to total clearance in subjects with normal renal function was 90%, with 4% of lenalidomide eliminated in faeces.

Lenalidomide is poorly metabolized as 82% of the dose is excreted unchanged in urine. Hydroxy-lenalidomide and N-acetyl-lenalidomide represent 4.59% and 1.83% of the excreted dose, respectively. The renal clearance of lenalidomide exceeds the glomerular filtration rate and therefore is at least actively secreted to some extent.

At doses of 5 to 25 mg/day, half-life in plasma is approximately 3 hours in healthy volunteers and ranges from 3 to 5 hours in patients with multiple myeloma.

#### Older people

No dedicated clinical studies have been conducted to evaluate pharmacokinetics of lenalidomide in the elderly. Population pharmacokinetic analyses included patients with ages ranging from 39 to 85 years old and indicate that age does not influence lenalidomide clearance (exposure in plasma). Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection and it would be prudent to monitor renal function.

#### Renal impairment

The pharmacokinetics of lenalidomide was studied in subjects with renal impairment due to nonmalignant conditions. In this study, two methods were used to classify renal function: the urinary creatinine clearance measured over 24 hours and the creatinine clearance estimated by Cockcroft-Gault formula. The results indicate that as renal function decreases (< 50 mL/min), the total lenalidomide clearance decreases proportionally resulting in an increase in AUC. The AUC was increased by approximately 2.5, 4 and 5-fold in subjects with moderate renal impairment, severe renal impairment, and end-stage renal disease, respectively, compared to the group combining subjects with

normal renal function and subjects with mild renal impairment. The half-life of lenalidomide increased from approximately 3.5 hours in subjects with creatinine clearance > 50 mL/min to more than 9 hours in subjects with reduced renal function < 50 mL/min. However, renal impairment did not alter the oral absorption of lenalidomide. The  $C_{max}$  was similar between healthy subjects and patients with renal impairment. Approximately 30% of the medicinal product in the body was removed during a single 4-hour dialysis session. Recommended dose adjustments in patients with impaired renal function are described in section 4.2.

#### Hepatic impairment

Population pharmacokinetic analyses included patients with mild hepatic impairment (N=16, total bilirubin >1 to  $\leq 1.5 \times$  ULN or AST > ULN) and indicate that mild hepatic impairment does not influence lenalidomide clearance (exposure in plasma). There are no data available for patients with moderate to severe hepatic impairment.

#### Other intrinsic factors

Population pharmacokinetic analyses indicate that body weight (33 - 135 kg), gender, race and type of haematological malignancy (MM, MDS or MCL) do not have a clinically relevant effect on lenalidomide clearance in adult patients.

### **5.3 Preclinical safety data**

An embryofoetal development study has been conducted in monkeys administered lenalidomide at doses from 0.5 and up to 4 mg/kg/day. Findings from this study indicate that lenalidomide produced external malformations including non-patent anus and malformations of upper and lower extremities (bent, shortened, malformed, malrotated and/or absent part of the extremities, oligo and/or polydactyly) in the offspring of female monkeys who received the active substance during pregnancy.

Various visceral effects (discoloration, red foci at different organs, small colorless mass above atrio-ventricular valve, small gall bladder, malformed diaphragm) were also observed in single fetuses.

Lenalidomide has a potential for acute toxicity; minimum lethal doses after oral administration were > 2000 mg/kg/day in rodents. Repeated oral administration of 75, 150 and 300 mg/kg/day to rats for up to 26 weeks produced a reversible treatment-related increase in kidney pelvis mineralisation in all 3 doses, most notably in females. The no observed adverse effect level (NOAEL) was considered to be less than 75 mg/kg/day, and is approximately 25-fold greater than the human daily exposure based on AUC exposure. Repeated oral administration of 4 and 6 mg/kg/day to monkeys for up to 20 weeks produced mortality and significant toxicity (marked weight loss, reduced red and white blood cell and platelet counts, multiple organ haemorrhage, gastrointestinal tract inflammation, lymphoid, and bone marrow atrophy). Repeated oral administration of 1 and 2 mg/kg/day to monkeys for up to 1 year produced reversible changes in bone marrow cellularity, a slight decrease in myeloid/erythroid cell ratio and thymic atrophy. Mild suppression of white blood cell count was observed at 1 mg/kg/day corresponding to approximately the same human dose based on AUC comparisons.

*In vitro* (bacterial mutation, human lymphocytes, mouse lymphoma, Syrian Hamster Embryo cell transformation) and *in vivo* (rat micronucleus) mutagenicity studies revealed no drug related effects at either the gene or chromosomal level. Carcinogenicity studies with lenalidomide have not been conducted.

Developmental toxicity studies were previously conducted in rabbits. In these studies, rabbits were administered 3, 10 and 20 mg/kg/day orally. An absence of the intermediate lobe of the lung was observed at 10 and 20 mg/kg/day with dose dependence and displaced kidneys were observed at 20 mg/kg/day. Although it was observed at maternotoxic levels they may be attributable to a direct effect. Soft tissue and skeletal variations in the foetuses were also observed at 10 and 20 mg/kg/day.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**



Revlimid® 5 mg hard capsules

Capsule contents

Anhydrous lactose  
Microcrystalline cellulose  
Croscarmellose sodium  
Magnesium stearate

Capsule shell

Gelatin  
Titanium dioxide (E171)

Printing ink

Shellac  
Propylene glycol  
(E1520)  
Black iron oxide (E172)  
Potassium hydroxide

Revlimid® 10 mg hard capsules

Capsule contents

Anhydrous lactose  
Microcrystalline cellulose  
Croscarmellose sodium  
Magnesium stearate

Capsule shell

Gelatin  
Titanium dioxide (E171)  
Indigo carmine (E132)  
Yellow iron oxide (E172)

Printing ink

Shellac  
Propylene glycol  
(E1520)  
Black iron oxide (E172)  
Potassium hydroxide

Revlimid® 15 mg hard capsules

Capsule contents

Anhydrous lactose  
Microcrystalline cellulose  
Croscarmellose sodium  
Magnesium stearate

Capsule shell

Gelatin  
Titanium dioxide (E171)  
Indigo carmine (E132)

Printing ink

Shellac  
Propylene glycol  
(E1520)

Black iron oxide (E172)  
Potassium hydroxide

Revlimid® 25 mg hard capsules

Capsule contents

Anhydrous lactose  
Microcrystalline cellulose  
Croscarmellose sodium  
Magnesium stearate

Capsule shell

Gelatin  
Titanium dioxide (E171)

Printing ink

Shellac  
Propylene glycol  
(E1520)  
Black iron oxide (E172)  
Potassium hydroxide

**6.2 Incompatibilities**

Not applicable.

**6.3 Special precautions for storage**

Do not store above 25 °C.

**6.4 Nature and contents of container**

The capsules are provided in carton packs. Each pack contains three Polyvinylchloride (PVC) / Polychlorotrifluoroethylene (PCTFE) / Aluminium foil blisters containing 7 hard capsules. This gives a total of 21 capsules per pack.

**6.5 Special precautions for disposal and other handling**

Capsules should not be opened or crushed. If powder from lenalidomide makes contact with the skin, the skin should be washed immediately and thoroughly with soap and water. If lenalidomide makes contact with the mucous membranes, they should be thoroughly flushed with water.

Healthcare professionals and caregivers should wear disposable gloves when handling the blister or capsule. Gloves should then be removed carefully to prevent skin exposure, placed in a sealable plastic polyethylene bag and disposed of in accordance with local requirements. Hands should then be washed thoroughly with soap and water. Women who are pregnant or suspect they may be pregnant should not handle the blister or capsule (see section 4.4).

Any unused medicinal product or waste material should be returned to the pharmacist for safe disposal in accordance with local requirements.

**7. NAME OF MANUFACTURER**

- a. Penn Pharmaceutical Services Limited.  
Tafarnaubach Industrial Estate,  
Tredegar, Gwent NP22 3AA,  
United Kingdom

- b. Celgene International Sarl  
Route de Perreux 1,  
2017 Boudry,  
Switzerland

**8. DATE OF REVISION OF THE TEXT**

8 November 2022

## PACKAGE LEAFLET: INFORMATION FOR THE PATIENT

**Revlimid®**  
**5 mg, 10 mg, 15 mg, 25 mg**  
**hard capsules**  
lenalidomide

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- **This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.**
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Revlimid® is and what it is used for
2. What you need to know before you take Revlimid®
3. How to take Revlimid®
4. Possible side effects
5. How to store Revlimid®
6. Content of the pack and other information

## **1. WHAT REVLIMID® IS AND WHAT IT IS USED FOR**

### **What Revlimid® is**

Revlimid® contains the active substance 'lenalidomide'. This medicine belongs to a group of medicines which affect how your immune system works.

### **What Revlimid® is used for**

Revlimid® is used in adults for multiple myeloma.

### **Multiple myeloma**

Multiple myeloma is a type of cancer which affects a certain kind of white blood cell, called the plasma cell. These cells collect in the bone marrow and divide, becoming out of control. This can damage the bones and kidneys.

Multiple myeloma generally cannot be cured. However, the signs and symptoms can be greatly reduced or disappear for a period of time. This is called a 'response'.

### **Newly diagnosed multiple myeloma – in patients who have had a bone marrow transplant**

Revlimid® is used on its own as maintenance therapy after patients have recovered enough following a bone marrow transplant.

### **Newly diagnosed multiple myeloma – in patients who cannot have a bone marrow transplant**

Revlimid® is taken with other medicine:

- an anti-inflammatory medicine called 'dexamethasone'.

You will take this other medicine at the start of treatment and then continue to take Revlimid® on its own.

If you are aged 75 years or older or have moderate to severe kidney problems – your doctor will check you carefully before starting treatment.

### **Multiple myeloma – in patients who have had treatment before**

Revlimid® is taken together with an anti-inflammatory medicine called 'dexamethasone'.

Revlimid® can stop the signs and symptoms of multiple myeloma getting worse. It has also been shown to delay multiple myeloma from coming back following treatment.

## How Revlimid® works

Revlimid® works by affecting the body's immune system and directly attacking the cancer. It works in a number of different ways:

- by stopping the cancer cells developing
- by stopping blood vessels growing in the cancer
- by stimulating part of the immune system to attack the cancer cells.

## 2. What you need to know before you take Revlimid®

### DO NOT TAKE Revlimid®:

- if you are pregnant, think you may be pregnant or are planning to become pregnant, **as Revlimid® is expected to be harmful to an unborn child** (see section 2, “Pregnancy, breast-feeding and contraception – information for women and men”).
- if you are able to become pregnant, unless you follow all the necessary measures to prevent you from becoming pregnant (see section 2, “Pregnancy, breast-feeding and contraception – information for women and men”).  
If you are able to become pregnant, your doctor will record with each prescription that the necessary measures have been taken and will provide you with this confirmation.
- if you are allergic to lenalidomide or any of the other ingredients of Revlimid® listed in section 6.  
If you think you may be allergic, ask your doctor for advice.

If any of these apply to you, do not take Revlimid®. Talk to your doctor if you are not sure.

## Warnings and precautions

### Tell your doctor before starting treatment if you have:

- had blood clots in the past - you have an increased risk of developing blood clots in the veins and arteries during treatment
- any signs of an infection, such as a cough or fever
- or have ever had previous viral infection, particularly hepatitis B infection, varicella zoster, HIV. If you are in doubt, talk to your doctor. Treatment with Revlimid® may cause virus to become active again, in patients who carry the virus, resulting in a recurrence of the infection. Your doctor should check whether you have ever had hepatitis B infection.
- kidney problems - your doctor may adjust your dose of Revlimid®
- had a heart attack, have ever had a blood clot, or if you smoke, have high blood pressure or high cholesterol levels
- had an allergic reaction whilst taking thalidomide (another medicine used to treat multiple myeloma) such as rash, itching, swelling, dizziness or trouble breathing
- experienced in the past a combination of any of the following symptoms: rash on face or extended rash, red skin, high fever, flu-like symptoms, enlarged lymph nodes (signs of severe skin reaction called drug reaction with eosinophilia and Systemic Symptoms (DRESS), see also section 4 “Possible side effects”).

If any of the above apply to you, tell your doctor before starting treatment.

At any time during or after your treatment, tell your doctor or nurse immediately if you:

- experience blurred, loss of or double vision, difficulty speaking, weakness in an arm or a leg, a change in the way you walk or problems with your balance, persistent numbness, decreased sensation or loss of sensation, memory loss or confusion. These may all be symptoms of a serious and potentially fatal brain condition known as progressive multifocal leukoencephalopathy (PML). If you had these symptoms prior to treatment with Revlimid, tell your doctor about any change in these symptoms.
- experience shortness of breath, tiredness, dizziness, pain in the chest, a faster heartbeat, or swelling in the legs or ankles. These may be symptoms of a serious condition known as pulmonary hypertension (see section 4).

## Tests and checks

Before and during the treatment with Revlimid® you will have regular blood tests as Revlimid® may cause a fall in the blood cells that help fight infection (white blood cells) and help the blood to clot (platelets). Your doctor will ask you to have a blood test:

- before treatment
- every week for the first 8 weeks of treatment
- then at least every month after that.

You may be evaluated for signs of cardiopulmonary problems before and during the treatment with lenalidomide.

Your doctor may check if you have a high total amount of tumour throughout the body, including your bone marrow. This could lead to a condition where the tumours break down and cause unusual levels of chemicals in the blood which can lead to kidney failure (this condition is called “Tumour Lysis Syndrome”).

Your doctor may check you for changes to your skin such as red spots or rashes.

Your doctor may adjust your dose of Revlimid<sup>®</sup> or stop your treatment based on the results of your blood tests and on your general condition. If you are newly diagnosed, your doctor may also assess your treatment based on your age and other conditions you already have.

### **Blood donation**

You should not donate blood during treatment and for at least 7 days after the end of treatment.

### **Children and adolescents**

Revlimid<sup>®</sup> is not recommended for use in children and adolescents under 18 years.

### **Elderly and people with kidney problems**

If you are aged 75 years or older or have moderate to severe kidney problems - your doctor will check you carefully before starting treatment.

### **Other medicines and Revlimid<sup>®</sup>**

Tell your doctor or nurse if you are taking or have recently taken any other medicines. This is because Revlimid<sup>®</sup> can affect the way some other medicines work. Also, some other medicines can affect the way Revlimid<sup>®</sup> works.

In particular, tell your doctor or nurse if you are taking any of the following medicines:

- some medicines used to prevent pregnancy such as oral contraceptives, as they may stop working
- some medicines used for heart problems – such as digoxin
- some medicines used to thin the blood – such as warfarin

### **Pregnancy, breast-feeding and contraception - information for women and men**

#### **Pregnancy**

##### **For women taking Revlimid<sup>®</sup>**

- You must not take Revlimid<sup>®</sup> if you are pregnant, as it is expected to be harmful to an unborn baby.
- You must not become pregnant while taking Revlimid<sup>®</sup>. Therefore you must use effective methods of contraception if you are a woman of childbearing potential (see “Contraception” below).
- If you do become pregnant during your treatment with Revlimid<sup>®</sup>, you must stop the treatment and inform your doctor immediately.

##### **For men taking Revlimid<sup>®</sup>**

- If your partner becomes pregnant whilst you are taking Revlimid<sup>®</sup>, you should inform your doctor immediately. It is recommended that your partner seeks medical advice.
- You must also use effective methods of contraception (see “Contraception” below).

#### **Breast-feeding**

You must not breast-feed when taking Revlimid<sup>®</sup>, as it is not known if Revlimid<sup>®</sup> passes into human milk.

#### **Contraception**

#### For women taking Revlimid®

Before starting the treatment, you should ask your doctor if you are able to become pregnant, even if you think this is unlikely.

If you are able to become pregnant

- you will have pregnancy tests under the supervision of your doctor (before every treatment, at least every 4 weeks during treatment, and at least 4 weeks after the treatment has finished) except where it has been confirmed that the fallopian tubes have been severed and sealed, to stop eggs from reaching the uterus (tubal sterilisation)

AND

- you must use at least one effective method of contraception for at least 4 weeks before starting treatment, during treatment, and until at least 4 weeks after stopping treatment. Your doctor will advise you on appropriate methods of contraception.

#### For men taking Revlimid®

Revlimid® passes into human semen. If your female partner is pregnant or able to become pregnant, and she does not use effective methods of contraception, you must use condoms during treatment and for at least 7 days after the end of treatment, even if you have had a vasectomy.

You should not donate semen or sperm during treatment (including dose interruption) and for at least 7 days after the end of treatment.

#### **Driving and using machines:**

Do not drive or operate machines if you feel dizzy, tired, sleepy, have vertigo or blurred vision after taking Revlimid®.

#### **Revlimid® contains lactose**

Revlimid® contains lactose. If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicine.

### **3. HOW TO TAKE REVLIMID®**

Revlimid® must be given to you by healthcare professionals with experience in treating multiple myeloma.

- When Revlimid® is used to treat multiple myeloma in patients who cannot have a bone marrow transplant or have had other treatments before, it is taken with dexamethasone (see section 1 'What Revlimid® is used for').
- When Revlimid® is used to treat multiple myeloma in patients who have had a bone marrow transplant, it is taken alone.

Always take Revlimid® exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

If you are taking Revlimid® in combination with dexamethasone, you should refer to the package leaflet of dexamethasone for further information on its use and effects.

#### **Treatment cycle**

Revlimid® is taken on certain days over 4 weeks (28 days).

- Each 28 days is called a 'treatment cycle'.
- Depending on the day of the cycle, you will take one or more of the medicines. However, on some days you do not take any of the medicines.
- After completing each 28-day cycle, you should start a new 'cycle' over the next 28 days.

#### **How much Revlimid® to take**

Before you start treatment, your doctor will tell you:

- how much Revlimid® you should take
- how much of the other medicines you should take in combination with Revlimid®, if any

- on what days of your treatment cycle to take each medicine.


#### **How and when to take Revlimid®**

- Swallow the capsules whole, preferably with water.
- Do not break, open or chew the capsules. If powder from a broken Revlimid® capsule makes contact with the skin, wash the skin immediately and thoroughly with soap and water.
  - Healthcare professionals, caregivers and family members should wear disposable gloves when handling the blister or capsule. Gloves should then be removed carefully to prevent skin exposure, placed in a sealable plastic polyethylene bag and disposed of in accordance with local requirements. Hands should then be washed thoroughly with soap and water. Women who are pregnant or suspect they may be pregnant should not handle the blister or capsule.
- The capsules can be taken either with or without food.
- You should take Revlimid® at about the same time on the scheduled days.

#### **Taking this medicine**

To remove the capsule from the blister:

- press only one end of the capsule out to push it through the foil
- do not put pressure on the centre of the capsule, as this can cause it to break.



#### **Duration of the treatment with Revlimid®**

Revlimid® is taken in treatment cycles, each cycle lasting 28 days (see above “Treatment cycle”). You should continue the cycles of treatment until your doctor tells you to stop.

#### **If you take more Revlimid® than you should**

If you take more Revlimid® than was prescribed, tell your doctor immediately.

#### **If you forget to take Revlimid®**

If you forget to take Revlimid® at your regular time and

- less than 12 hours have passed- take your capsule immediately.
- more than 12 hours have passed- do not take your capsule. Take your next capsule at the usual time the next day.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Revlimid® can cause side effects, although not everybody gets them.

**Stop taking Revlimid and see a doctor straight away if you notice any of the following serious side effects – you may need urgent medical treatment:**



- Hives, rashes, swelling of eyes, mouth or face, difficulty breathing, or itching, which may be symptoms of serious types of allergic reactions called angioedema and anaphylactic reaction.
- A serious allergic reaction that may begin as a rash in one area but spread with extensive loss of skin over the whole body (Stevens-Johnson syndrome and/or toxic epidermal necrolysis).
- Widespread rash, high body temperature, liver enzyme elevations, blood abnormalities (eosinophilia), enlarged lymph nodes and other body organs involvement (Drug Reaction with Eosinophilia and Systemic Symptoms which is also known as DRESS or drug hypersensitivity syndrome). See also section 2.

**Tell your doctor straight away if you notice any of the following serious side effects:**

- fever, chills, sore throat, cough, mouth ulcers or any other symptoms of infection including within the bloodstream (sepsis)
- bleeding or bruising in the absence of injury
- chest pain or leg pain
- shortness of breath
- Bone pain, muscle weakness, confusion or tiredness that might be due to high level of calcium in the blood.

Revlimid may reduce the number of white blood cells that fight infection and also the blood cells which help the blood to clot (platelets) which may lead to bleeding disorders such as nosebleeds and bruising.

Revlimid may also cause blood clots in the veins (thrombosis).

**Other side effects**

It is important to note that a small number of patients may develop additional types of cancer, and it is possible that this risk may be increased with Revlimid® treatment, therefore your doctor should carefully evaluate the benefit and risk when you are prescribed Revlimid®.

**Very common** side effects (may affect more than 1 in 10 people):

- A fall in the number of red blood cells which may cause anemia leading to tiredness and weakness
- Constipation, diarrhoea, nausea, redness of skin, rashes, vomiting, muscle cramps, muscle aches, bone pain, joint pain, tiredness, generalised swelling including swelling of your arms and legs
- Fever and flu like symptoms including fever, muscle ache, headache, earache and chills
- Numbness, tingling or burning sensation to the skin, pains in hands or feet, dizziness, tremor, changes in the way things taste
- Chest pain spreading to the arms, neck, jaw, back or stomach, feeling sweaty and breathless, feeling sick or vomiting, which may be symptoms of a heart attack (myocardial infarction)
- Decreased appetite
- Low levels of potassium in the blood
- Leg pain (which could be a symptom of thrombosis), chest pain or shortness of breath (which may be a symptom of blood clots in the lungs, called pulmonary embolism)
- Infections of all types
- Infection of the lung and the upper respiratory tract, shortness of breath
- Blurred vision
- Clouding of your eye (cataract)
- Kidney problems
- Changes to a protein in the blood that can cause swelling of the arteries (vasculitis)
- Increases in your blood sugar level (diabetes)
- Headache
- Dry skin
- Stomach pain
- Mood change, difficulty sleeping

**Common** side effects (may affect up to 1 in 10 people):

- Infection of the sinuses that surround the nose
- Bleeding from the gums, stomach, or bowels
- Increase in pain, tumour size, redness around the tumour
- Increased blood pressure or a fall in blood pressure, slow, fast or irregular heart beat
- Darkening of your skin
- Skin eruptions, skin cracking, flaking or peeling skin
- Hives, itching, increased sweating, dehydration
- Sore inflamed mouth, dry mouth, difficulty swallowing
- Heartburn
- Production of much more or much less urine than usual (which may be a symptom of kidney failure), passing blood in the urine
- Shortness of breath especially when lying down (which may be a symptom of heart failure)
- Difficulty getting an erection
- Stroke, fainting
- Muscle weakness
- Joint swelling
- Changes to blood thyroid hormone, low levels of calcium, phosphate or magnesium in the blood
- Depression
- Deafness
- Abnormal liver test results
- Impaired balance, difficulty moving
- Ringing in the ears (tinnitus)
- An excess of iron in the body
- Thirst
- Confusion
- Toothache
- Weight loss

**Uncommon** side effects (may affect up to 1 in 100 people):

- Bleeding within the skull
- Circulatory problems
- Loss of vision
- Loss of sex drive (libido)
- Passing large amount of urine with bone pain and weakness, which may be symptoms of a kidney disorder (Fanconi syndrome)
- Yellow pigmentation to the skin, mucus membrane or eyes (jaundice), pale coloured stools, dark coloured urine, skin itch, rash, pain or swelling of the stomach—these may be symptoms of injury to the liver (hepatic failure).
- Stomach pain, bloating, or diarrhoea, which may be symptoms of inflammation in the large intestine (called colitis or caecitis)
- Damage to the cells of the kidney (called renal tubular necrosis)
- Changes to the colour of your skin, sensitivity to sunlight
- Certain types of skin tumour
- Tumour lysis syndrome - metabolic complications that can occur during treatment of cancer and sometimes even without treatment. These complications are caused by the break-down products of dying cancer cells and may include the following: changes to blood chemistry; high potassium, phosphorus, uric acid, and low calcium consequently leading to changes in kidney function, heart beat, seizures, and sometimes death.
- Increase in blood pressure within blood vessels that supply the lungs (pulmonary hypertension).

**Not known** (frequency cannot be estimated from the available data):

- Sudden, or mild but worsening pain in the upper stomach and/or back, which remains for a few days, possibly accompanied by nausea, vomiting, fever and a rapid pulse. These symptoms may be due to inflammation of the pancreas.
- Wheezing, shortness of breath or a dry cough, which may be symptoms caused by inflammation of the

tissue in the lungs.

- Rare cases of muscle breakdown (muscle pain, weakness or swelling) which can lead to kidney problems (rhabdomyolysis) have been observed, some of them when Revlimid® is administered with a statin (a type of cholesterol lowering medication).
- A condition affecting the skin caused by inflammation of small blood vessels, along with pain in the joints and fever (leukocytoclastic vasculitis).
- Breakdown of the wall of the stomach or gut. This may lead to very serious infection. Tell your doctor if you have severe stomach pain, fever, nausea, vomiting, blood in your stool, or changes in bowel habits.
- Viral infections, including herpes zoster (also known as ‘shingles’, a viral disease that causes a painful skin rash with blisters) and recurrence of hepatitis B infection (which can cause yellowing of the skin and eyes, dark brown-colored urine, right-sided stomach pain, fever and feeling nauseous or being sick).
- Rejection of solid organ transplant (such as kidney, heart).

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

## **5. HOW TO STORE REVLIMID®**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date, which is stated on the blister and on the carton after “EXP”. The expiry date refers to the last day of that month.
- Do not use this medicine if you notice any damage or signs of tampering to the pack.
- Do not throw away any medicines via wastewater or household waste. All unused Revlimid® capsules should be returned to the pharmacist. These measures will help protect the environment.

## **6. CONTENTS OF THE PACK AND OTHER INFORMATION**

### **What Revlimid® contains**

Revlimid® 5 mg hard capsules:

- The active substance is lenalidomide. Each capsule contains 5 mg of lenalidomide.
- The other ingredients are:
  - capsule contents: anhydrous lactose (see section 2 of the package insert), microcrystalline cellulose, croscarmellose sodium and magnesium stearate
  - capsule shell: gelatine and titanium dioxide (E171)
  - printing ink: shellac, propylene glycol (E1520), potassium hydroxide and black iron oxide (E172).

Revlimid® 10 mg hard capsules:

- The active substance is lenalidomide. Each capsule contains 10 mg of lenalidomide.
- The other ingredients are:
  - capsule contents: anhydrous lactose (see section 2 of the package insert), microcrystalline cellulose, croscarmellose sodium and magnesium stearate
  - capsule shell: gelatine, titanium dioxide (E171), indigo carmine (E132) and yellow iron oxide (E172)
  - printing ink: shellac, propylene glycol (E1520), potassium hydroxide and black iron oxide (E172).

Revlimid® 15 mg hard capsules:

- The active substance is lenalidomide. Each capsule contains 15 mg of lenalidomide.
- The other ingredients are:
  - capsule contents: anhydrous lactose (see section 2 of the package insert), microcrystalline cellulose, croscarmellose sodium and magnesium stearate
  - capsule shell: gelatine, titanium dioxide (E171) and indigo carmine (E132)
  - printing ink: shellac, propylene glycol (E1520), potassium hydroxide and black iron oxide (E172).

Revlimid® 25 mg hard capsules:

- The active substance is lenalidomide. Each capsule contains 25 mg of lenalidomide.
- The other ingredients are:
  - capsule contents: anhydrous lactose (see section 2 of the package insert), microcrystalline cellulose, croscarmellose sodium and magnesium stearate
  - capsule shell: gelatine and titanium dioxide (E171)
  - printing ink: shellac, propylene glycol (E1520), potassium hydroxide and black iron oxide (E172).

**What Revlimid® looks like and contents of the pack**

The capsules are provided in packs. Each pack contains three blisters, each blister with seven capsules. This gives a total of 21 capsules per pack.

This leaflet was last revised on 8 November 2022