

1. NAME OF THE MEDICINAL PRODUCT

TEROPROS FILM COATED TABLET 1000MG
TEROPROS FILM COATED TABLET 500MG
TEROPROS FILM COATED TABLET 250MG

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

TEROPROS FILM COATED TABLET 1000MG

Each film-coated tablet contains 1000 mg of abiraterone acetate.

Excipients with known effect: Each film-coated tablet contains 136 mg of lactose monohydrate and 21.24 mg of sodium.

TEROPROS FILM COATED TABLET 500MG

Each film-coated tablet contains 500 mg of abiraterone acetate.

Excipients with known effect: Each film-coated tablet contains 68 mg of lactose monohydrate and 10.6 mg of sodium.

TEROPROS FILM COATED TABLET 250MG

Each film-coated tablet contains 250 mg of abiraterone acetate.

Excipients with known effect: Each film-coated tablet contains 34 mg of lactose monohydrate and 5.3 mg of sodium.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet.

TEROPROS FILM COATED TABLET 250MG: White to off-white, oval-shaped film-coated tablets, debossed with “250” on one side, with dimensions of 14.2 mm x 7.2 mm.

TEROPROS FILM COATED TABLET 500MG: Purple, oval-shaped film-coated tablets, debossed with “500” on one side, with dimensions of 18.9 mm x 9.5 mm.

TEROPROS FILM COATED TABLET 1000MG: White to off-white, oval-shaped film-coated tablets, with a break line on one side and plain on the other side, with dimensions 23.1 mm x 11.1 mm. The break line only serves to facilitate breaking for ease of swallowing and does not divide the tablet into equal half-doses.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

TEROPROS is indicated with prednisone or prednisolone for:

- The treatment of newly diagnosed high risk metastatic hormone sensitive prostate cancer (mHSPC) in adult men in combination with androgen deprivation therapy (ADT).
- the treatment of metastatic castration resistant prostate cancer (mCRPC) in adult men who are asymptomatic or mildly symptomatic after failure of androgen deprivation therapy in whom chemotherapy is not yet clinically indicated (see section 5.1).
- the treatment of metastatic castration resistant prostate cancer (mCRPC) in adult men whose disease has progressed on or after a docetaxel-based chemotherapy regimen.

4.2 Posology and method of administration

Dosage

The recommended dosage is 1000 mg as a single daily dose that must not be taken with food.

Abiraterone acetate must be taken on an empty stomach, at least one hour before or at least two hours after a meal. Taking the tablets with food increases systemic exposure to abiraterone. The tablets should be swallowed with water (see section 5.2 – Absorption). If splitting the 1000 mg tablet is

necessary to facilitate swallowing, break the tablet using a dedicated pill cutter, only at the moment of administration. Swallow both generated halves of the tablet consecutively at the same moment without chewing.

Medical castration with LHRH analogue should be continued during treatment in patients not surgically castrated.

Dosage of prednisone or prednisolone

For metastatic hormone sensitive prostate cancer (mHSPC), abiraterone acetate is used with 5 mg prednisone or prednisolone daily.

For metastatic castration-resistant prostate cancer (mCRPC), abiraterone acetate is used with 10 mg prednisone or prednisolone daily.

Recommended monitoring

Serum transaminases should be measured prior to starting treatment, every two weeks for the first three months of treatment and monthly thereafter. Blood pressure, serum potassium and fluid retention should be monitored monthly. However, patients with a significant risk for congestive heart failure should be monitored every 2 weeks for the first three months of treatment and monthly thereafter (see section 4.4 – Hypertension, hypokalemia, fluid retention and cardiac failure due to mineralocorticoid excess and Hepatotoxicity and Hepatic impairment).

In patients with pre-existing hypokalemia or those that develop hypokalemia whilst being treated with abiraterone acetate, consider maintaining the patient's potassium level at ≥ 4.0 mM.

For patients who develop Grade ≥ 3 toxicities including hypertension, hypokalemia, oedema and other non-mineralocorticoid toxicities, treatment should be withheld and appropriate medical management should be instituted. Treatment with abiraterone acetate should not be reinitiated until symptoms of the toxicity have resolved to Grade 1 or baseline.

In the event of a missed daily dose of either abiraterone acetate, prednisone or prednisolone, treatment should be resumed the following day with the usual daily dose.

Hepatic impairment

No dose adjustment is necessary for patients with pre-existing mild hepatic impairment, Child-Pugh Class A. There are no data on the clinical safety and efficacy of multiple doses of abiraterone acetate when administered to patients with moderate or severe hepatic impairment (Child-Pugh Class B or C). No dose adjustment can be predicted. abiraterone acetate should be used with caution in patients with moderate hepatic impairment, only if the benefit clearly outweighs the possible risk (see section 4.4 – Hepatotoxicity and Hepatic impairment and 5.2 – Special populations). Abiraterone acetate should not be used in patients with severe hepatic impairment (see section 4.4 – Hepatotoxicity and Hepatic impairment and 5.2 – Special populations).

For patients who develop hepatotoxicity during treatment (alanine aminotransferase (ALT) increases or aspartate aminotransferase (AST) increases above 5 times the upper limit of normal, treatment should be withheld immediately until liver function tests normalize (see section 4.4 – Hepatotoxicity and Hepatic impairment). Re-treatment following return of liver function tests to the patient's baseline may be given at a reduced dose of 500 mg once daily. For patients being re-treated, serum transaminases should be monitored at a minimum of every two weeks for three months and monthly thereafter. If hepatotoxicity recurs at the reduced dose of 500 mg daily, treatment should be discontinued. Reduced doses should not be taken with food (see section 4.2 – Dosage).

If patients develop severe hepatotoxicity (ALT or AST 20 times the upper limit of normal) anytime while on therapy, treatment should be discontinued and patients should not be re-treated with abiraterone acetate.

Moderate hepatic impairment (Child-Pugh Class B) has been shown to increase the systemic exposure to abiraterone by approximately four-fold following single oral doses of abiraterone acetate 1,000 mg

(see section 5.2).

Renal impairment

No dosage adjustment is necessary for patients with renal impairment (see section 5.2 – Special populations). However, there is no clinical experience in patients with prostate cancer and severe renal impairment. Caution is advised in these patients (see section 4.4).

Paediatric population

There is no relevant use of this medicinal product in the paediatric population, as prostate cancer is not present in children and adolescents.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients (see section 6.1).
- Women who are or may potentially be pregnant (see section 4.6 – Pregnancy)
- Severe hepatic impairment [Child-Pugh Class C (see sections 4.2, 4.4 and 5.2)].

4.4 Special warnings and precautions for use

Hypertension, hypokalemia and fluid retention due to mineralocorticoid excess

Abiraterone acetate may cause hypertension, hypokalemia and fluid retention (see section 4.8) as a consequence of increased mineralocorticoid levels resulting from CYP17 inhibition (see section 5 – Mechanism of action). Co-administration of a corticosteroid suppresses adrenocorticotrophic hormone (ACTH) drive, resulting in a reduction in incidence and severity of these adverse reactions. Caution is required in treating patients whose underlying medical conditions might be compromised by increases in blood pressure, hypokalemia (e.g., those on cardiac glycosides), or fluid retention (e.g., those with heart failure), severe or unstable angina pectoris, recent myocardial infarction or ventricular arrhythmia and those with severe renal impairment. In postmarketing experience, QT prolongation and Torsades de Pointes have been observed in patients who develop hypokalemia or have underlying cardiovascular conditions while taking abiraterone acetate.

Abiraterone acetate should be used with caution in patients with a history of cardiovascular disease. The phase 3 studies conducted with abiraterone acetate excluded patients with uncontrolled hypertension, clinically significant heart disease as evidenced by myocardial infarction, or arterial thrombotic events in the past 6 months, severe or unstable angina, or New York Heart Association (NYHA) Class III or IV heart failure (Study 301) or Class II to IV heart failure (Study 302) or cardiac ejection fraction measurement of < 50%. In Study 302 patients with atrial fibrillation, or other cardiac arrhythmia requiring medical therapy were excluded. Safety in patients with left ventricular ejection fraction (LVEF) < 50% or NYHA Class III or IV heart failure (in Study 301) or NYHA Class II to IV heart failure (in Studies 301 and 302) was not established (see sections 4.8 and 5).

Before treating patients with a significant risk for congestive heart failure (e.g. a history of cardiac failure, uncontrolled hypertension, or cardiac events such as ischaemic heart disease), consider obtaining an assessment of cardiac function (e.g. echocardiogram). Before treatment with abiraterone acetate, cardiac failure should be treated and cardiac function optimised. Hypertension, hypokalemia and fluid retention should be corrected and controlled. During treatment, blood pressure, serum potassium, fluid retention (weight gain, peripheral oedema), and other signs and symptoms of congestive heart failure should be monitored every 2 weeks for 3 months, then monthly thereafter and abnormalities corrected. Assess cardiac function as clinically indicated, institute appropriate management and consider discontinuation of abiraterone acetate treatment if there is a clinically significant decrease in cardiac function (see section 4.2).

Hepatotoxicity and hepatic Impairment

Marked increases in liver enzymes leading to treatment discontinuation or dose modification occurred in controlled clinical studies (see section 4.8). Serum transaminase and bilirubin levels

should be measured prior to starting treatment, every two weeks for the first three months of treatment, and monthly thereafter. If clinical symptoms or signs suggestive of hepatotoxicity develop, serum transaminases should be measured immediately. If at any time the ALT or AST rises above 5 times the upper limit of normal or the bilirubin rises above 3 times the upper limit of normal, treatment should be interrupted immediately and liver function closely monitored.

Re-treatment may only take place after the return of liver function tests to the patient's baseline and at a reduced dose level (see section 4.2 – Hepatic impairment).

If patients develop severe hepatotoxicity (ALT or AST 20 times the upper limit of normal) anytime while on therapy, treatment should be discontinued and patients should not be re-treated.

Patients with active or symptomatic viral hepatitis were excluded from clinical trials; thus, there are no data to support the use of abiraterone acetate in this population.

There are no data on the clinical safety and efficacy of multiple doses of abiraterone acetate when administered to patients with moderate or severe hepatic impairment (Child-Pugh Class B or C). No dose adjustment can be predicted. abiraterone acetate should be used with caution in patients with moderate hepatic impairment only if the benefit clearly outweighs the possible risk (see section 4.2 – Hepatic impairment and 5.2 – Special populations). Abiraterone acetate should not be used in patients with severe hepatic impairment (see section 4.2 – Hepatic impairment, 4.3 and 5.2 – Special populations).

There have been rare post-marketing reports of acute liver failure and hepatitis fulminant, some with fatal outcome (see section 4.8).

Corticosteroid withdrawal and coverage of stress situations

Caution is advised and monitoring for adrenocortical insufficiency should occur if patients are withdrawn from prednisone or prednisolone. If abiraterone acetate is continued after corticosteroids are withdrawn, patients should be monitored for symptoms of mineralocorticoid excess (see section 4.4 – Hypertension, hypokalemia and fluid retention due to mineralocorticoid excess).

In patients on prednisone or prednisolone who are subjected to unusual stress, an increased dose of corticosteroids may be indicated before, during and after the stressful situation.

Bone density

Decreased bone density may occur in men with metastatic advanced prostate cancer (castration resistant prostate cancer). The use of abiraterone acetate in combination with a glucocorticoid could increase this effect.

Prior use of ketoconazole

Lower rates of response might be expected in patients previously treated with ketoconazole for prostate cancer.

Hyperglycaemia

The use of glucocorticoids could increase hyperglycaemia, therefore blood sugar should be measured frequently in patients with diabetes.

Hypoglycaemia

Isolated cases of hypoglycaemia have been reported when abiraterone acetate plus prednisone/prednisolone was administered to patients with pre-existing diabetes receiving pioglitazone or repaglinide (see section 4.5 Interaction with other medicinal products and other forms of interaction – Interactions with other drugs – Potential for abiraterone acetate to affect exposures to other drugs). Blood glucose should be monitored in patients with diabetes.

Use with chemotherapy

The safety and efficacy of concomitant use of abiraterone acetate with cytotoxic chemotherapy has not been established (see section 5 – Clinical studies).

Use in combination with radium 223 dichloride

In a randomized clinical trial in patients with asymptomatic or mildly symptomatic bone-predominant metastatic castration resistant prostate cancer, at the time of unblinding, the addition of radium 223 dichloride to abiraterone acetate plus prednisone/prednisolone showed an increase in mortality and an increased rate of fracture. Radium 223 dichloride is not recommended for use in combination with abiraterone acetate plus prednisone/prednisolone outside of clinical trials.

Intolerance to excipients

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Potential risks

Anaemia and sexual dysfunction may occur in men with metastatic castration resistant prostate cancer including those undergoing treatment with abiraterone acetate.

4.5 Interaction with other medicinal products and other forms of interaction

Effect of food on abiraterone acetate

Administration with food significantly increases the absorption of abiraterone acetate. The efficacy and safety of abiraterone acetate given with food have not been established. Abiraterone acetate must not be taken with food (see section 4.2 and 5.2 – Absorption).

Interactions with other drugs

Potential for other drugs to affect abiraterone exposures

Based on in vitro data, abiraterone is a substrate of CYP3A4.

In a clinical pharmacokinetic interaction study of healthy subjects pretreated with a strong CYP3A4 inducer (rifampicin, 600 mg daily for 6 days) followed by a single dose of abiraterone acetate 1000 mg, the mean plasma AUC_{∞} of abiraterone was decreased by 55%.

Strong inducers of CYP3A4 (e.g., phenytoin, carbamazepine, rifampicin, rifabutin, rifapentine, phenobarbital) during treatment with abiraterone acetate are to be avoided, or used with careful evaluation of clinical efficacy.

In a separate clinical pharmacokinetic interaction study of healthy subjects, co-administration of ketoconazole, a strong inhibitor of CYP3A4, had no clinically meaningful effect on the pharmacokinetics of abiraterone.

Potential for abiraterone acetate to affect exposures to other drugs

Abiraterone is an inhibitor of the hepatic drug-metabolizing enzymes CYP2D6 and CYP2C8.

In a clinical study to determine the effects of abiraterone acetate (plus prednisone) on a single dose of the CYP2D6 substrate dextromethorphan, the systemic exposure (AUC) of dextromethorphan was increased approximately 2.9 fold. The AUC_{24} for dextromethorphan, the active metabolite of dextromethorphan, increased approximately 33%.

Caution is advised when abiraterone acetate is administered with drugs activated by or metabolized by CYP2D6, particularly with drugs that have a narrow therapeutic index. Dose reduction of narrow therapeutic index drugs metabolized by CYP2D6 should be considered. Examples of medicinal products metabolised by CYP2D6 include metoprolol, propranolol, desipramine, venlafaxine, haloperidol, risperidone, propafenone, flecanide, codeine, oxycodone and tramadol (the latter three products requiring CYP2D6 to form their active analgesic metabolites).

In vitro, abiraterone was shown to inhibit the hepatic drug-metabolizing enzyme CYP1A2. However, in a clinical study to determine the effects of abiraterone acetate (plus prednisone) on a single dose of the CYP1A2 substrate theophylline, no increase in systemic exposure of theophylline was observed.

In a CYP2C8 drug-drug interaction trial in healthy subjects, the AUC of pioglitazone was increased by 46% and the AUCs for M-III and M-IV, the active metabolites of pioglitazone, each decreased by 10%, when pioglitazone was given together with a single dose of 1000 mg abiraterone acetate. Patients should be monitored for signs of toxicity related to a CYP2C8 substrate with a narrow therapeutic index if used concomitantly with abiraterone acetate. Examples of medicinal products metabolised by CYP2C8 include pioglitazone and repaglinide.

4.6 Fertility, pregnancy and lactation

Pregnancy

Abiraterone acetate is not for use in women.

Abiraterone acetate is contraindicated in women who are or may potentially be pregnant (see section 4.3).

There are no human data on the use of abiraterone acetate in pregnancy and abiraterone acetate is not for use in women of child-bearing potential. Maternal use of a CYP17 inhibitor is expected to produce changes in hormone levels that could affect development of the fetus (see section 5 – Mechanism of action and 5.3 – Reproductive Toxicology).

It is not known whether abiraterone or its metabolites are present in semen. A condom is required if the patient is engaged in sexual activity with a pregnant woman. If the patient is engaged in sex with a woman of childbearing potential, a condom is required along with another effective contraceptive method.

Based on its mechanism of action, abiraterone acetate may harm a developing fetus; therefore, to avoid inadvertent exposure women who are pregnant or women whom may be pregnant should not handle abiraterone acetate without protection, e.g., gloves.

Breast-feeding

Abiraterone acetate is not for use in women.

It is not known if either abiraterone acetate or its metabolites are excreted in human breast milk.

Fertility

Abiraterone affected fertility in male and female rats, but these effects were fully reversible (see section 5.3).

4.7 Effects on ability to drive and use machines

Abiraterone acetate has no or negligible influence on the ability to drive or use machines.

4.8 Undesirable effects

Throughout this section, adverse reactions are presented. Adverse reactions are adverse events that were considered to be reasonably associated with the use of abiraterone acetate based on the comprehensive assessment of the available adverse event information. A causal relationship with abiraterone acetate usually cannot be reliably established in individual cases. Further, because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

In an analysis of adverse reactions of composite Phase 3 studies with abiraterone acetate, adverse reactions that were observed in $\geq 10\%$ of patients were hypertension, peripheral edema, hypokalemia, urinary tract infection, and aspartate aminotransferase increased and/or alanine aminotransferase

increased.

Abiraterone acetate may cause hypertension, hypokalemia and fluid retention as a pharmacodynamic consequence of its mechanism of action. In Phase 3 studies, anticipated mineralocorticoid adverse reactions were seen more commonly in patients treated with abiraterone acetate than in patients treated with placebo: hypokalemia 18% versus 8%, hypertension 22% versus 16% and fluid retention (peripheral oedema) 23% versus 17%, respectively. In patients treated with abiraterone acetate, CTCAE (version 3.0) Grades 3 and 4 hypokalemia were observed in 6% and 2% of patients, CTCAE (version 3.0) Grades 3 and 4 hypertension were observed in 8% and 5% of patients, and grades 3 and 4 fluid retention edema were observed in 1% and 1% of patients, respectively. Mineralocorticoid reactions generally were able to be successfully managed medically. Concomitant use of a corticosteroid reduces the incidence and severity of these adverse reactions (see section 4.4 – Hypertension, hypokalemia and fluid retention due to mineralocorticoid excess).

In studies of patients with metastatic advanced prostate cancer who were using a LHRH agonist, or were previously treated with orchiectomy, abiraterone acetate was administered at a dose of 1,000 mg daily in combination with low dose prednisone or prednisolone (5 or 10 mg daily).

Adverse reactions that occurred at a rate of $\geq 1\%$ (all grades) are shown in Table 1.

Table 1: Adverse Reactions in $\geq 1\%$ of Patients in Clinical Studies^a

| | Abiraterone acetate 1000 mg daily with prednisone or prednisolone n=2659 ^b | | |
|---|--|--------------|--------------|
| System Organ Class Adverse Reaction | All grades % | Grade 3 % | Grade 4 % |
| General Disorders and Administration Site Conditions | | | |
| Edema peripheral | 20 | <1 | 0 |
| Metabolism and Nutrition Disorders | | | |
| Hypokalemia | 20 | 5 | <1 |
| Hypertriglyceridemia | 1 | <1 | 0 |
| Infections and Infestations | | | |
| Urinary tract infection | 10 | 2 | <1 |
| Hepatobiliary Disorders | | | |
| ALT increased and/or AST increased ^c | 13 | 4 | <1 |
| Vascular Disorders | | | |
| Hypertension | 21 | 6 | 0 |
| Injury, Poisoning and Procedural Complications | | | |
| Fractures ^d | 7 | 2 | <1 |
| Cardiac Disorders | | | |
| Cardiac failure ^e | 1 | <1 | <1 |
| Angina pectoris | 2 | <1 | 0 |
| Arrhythmia | 1 | 0 | 0 |
| Atrial fibrillation | 3 | 1 | <1 |
| Tachycardia | 2 | <1 | 0 |
| Renal and Urinary Disorders | | | |
| Hematuria | 7 | 1 | 0 |
| Gastrointestinal Disorders | | | |
| Dyspepsia | 6 | 0 | 0 |

^a All patients were using an LHRH agonist or had undergone orchiectomy.

^b n=patients assessed for safety.

^c ALT increased and/or AST increased includes ALT increased, AST increased, and hepatic function abnormal.

^d Fractures includes osteoporosis and all fractures with the exception of pathological fracture.

^e Cardiac failure includes congestive heart failure, left ventricular dysfunction and ejection fraction

decreased.

The adverse reaction, adrenal insufficiency, occurred in phase 3 clinical studies at a rate of 0.3% in patients taking abiraterone acetate and at a rate of 0.1% in patients taking placebo.

Cardiovascular effects

The three phase 3 studies excluded patients with uncontrolled hypertension, clinically significant heart disease as evidenced by myocardial infarction, or arterial thrombotic events in the past 6 months, severe or unstable angina, or NYHA Class III or IV heart failure (Study 301) or Class II to IV heart failure (Studies 3011 and 302) or cardiac ejection fraction measurement of < 50%. All patients enrolled (both active and placebo-treated patients) were concomitantly treated with androgen deprivation therapy, predominantly with the use of LHRH agonists, which has been associated with diabetes, myocardial infarction, cerebrovascular accident and sudden cardiac death. The incidence of cardiovascular adverse reactions in the phase 3 studies in patients taking abiraterone acetate versus patients taking placebo were as follows: hypertension 14.5% vs. 10.5%, atrial fibrillation 2.6% vs. 2.0%, tachycardia 1.9% vs. 1.0%, angina pectoris 1.7% vs. 0.8%, cardiac failure 0.7% vs. 0.2% and arrhythmia 0.7% vs. 0.5%.

Hepatotoxicity

Drug-associated hepatotoxicity with elevated ALT, AST and total bilirubin has been reported in patients treated with abiraterone acetate. Across Phase 3 clinical studies, hepatotoxicity grades 3 and 4 (e.g., ALT or AST increases of > 5 x ULN or bilirubin increases > 1.5 x ULN) were reported in approximately 6% of patients who received abiraterone acetate, typically during the first 3 months after starting treatment. In Study 3011, grade 3 or 4 hepatotoxicity was observed in 8.4% of patients treated with abiraterone acetate. Ten patients who received abiraterone acetate were discontinued because of hepatotoxicity; two had Grade 2 hepatotoxicity, six had Grade 3 hepatotoxicity, and two had Grade 4 hepatotoxicity. No patient died of hepatotoxicity in Study 3011. In the Phase 3 clinical studies, patients whose baseline ALT or AST were elevated were more likely to experience liver function test elevations than those beginning with normal values. When elevations of either ALT or AST > 5 x ULN, or elevations in bilirubin > 3 x ULN were observed, abiraterone acetate was withheld or discontinued. In two instances marked increases in liver function tests occurred (see section 4.4). These two patients with normal baseline hepatic function, experienced ALT or AST elevations 15 to 40 x ULN and bilirubin elevations 2 to 6 x ULN. Upon discontinuation of abiraterone acetate, both patients had normalisation of their liver function tests and one patient was re-treated without recurrence of the elevations. In Study 302, grade 3 or 4 ALT or AST elevations were observed in 35 (6.5%) patients treated with abiraterone acetate. Aminotransferase elevations resolved in all but 3 patients (2 with new multiple liver metastases and 1 with AST elevation approximately 3 weeks after the last dose of abiraterone acetate). In Phase 3 clinical studies, treatment discontinuations due to ALT and AST increases or abnormal hepatic function were reported in 1.1% of patients treated with abiraterone acetate and 0.6% of patients treated with placebo; no deaths were reported due to hepatotoxicity events.

In clinical trials, the risk for hepatotoxicity was mitigated by exclusion of patients with baseline hepatitis or significant abnormalities of liver function tests. In the 3011 trial, patients with baseline ALT and AST > 2.5 X ULN, bilirubin > 1.5 X ULN or those with active or symptomatic viral hepatitis or chronic liver disease; ascites or bleeding disorders secondary to hepatic dysfunction were excluded. In the 301 trial, patients with baseline ALT and AST \geq 2.5 x ULN in the absence of liver metastases and > 5 x ULN in the presence of liver metastases were excluded. In the 302 trial, patients with liver metastases were not eligible and patients with baseline ALT and AST \geq 2.5 x ULN were excluded. Abnormal liver function tests developing in patients participating in clinical trials were vigorously managed by requiring treatment interruption and permitting re-treatment only after return of liver function tests to the patient's baseline (see section 4.2 – Hepatic impairment). Patients with elevations of ALT or AST > 20 x ULN were not re-treated. The safety of re-treatment in such patients is unknown. The mechanism for hepatotoxicity is not understood.

Post-marketing experience

Adverse reactions identified during the post-marketing experience based on spontaneous reports with abiraterone acetate are described below. The frequencies are provided according to the following convention:

| | |
|-----------|-------------------------------|
| Uncommon | $\geq 1/1000$ and $< 1/100$ |
| Rare | $\geq 1/10000$ and $< 1/1000$ |
| Very Rare | $< 1/10000$ |

System Organ Class: Respiratory, thoracic and mediastinal disorders

Rare: Allergic alveolitis

System Organ Class: Musculoskeletal and connective tissue disorders

Uncommon: Rhabdomyolysis, Myopathy

System Organ Class: Hepatobiliary disorders

Rare: Hepatitis fulminant, Acute hepatic failure

System Organ Class: Cardiac disorders

Very rare: QT prolongation and Torsades de Pointes (observed in patients who developed hypokalemia or had underlying cardiovascular conditions).

System Organ Class: Immune System Disorders – Hypersensitivity

Very rare: Anaphylactic reaction (severe allergic reactions that include, but are not limited to difficulty swallowing or breathing, swollen face, lips, tongue or throat, or an itchy rash (urticaria)).

4.9 Overdose

Human experience of overdose with abiraterone acetate is limited.

There is no specific antidote. In the event of an overdose, administration should be withheld and general supportive measures undertaken, including monitoring for arrhythmias, hypokalemia and for signs and symptoms of fluid retention. Liver function also should be assessed.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: endocrine therapy, other hormone antagonists and related agents, ATC code: L02BX03

Mechanism of action

Abiraterone acetate is converted *in vivo* to abiraterone, an androgen biosynthesis inhibitor. Specifically, abiraterone selectively inhibits the enzyme 17 α -hydroxylase/C17,20-lyase (CYP17). This enzyme is expressed in and is required for androgen biosynthesis in testicular, adrenal and prostatic tumour tissues. CYP17 catalyses the conversion of pregnenolone and progesterone into testosterone precursors, DHEA and androstenedione, respectively, by 17 α -hydroxylation and cleavage of the C17,20 bond. CYP17 inhibition also results in increased mineralocorticoid production by the adrenals (see section 4.4 – Hypertension, hypokalemia and fluid retention due to mineralocorticoid excess).

Androgen-sensitive prostatic carcinoma responds to treatment that decreases androgen levels.

Androgen deprivation therapies, such as treatment with LHRH agonists or orchiectomy, decrease androgen production in the testes but do not affect androgen production by the adrenals or in the tumour. Treatment with abiraterone acetate decreases serum testosterone to undetectable levels (using commercial assays) when given with LHRH agonists (or orchiectomy).

Pharmacodynamic effects

Abiraterone acetate decreases serum testosterone and other androgens to levels lower than those achieved by the use of LHRH agonists alone or by orchiectomy. This results from the selective inhibition of the CYP17 enzyme required for androgen biosynthesis. Prostate specific antigen (PSA) serves as a biomarker in patients with prostate cancer. In a Phase 3 clinical study of patients who failed prior chemotherapy with taxanes, 38% of patients treated with abiraterone acetate, versus 10% of patients treated with placebo, had at least a 50% decline from baseline in PSA levels.

Use of Spironolactone

Patients in pivotal clinical trials with abiraterone acetate were not allowed to use spironolactone as spironolactone binds to the androgen receptor and may increase PSA levels.

Clinical studies

The efficacy of abiraterone acetate was established in three randomised placebo-controlled multicenter phase 3 clinical studies (Studies 3011, 302, and 301) of patients with hormone sensitive metastatic prostate cancer and metastatic castration-resistant prostate cancer.

Study 3011 enrolled patients who were newly diagnosed (within 3 months of randomization) mHNPC who had high-risk prognostic factors. High-risk prognosis was defined as having at least 2 of the following 3 risk factors: (1) Gleason score of ≥ 8 ; (2) presence of 3 or more lesions on bone scan; (3) presence of measurable visceral (excluding lymph node disease) metastasis. In the active arm, abiraterone acetate was administered at a dose of 1000 mg daily in combination with low dose prednisone or prednisolone 5 mg once daily in addition to ADT (LHRH agonist or orchiectomy), which was the standard of care treatment. Patients in the control arm received ADT and placebos for both abiraterone acetate and prednisone.

Study 302 enrolled docetaxel naïve patients, whereas Study 301 enrolled patients who had received prior docetaxel. In both studies, patients were using an LHRH agonist or were previously treated with orchiectomy. In the active treatment arm, abiraterone acetate was administered at a dose of 1,000 mg daily in combination with low dose prednisone or prednisolone 5 mg twice daily. Control patients received placebo and low dose prednisone or prednisolone 5 mg twice daily.

Because changes in PSA serum concentration do not always predict clinical benefit in all studies patients were maintained on abiraterone acetate until discontinuation criteria were met as specified below for each study.

Study 3011 (patients with newly diagnosed high-risk metastatic hormone sensitive prostate cancer (mHSPC))

In Study 3011, (n=1199) the median age of enrolled patients was 67 years. The ECOG performance status was 0 or 1 for 97% of patients. Patients with uncontrolled hypertension, significant heart disease, or NYHA Class II or worse heart failure were excluded. Co-primary efficacy endpoints were overall survival (OS) and radiographic progression-free survival (rPFS). The median baseline pain score, as measured by the Brief Pain Inventory Short Form (BPI-SF) was 2.0 in both the treatment and placebo groups. In addition to the co-primary endpoint measures, benefit was also assessed using time to skeletal-related event (SRE), time to subsequent therapy for prostate cancer, time to initiation of chemotherapy, time to pain progression and time to PSA progression.

In the 3011 study, treatment continued until disease progression, withdrawal of consent, the occurrence of unacceptable toxicity, or death.

Radiographic progression-free survival was defined as the time from randomization to the occurrence of radiographic progression or death from any cause. Radiographic progression included progression by bone scan (according to modified PCWG2) or progression of soft tissue lesions by CT or MRI (according to RECIST 1.1).

At the planned rPFS analysis there were 593 events; 239 (40.0%) of patients treated with abiraterone acetate and 354 (58.8%) of patients treated with placebo had radiographic evidence of progression or had died. A significant difference in rPFS between treatment groups was observed (see Table 2 and Figure 1).

Table 2: Radiographic Progression-Free Survival - Stratified Analysis; Intent-to-treat Population (Study PCR3011)

| | AA-P | Placebo |
|--------------------------|----------------------|--------------------|
| Subjects randomized | 597 | 602 |
| Event | 239 (40.0%) | 354 (58.8%) |
| Censored | 358 (60.0%) | 248 (41.2%) |
| Time to Event (months) | | |
| 25th percentile (95% CI) | 14.59 (11.47, 15.61) | 7.43 (7.29, 10.58) |


| | | |
|------------------------------------|----------------------|----------------------|
| Median (95% CI) | 33.02 (29.57, NE) | 14.78 (14.69, 18.27) |
| 75th percentile (95% CI) | NE (NE, NE) | 30.36 (29.24, 39.95) |
| Range | (0.0+, 41.0+) | (0.0+, 40.6+) |
| 6-month event-free rate (95% CI) | 0.941 (0.918, 0.957) | 0.867 (0.836, 0.892) |
| 12-month event-free rate (95% CI) | 0.779 (0.742, 0.812) | 0.611 (0.567, 0.652) |
| 18-month event-free rate (95% CI) | 0.702 (0.661, 0.739) | 0.476 (0.431, 0.520) |
| 24-month event-free rate (95% CI) | 0.611 (0.568, 0.652) | 0.347 (0.303, 0.391) |
| 30-month event-free rate (95% CI) | 0.532 (0.483, 0.579) | 0.250 (0.206, 0.296) |
| 36-month event-free rate (95% CI) | 0.471 (0.414, 0.526) | 0.209 (0.162, 0.260) |
| p value ^a | < 0.0001 | |
| Hazard ratio (95% CI) ^b | 0.466 (0.394, 0.550) | |

Note: += censored observation, NE=not estimable. The radiographic progression and death are considered in defining the rPFS event. AA-P= subjects who received abiraterone acetate and prednisone.

^a p value is from a log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^b Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Figure 1: Kaplan-Meier Plot of Radiographic Progression-free Survival; Intent-to-treat Population (Study PCR3011)



This final analysis of OS was conducted based on a clinical cutoff (CCO) of 15 August 2018, after 618 deaths were reported (median follow up of 51.8 months). Significant improvement in OS was demonstrated in the AA-P group compared with the Placebos group, showing a consistent and robust treatment effect in favor of AA-P treatment. In addition, updated data for key secondary efficacy endpoints (time to initiation of chemotherapy, time to subsequent therapy for prostate cancer, time to pain progression, time to skeletal-related events, time to prostate specific antigen [PSA] progression) were consistent with earlier results and continue to show the favorable treatment effect of AA-P. Across these secondary endpoints, statistically significant and clinically meaningful benefits in favor of AA-P plus ADT were observed.

Primary Efficacy Endpoint Analysis: Overall Survival

The co-primary efficacy endpoint of OS is defined as the time interval from the date of randomization

to date of death from any cause. Results from the final OS analysis are presented in Table 3 and Figure 2. As of the CCO of 15 August 2018, 618 deaths were reported: 275 (46%) in the AA-P group and 343 (57%) in the Placebos group. The median follow-up time for all subjects was 51.8 months (PCR3011/Attachment TEFOS08). The hazard ratio (HR) for OS was 0.661 (95% confidence interval [CI]: 0.564, 0.775; $p < 0.0001$), representing a 34% reduction in the risk of death. The median survival was 53.3 months in the AA-P group and 36.5 months in the Placebos group, for a difference in median survival between groups of 16.8 months.

Table 3: Overall Survival, Stratified Analysis; Intent-to-treat Population (Study 212082PCR3011)


| | AA-P | Placebo |
|------------------------------------|----------------------|----------------------|
| Subjects randomized | 597 | 602 |
| Event | 275 (46.1%) | 343 (57.0%) |
| Censored | 322 (53.9%) | 259 (43.0%) |
| Overall Survival (months) | | |
| 25th percentile (95% CI) | 26.25 (22.74, 29.93) | 19.75 (17.91, 21.85) |
| Median (95% CI) | 53.32 (48.23, NE) | 36.53 (33.54, 39.95) |
| 75th percentile (95% CI) | NE (NE, NE) | NE (60.19, NE) |
| Range | (0.1, 65.2+) | (1.4+, 64.6+) |
| 12-month event-free rate (95% CI) | 0.931 (0.908, 0.949) | 0.892 (0.863, 0.914) |
| 24-month event-free rate (95% CI) | 0.771 (0.734, 0.803) | 0.687 (0.646, 0.723) |
| 36-month event-free rate (95% CI) | 0.650 (0.608, 0.688) | 0.503 (0.460, 0.544) |
| 48-month event-free rate (95% CI) | 0.544 (0.501, 0.585) | 0.399 (0.356, 0.441) |
| 60-month event-free rate (95% CI) | 0.458 (0.408, 0.506) | 0.315 (0.265, 0.367) |
| p value ^a | < 0.0001 | |
| Hazard ratio (95% CI) ^b | 0.661 (0.564, 0.775) | |

Note: += censored observation, NE = not estimable

^ap value is from log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^bHazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Figure 2: Kaplan-Meier Plot of Overall Survival; Intent-to-treat Population (Study 212082PCR3011)



Note: Abiraterone acetate indicates abiraterone acetate plus prednisone

Secondary Efficacy Endpoint Analysis: Time to Initiation of Chemotherapy

Time to initiation of chemotherapy was defined as the time interval from the date of randomization to the date of initiation of chemotherapy for prostate cancer. Updated information from the final analysis is presented in Table 4 and Figure 3. Initiation of chemotherapy was documented for 25% of subjects in the AA-P group and 36% of subjects in the Placebos group. There was a 49% reduction in risk of initiation of chemotherapy (HR=0.508; 95% CI: 0.412, 0.627, $p<0.0001$). The median time to initiation of chemotherapy was not reached in the AA-P group and was 57.6 months in the Placebos group, demonstrating that AA-P delayed the need for initiation of chemotherapy. The 60-month event-free rate (ie, percent of subjects for whom chemotherapy was not required at 5 years after initiation of study treatment) was 66% for AA-P versus 48% for Placebos.

Table 4: Time to Initiation of Chemotherapy - Stratified Analysis; Intent-to-treat Population (Study 212082PCR3011)


| | AA-P | Placebo |
|------------------------------------|----------------------|----------------------|
| Subjects randomized | 597 | 602 |
| Event | 150 (25.1%) | 218 (36.2%) |
| Censored | 447 (74.9%) | 384 (63.8%) |
| Time to Event (months) | | |
| 25th percentile (95% CI) | 36.63 (30.16, 44.45) | 18.63 (16.07, 21.22) |
| Median (95% CI) | NE (62.62, NE) | 57.59 (38.18, NE) |
| 75th percentile (95% CI) | NE (NE, NE) | NE (NE, NE) |
| Range | (0.1+, 64.2+) | (1.4+, 64.6+) |
| 6-month event-free rate (95% CI) | 0.988 (0.975, 0.994) | 0.945 (0.923, 0.961) |
| 12-month event-free rate (95% CI) | 0.951 (0.930, 0.966) | 0.857 (0.825, 0.884) |
| 18-month event-free rate (95% CI) | 0.899 (0.871, 0.922) | 0.766 (0.727, 0.800) |
| 24-month event-free rate (95% CI) | 0.847 (0.812, 0.875) | 0.675 (0.631, 0.715) |
| 30-month event-free rate (95% CI) | 0.791 (0.752, 0.824) | 0.605 (0.558, 0.649) |
| 36-month event-free rate (95% CI) | 0.755 (0.714, 0.791) | 0.563 (0.514, 0.609) |
| 42-month event-free rate (95% CI) | 0.721 (0.678, 0.760) | 0.526 (0.475, 0.574) |
| 48-month event-free rate (95% CI) | 0.693 (0.647, 0.733) | 0.515 (0.464, 0.563) |
| 54-month event-free rate (95% CI) | 0.667 (0.618, 0.711) | 0.501 (0.447, 0.552) |
| 60-month event-free rate (95% CI) | 0.660 (0.610, 0.705) | 0.481 (0.417, 0.542) |
| p value ^a | < 0.0001 | |
| Hazard ratio (95% CI) ^b | 0.508 (0.412, 0.627) | |

Note: += censored observation, NE=not estimable.

^a p value is from a log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^b Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Figure 3: Kaplan-Meier Plot of Time to Initiation of Chemotherapy; Intent-to-treat Population (Study 212082PCR3011)



Note: Abiraterone acetate indicates abiraterone acetate plus prednisone

Time to Life-Extending Subsequent Therapy for Prostate Cancer

Time to life-extending subsequent therapy for prostate cancer (ie, docetaxel, cabazitaxel, abiraterone acetate, enzalutamide, sipuleucel-T, and radium-223) is summarized for the final analysis in Table 5 and Figure 4. There were 29% of subjects in the AA-P group and 45% of subjects in the Placebos group who received life-extending subsequent therapy. The median time to life-extending subsequent therapy was not reached in the AA-P group and was 29.6 months in the Placebos group, demonstrating that AA-P delayed the need for initiation of life-extending subsequent therapy (HR=0.431; 95% CI: 0.356, 0.522; $p<0.0001$). The 60-month event-free rate was 60% for AA-P and 37% for Placebos.

Table 5: Time to life-extending Subsequent Therapy for Prostate Cancer- Stratified Analysis; Intent-to-treat Population (Study 212082PCR3011)

| | AA-P | Placebo |
|-----------------------------------|----------------------|----------------------|
| Subjects randomized | 597 | 602 |
| Event | 176 (29.5%) | 273 (45.3%) |
| Censored | 421 (70.5%) | 329 (54.7%) |
| Time to Event (months) | | |
| 25th percentile (95% CI) | 30.62 (26.61, 34.99) | 14.85 (13.21, 16.66) |
| Median (95% CI) | NE (60.78, NE) | 29.63 (25.86, 34.99) |
| 75th percentile (95% CI) | NE (NE, NE) | NE (NE, NE) |
| Range | (0.1+, 64.2+) | (1.4+, 63.8+) |
| 6-month event-free rate (95% CI) | 0.988 (0.975, 0.994) | 0.938 (0.915, 0.955) |
| 12-month event-free rate (95% CI) | 0.946 (0.924, 0.962) | 0.822 (0.787, 0.852) |
| 18-month event-free rate (95% CI) | 0.888 (0.858, 0.912) | 0.685 (0.643, 0.724) |
| 24-month event-free rate (95% CI) | 0.819 (0.782, 0.849) | 0.577 (0.531, 0.620) |
| 30-month event-free rate (95% CI) | 0.754 (0.713, 0.790) | 0.494 (0.446, 0.540) |
| 36-month event-free rate (95% CI) | 0.709 (0.665, 0.747) | 0.439 (0.390, 0.486) |
| 42-month event-free rate (95% CI) | 0.671 (0.625, 0.712) | 0.408 (0.360, 0.457) |
| 48-month event-free rate (95% CI) | 0.651 (0.604, 0.693) | 0.388 (0.339, 0.437) |
| 54-month event-free rate (95% CI) | 0.621 (0.572, 0.666) | 0.388 (0.339, 0.437) |
| 60-month event-free rate (95% CI) | 0.599 (0.546, 0.648) | 0.367 (0.305, 0.429) |

p value^a < 0.0001
Hazard ratio (95% CI)^b 0.431 (0.356, 0.522)


Note: += censored observation, NE=not estimable.

^a p value is from a log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^b Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Life-extending therapies include: Docetaxel, Cabazitaxel, Abiraterone acetate, Enzalutamide, Sipuleucel-T, and Ra-223.

Figure 4: Kaplan-Meier Plot of Time to Life-extending Subsequent Therapy for Prostate Cancer; Intent-to-treat Population (Study 212082PCR3011)



Life-extending therapies include: Docetaxel, Cabazitaxel, Abiraterone acetate plus prednisone, Enzalutamide, Sipuleucel-T, and Radium-223.

Note: Abiraterone acetate indicates abiraterone acetate plus prednisone

Time to Pain Progression

Time to pain progression was defined as the time interval from randomization to the first date a subject experienced a $\geq 30\%$ increase from baseline in the BPI-SF worst pain intensity (Item 3) observed at 2 consecutive evaluations ≥ 4 weeks apart. Updated data for time to pain progression from the final analysis is presented in Table 6 and Figure 5. Pain progression was documented for 41% of subjects in the AA-P group and 49% of subjects in the Placebos group. There was a 28% reduction in risk of pain progression (HR=0.721; 95% CI: 0.607, 0.857; p=0.0002). The median time to pain progression was 47.4 months in the AA-P group and 16.6 months in the Placebos group. The 48-month event-free rate was 50% for AA-P and 39% for the Placebos group.

Table 6: Time to Pain Progression (BPI3) - Stratified Analysis; Intent-to-treat Population (Study 212082PCR3011)

| | AA-P | Placebo |
|--------------------------|-------------------|----------------------|
| Subjects randomized | 597 | 602 |
| Event | 245 (41.0%) | 292 (48.5%) |
| Censored | 352 (59.0%) | 310 (51.5%) |
| Time to Event (months) | | |
| 25th percentile (95% CI) | 4.67 (2.89, 6.47) | 2.79 (2.73, 3.71) |
| Median (95% CI) | 47.41 (33.15, NE) | 16.62 (11.07, 23.95) |


| 75th percentile (95% CI) Range | NE (NE, NE) (0.0+, 60.7+) | NE (NE, NE) (0.0+, 55.8+) |
|------------------------------------|------------------------------|------------------------------|
| 6-month event-free rate (95% CI) | 0.723 (0.685, 0.758) | 0.635 (0.594, 0.673) |
| 12-month event-free rate (95% CI) | 0.630 (0.588, 0.669) | 0.534 (0.490, 0.575) |
| 18-month event-free rate (95% CI) | 0.603 (0.560, 0.643) | 0.496 (0.451, 0.540) |
| 24-month event-free rate (95% CI) | 0.577 (0.533, 0.618) | 0.448 (0.400, 0.495) |
| 30-month event-free rate (95% CI) | 0.556 (0.510, 0.598) | 0.417 (0.367, 0.467) |
| 36-month event-free rate (95% CI) | 0.541 (0.495, 0.585) | 0.410 (0.358, 0.461) |
| 42-month event-free rate (95% CI) | 0.526 (0.478, 0.572) | 0.410 (0.358, 0.461) |
| 48-month event-free rate (95% CI) | 0.499 (0.439, 0.557) | 0.388 (0.324, 0.451) |
| 54-month event-free rate (95% CI) | 0.499 (0.439, 0.557) | 0.388 (0.324, 0.451) |
| 60-month event-free rate (95% CI) | 0.499 (0.439, 0.557) | NE (NE, NE) |
| p value ^a | 0.0002 | |
| Hazard ratio (95% CI) ^b | 0.721 (0.607, 0.857) | |

Note: += censored observation, NE=not estimable.

^a p value is from a log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^b Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Figure 5: Kaplan-Meier Plot of Time to Pain Progression (BPI3); Intent-to-treat Population (Study 212082PCR3011)



Note: Abiraterone acetate indicates abiraterone acetate plus prednisone

Time to Skeletal-related Event

Time to skeletal-related event was defined as the earliest of the following: clinical or pathological fracture, spinal cord compression, palliative radiation to bone, or surgery to bone. Updated information from the final analysis is presented for time to skeletal-related event in Table 7 and Figure 6. Skeletal-related events were reported for 22% of subjects in the AA-P group and 25% of subjects in the Placebos group. There was a 24% reduction in the risk of skeletal-related event (HR=0.759; 95% CI: 0.601, 0.960; p=0.0208). The 25th percentile time to skeletal-related event was 43.0 months for the AA-P group and 31.3 months for Placebos. The 60-month event-free rate was 71% for AA-P and 64% for Placebos.

Table 7: Time to Skeletal-Related Event - Stratified Analysis; Intent-to-treat Population (Study 212082PCR3011)


| | AA-P | Placebo |
|------------------------------------|----------------------|----------------------|
| Subjects randomized | 597 | 602 |
| Event | 133 (22.3%) | 150 (24.9%) |
| Censored | 464 (77.7%) | 452 (75.1%) |
| Time to Event (months) | | |
| 25th percentile (95% CI) | 43.04 (34.56, 53.52) | 31.34 (23.29, 41.92) |
| Median (95% CI) | NE (NE, NE) | NE (NE, NE) |
| 75th percentile (95% CI) | NE (NE, NE) | NE (NE, NE) |
| Range | (0.1, 64.2+) | (0.8, 64.6+) |
| 6-month event-free rate (95% CI) | 0.965 (0.947, 0.977) | 0.945 (0.924, 0.961) |
| 12-month event-free rate (95% CI) | 0.935 (0.911, 0.952) | 0.877 (0.847, 0.902) |
| 18-month event-free rate (95% CI) | 0.892 (0.863, 0.915) | 0.827 (0.792, 0.856) |
| 24-month event-free rate (95% CI) | 0.841 (0.807, 0.870) | 0.786 (0.748, 0.819) |
| 30-month event-free rate (95% CI) | 0.807 (0.770, 0.839) | 0.753 (0.712, 0.789) |
| 36-month event-free rate (95% CI) | 0.783 (0.743, 0.817) | 0.728 (0.685, 0.767) |
| 42-month event-free rate (95% CI) | 0.753 (0.711, 0.790) | 0.707 (0.662, 0.748) |
| 48-month event-free rate (95% CI) | 0.736 (0.692, 0.774) | 0.674 (0.625, 0.718) |
| 54-month event-free rate (95% CI) | 0.706 (0.658, 0.748) | 0.667 (0.616, 0.713) |
| 60-month event-free rate (95% CI) | 0.706 (0.658, 0.748) | 0.643 (0.584, 0.697) |
| p value ^a | 0.0208 | |
| Hazard ratio (95% CI) ^b | 0.759 (0.601, 0.960) | |

Note: += censored observation, NE=not estimable.

^a p value is from a log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^b Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Figure 6: Kaplan-Meier Plot of Time to Skeletal-related event; Intent-to-treat Population (Study 212082PCR3011)



Note: Abiraterone acetate indicates abiraterone acetate plus prednisone

Time to PSA Progression

Time to PSA progression was defined as the time interval from the date of randomization to the date of PSA progression, according to Prostate Cancer Working Group 2 (PCWG2) criteria. At the time of this final analysis, PSA progression was documented for 46% of subjects in the AA-P group and 74% of subjects in the Placebos group (Table 8 and Figure 7). Treatment with AA-P statistically significantly decreased the risk of PSA progression by 69% compared with Placebos (HR=0.310; 95% CI: 0.266, 0.363; $p<0.0001$). The median time to PSA progression was 33.3 months in the AA-P group and 7.4 months in the Placebos group, a delay in PSA progression by >25 months in the AA-P group compared with the Placebos group. As shown in Figure 7, this separation continued over the course of time measured. The 60-month event-free rate was 42% for AA-P and 11% for Placebos.

Table 8: Time to PSA Progression - Stratified Analysis; Intent-to-treat Population (Study 212082PCR3011)


| | AA-P | Placebo |
|------------------------------------|----------------------|----------------------|
| Subjects randomized | 597 | 602 |
| Event | 273 (45.7%) | 448 (74.4%) |
| Censored | 324 (54.3%) | 154 (25.6%) |
| Time to Event (months) | | |
| 25th percentile (95% CI) | 12.71 (10.15, 14.75) | 3.75 (3.71, 4.63) |
| Median (95% CI) | 33.31 (29.44, 46.09) | 7.43 (7.20, 9.20) |
| 75th percentile (95% CI) | NE (NE, NE) | 18.37 (14.82, 20.30) |
| Range | (0.0+, 64.2+) | (0.0+, 62.1+) |
| 6-month event-free rate (95% CI) | 0.885 (0.856, 0.909) | 0.616 (0.574, 0.656) |
| 12-month event-free rate (95% CI) | 0.751 (0.712, 0.785) | 0.356 (0.315, 0.397) |
| 18-month event-free rate (95% CI) | 0.665 (0.623, 0.704) | 0.254 (0.217, 0.293) |
| 24-month event-free rate (95% CI) | 0.577 (0.532, 0.618) | 0.184 (0.151, 0.220) |
| 30-month event-free rate (95% CI) | 0.527 (0.481, 0.570) | 0.140 (0.110, 0.174) |
| 36-month event-free rate (95% CI) | 0.489 (0.443, 0.534) | 0.121 (0.092, 0.153) |
| 42-month event-free rate (95% CI) | 0.468 (0.422, 0.513) | 0.112 (0.084, 0.144) |
| 48-month event-free rate (95% CI) | 0.453 (0.406, 0.498) | 0.109 (0.081, 0.141) |
| 54-month event-free rate (95% CI) | 0.430 (0.382, 0.478) | 0.109 (0.081, 0.141) |
| 60-month event-free rate (95% CI) | 0.423 (0.373, 0.472) | 0.109 (0.081, 0.141) |
| p value ^a | < 0.0001 | |
| Hazard ratio (95% CI) ^b | 0.310 (0.266, 0.363) | |

Note: += censored observation, NE=not estimable.

^a p value is from a log-rank test stratified by ECOG PS score(0/1 or 2) and visceral (absent or present).

^b Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors AA-P.

Figure 7: Kaplan-Meier Plot of PSA Progression; Intent-to-treat Population (Study 212082PCR3011)



Note: Abiraterone acetate indicates abiraterone acetate plus prednisone

Study 302 (asymptomatic or mildly symptomatic patients who did not receive prior chemotherapy)

This study enrolled chemotherapy naïve patients who were asymptomatic or mildly symptomatic and for whom chemotherapy was not yet clinically indicated. A score of 0-1 on Brief Pain Inventory-Short Form (BPI-SF) worst pain in last 24 hours was considered asymptomatic, and a score of 2-3 was considered mildly symptomatic.

In Study 302, (n=1088) the median age of enrolled patients was 71 years for patients treated with abiraterone acetate plus prednisone or prednisolone and 70 years for patients treated with placebo plus prednisone or prednisolone. The number of patients treated with abiraterone acetate by racial group was Caucasian 520 (95.4%), Black 15 (2.8%), Asian 4 (0.7%) and other 6 (1.1%). The ECOG performance status was 0 for 76% of patients, and 1 for 24% of patients in both arms. Patients with visceral metastases were excluded. Fifty percent of patients had only bone metastases, an additional 31% of patients had bone and soft tissue or lymph node metastases and 19% of patients had only soft tissue or lymph node metastases. Patients with visceral metastases were excluded. Co-primary efficacy endpoints were overall survival and radiographic progression-free survival (rPFS). Baseline pain assessment was 0-1 (asymptomatic) in 66% of patients and 2-3 (mildly symptomatic) in 26% of patients as defined by the Brief Pain Inventory-Short Form (worst pain over the last 24 hours). In addition to the co-primary endpoint measures, benefit was also assessed using time to opiate use for cancer pain, time to initiation of cytotoxic chemotherapy, time to deterioration in ECOG performance score by ≥ 1 point and time to PSA progression based on Prostate Cancer Working Group-2 (PCWG2) criteria. Study treatments were discontinued at the time of unequivocal clinical progression. Treatments could also be discontinued at the time of confirmed radiographic progression at the discretion of the investigator.

Radiographic progression-free survival was assessed with the use of sequential imaging studies as defined by PCWG2 criteria (for bone lesions) and modified Response Evaluation Criteria In Solid Tumors (RECIST) criteria (for soft tissue lesions). Analysis of rPFS utilized centrally-reviewed radiographic assessment of progression.

At the planned rPFS analysis there were 401 events; 150 (28%) of patients treated with abiraterone acetate and 251 (46%) of patients treated with placebo had radiographic evidence of progression or had died. A significant difference in rPFS between treatment groups was observed (see Table 9 and Figure 8).

Table 9: Study 302: Radiographic Progression-free Survival of Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone or Prednisolone Plus LHRH Agonists or Prior Orchiectomy


| | Abiraterone acetate (N=546) | PLACEBO (N=542) |
|--|--|----------------------------|
| Radiographic Progression-free Survival (rPFS) | | |
| Progression or death | 150 (28%) | 251 (46%) |
| Median rPFS in months (95% CI) | Not reached (11.66, NE) | 8.3 (8.12, 8.54) |
| p value* | <0.0001 | |
| Hazard ratio** (95% CI) | 0.425 (0.347, 0.522) | |

NE=Not Estimated.

* p-value is derived from a log-rank test stratified by baseline ECOG score (0 or 1).

** Hazard ratio (HR) < 1 favours abiraterone acetate.

Figure 8: Kaplan Meier Curves of Radiographic Progression-free Survival in Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone or Prednisolone plus LHRH Agonists or Prior Orchiectomy



AA= abiraterone acetate

However, subject data continued to be collected through the date of the second interim analysis of Overall survival (OS). The investigator radiographic review of rPFS performed as a follow up sensitivity analysis is presented in Table 10 and Figure 9.

Six hundred and seven (607) subjects had radiographic progression or died: 271 (50%) in the abiraterone acetate group and 336 (62%) in the placebo group. Treatment with abiraterone acetate decreased the risk of radiographic progression or death by 47% compared with placebo (HR=0.530; 95% CI: [0.451, 0.623], $p < 0.0001$). The median rPFS was 16.5 months in the abiraterone acetate group and 8.3 months in the placebo group.


Table 10: Study 302: Radiographic progression-free survival of patients treated with either abiraterone acetate or placebo in combination with prednisone or prednisolone plus LHRH analogues or prior orchiectomy (At second interim analysis of OS-Investigator Review)

| | Abiraterone acetate (N=546) | PLACEBO (N=542) |
|--|--|----------------------------|
| Radiographic Progression-free Survival (rPFS) | | |
| Progression or death | 271 (50%) | 336 (62%) |
| Median rPFS in months (95% CI) | 16.5 (13.80, 16.79) | 8.3 (8.05, 9.43) |
| p-value* | < 0.0001 | |
| Hazard ratio** (95% CI) | 0.530 (0.451, 0.623) | |

* p-value is derived from a log-rank test stratified by baseline ECOG score (0 or 1).

** Hazard ratio < 1 favours abiraterone acetate.

Figure 9: Kaplan Meier curves of radiographic progression-free survival in patients treated with either abiraterone acetate or placebo in combination with prednisone or prednisolone plus LHRH analogues or prior orchiectomy (At second interim analysis of OS-Investigator Review)



AA= abiraterone acetate

A planned interim analysis (IA) for overall survival was conducted after 333 deaths were observed. The study was unblinded based on the magnitude of clinical benefit observed and patients in the placebo group were offered treatment with abiraterone acetate. Overall survival was longer for abiraterone acetate than placebo with a 25% reduction in risk of death (HR = 0.752; 95% CI: [0.606, 0.934], p = 0.0097), but OS was not mature and interim results did not meet the pre-specified stopping boundary for statistical significance (see Table 11). Survival continued to be followed after this IA.

The planned final analysis for OS was conducted after 741 deaths were observed (median follow up of 49 months). Sixty-five percent (354 of 546) of patients treated with abiraterone acetate, compared with 71% (387 of 542) of patients treated with placebo, had died. A statistically significant OS benefit in

favor of the abiraterone acetate-treated group was demonstrated with a 19.4% reduction in risk of death (HR=0.806; 95% CI: [0.697, 0.931], p = 0.0033) and an improvement in median OS of 4.4 months (abiraterone acetate 34.7 months, placebo 30.3 months) (see Table 11 and Figure 10). This improvement was demonstrated despite subsequent therapy being common, irrespective of whether patients initially received abiraterone acetate or placebo. Subsequent therapies in the abiraterone acetate and placebo patient groups included abiraterone acetate, 69 (13%) and 238 (44%); docetaxel, 311 (57%) and 331 (61%); cabazitaxel, 100 (18%) and 105 (19%); and enzalutamide 87 (16%) and 54 (10%) patients respectively.

Table 11: Study 302: Overall Survival of Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone or Prednisolone Plus LHRH Agonists or Prior Orchiectomy


| | Abiraterone acetate (N=546) | PLACEBO (N=542) |
|---|--------------------------------|---------------------|
| Interim Survival Analysis | | |
| Deaths | 147 (27%) | 186 (34%) |
| Median overall survival in months (95% CI) | Not reached (NE, NE) | 27.2 (25.95, NE) |
| p value* | 0.0097 | |
| Hazard ratio** (95% CI) | 0.752 (0.606, 0.934) | |
| Final Survival Analysis | | |
| Deaths | 354 (65%) | 387 (71%) |
| Median overall survival in months (95% CI) | 34.7 (32.7, 36.8) | 30.3 (28.7, 33.3) |
| p value* | 0.0033 | |
| Hazard ratio** (95% CI) | 0.806 (0.697, 0.931) | |

NE=Not Estimated.

* p-value is derived from a log-rank test stratified by baseline ECOG score (0 or 1).

** Hazard ratio < 1 favours abiraterone acetate.

Figure 10: Kaplan Meier Survival Curves of Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone or Prednisolone plus LHRH Agonists or Prior Orchiectomy, Final analysis



AA= abiraterone acetate

Subgroup analyses consistently favor treatment with abiraterone acetate (see Figure 11).

Figure 11: Overall Survival by Subgroup: Hazard Ratio and 95% Confidence Interval

| Variable | Subgroup | Median (months) | | HR | 95% C.I. | Events/N | |
|-------------------------------|----------|-----------------|---------|------|--------------|----------|---------|
| | | AA | Placebo | | | AA | Placebo |
| All subjects | ALL | 34.7 | 30.3 | 0.81 | (0.70, 0.93) | 354/546 | 387/542 |
| Baseline ECOG | 0 | 35.4 | 32.0 | 0.79 | (0.66, 0.93) | 261/416 | 292/414 |
| | 1 | 27.9 | 26.4 | 0.87 | (0.65, 1.16) | 93/130 | 95/128 |
| Baseline BPI | 0-1 | 38.1 | 33.4 | 0.77 | (0.64, 0.93) | 223/370 | 233/346 |
| | 2-3 | 26.4 | 27.4 | 0.97 | (0.75, 1.27) | 100/129 | 120/147 |
| Bone Metastasis Only At Entry | YES | 38.9 | 34.1 | 0.78 | (0.62, 0.97) | 147/238 | 162/241 |
| | NO | 31.6 | 29.0 | 0.83 | (0.69, 1.00) | 207/308 | 225/301 |
| Age | <65 | 34.5 | 30.2 | 0.78 | (0.59, 1.03) | 89/135 | 111/155 |
| | >=65 | 34.7 | 30.8 | 0.81 | (0.69, 0.96) | 265/411 | 276/387 |
| | >=75 | 29.3 | 25.9 | 0.79 | (0.61, 1.01) | 125/185 | 125/165 |
| Baseline PSA above median | YES | 28.5 | 25.8 | 0.86 | (0.71, 1.04) | 208/282 | 206/260 |
| | NO | 43.1 | 34.4 | 0.72 | (0.58, 0.90) | 146/264 | 181/282 |
| Baseline LDH above median | YES | 31.2 | 24.8 | 0.74 | (0.61, 0.90) | 192/278 | 203/259 |
| | NO | 38.3 | 35.8 | 0.85 | (0.69, 1.05) | 162/268 | 184/283 |
| Baseline ALK-P above median | YES | 28.6 | 26.8 | 0.92 | (0.76, 1.11) | 211/279 | 201/256 |
| | NO | 44.5 | 33.2 | 0.68 | (0.55, 0.85) | 143/267 | 186/286 |
| Region | N.A. | 37.0 | 31.2 | 0.74 | (0.61, 0.91) | 184/297 | 198/275 |
| | Other | 33.2 | 30.1 | 0.90 | (0.73, 1.11) | 170/249 | 189/267 |

0.2 0.75 1.5
 Favors AA ← → Favors Placebo

AA= abiraterone acetate; ALK-P=alkaline phosphatase; BPI=Brief Pain Inventory; C.I.=confidence interval; ECOG=Eastern Cooperative Oncology Group performance score; HR=hazard ratio; LDH=lactic dehydrogenase; N.A.=North America; NE=not evaluable

In addition to the observed improvements in overall survival and rPFS, benefit was demonstrated for

abiraterone acetate vs. placebo treatment in all prospectively-defined secondary endpoint measures as follows:

Time to PSA progression based on PCWG2 criteria

The median time to PSA progression was 11.1 months for patients receiving abiraterone acetate and 5.6 months for patients receiving placebo (HR=0.488; 95% CI: [0.420, 0.568], $p<0.0001$). The time to PSA progression was approximately doubled with abiraterone acetate treatment (HR=0.488). The proportion of subjects with a confirmed PSA response was greater in the abiraterone acetate group than in the placebo group (62% versus 24%; $p<0.0001$). In subjects with measurable soft tissue disease, significantly increased numbers of complete and partial tumor responses were seen with abiraterone acetate treatment.

Time to opiate use for cancer pain

The median time to opiate use for prostate cancer pain at the time of final analysis was 33.4 months for patients receiving abiraterone acetate and was 23.4 months for patients receiving placebo (HR=0.721; 95% CI: [0.614, 0.846], $p<0.0001$).

Time to initiation of cytotoxic chemotherapy

The median time to initiation of cytotoxic chemotherapy was 25.2 months for patients receiving abiraterone acetate and 16.8 months for patients receiving placebo (HR=0.580; 95% CI: [0.487, 0.691], $p<0.0001$).

Time to deterioration in ECOG performance score by ≥ 1 point

The median time to deterioration in ECOG performance score by ≥ 1 point was 12.3 months for patients receiving abiraterone acetate and 10.9 months for patients receiving placebo (HR=0.821; 95% CI: [0.714, 0.943], $p=0.0053$).

The following study endpoints demonstrated a statistically significant advantage in favor of abiraterone acetate treatment:

Objective response

Objective response was defined as the proportion of subjects with measurable disease achieving a complete or partial response according to RECIST criteria (baseline lymph node size was required to be ≥ 2 cm to be considered a target lesion). The proportion of subjects with measurable disease at baseline who had an objective response was 36% in the abiraterone acetate group and 16% in the placebo group ($p<0.0001$).

Pain

Treatment with abiraterone acetate significantly reduced the risk of average pain intensity progression by 18% compared with placebo ($p=0.0490$). The median time to progression was 26.7 months in the abiraterone acetate group and 18.4 months in the placebo group.

Time to degradation in the FACT-P (Total Score)

Treatment with abiraterone acetate decreased the risk of FACT-P (Total Score) degradation by 22% compared with placebo ($p=0.0028$). The median time to degradation in FACT-P (Total Score) was 12.7 months in the abiraterone acetate group and 8.3 months in the placebo group.

Study 301 (patients who had received prior chemotherapy)

Study 301 enrolled patients who had received prior docetaxel. Patients were not required to show disease progression on docetaxel, as toxicity from this chemotherapy may have led to discontinuation.

Patients were maintained on study treatments until there was PSA progression (confirmed 25% increase over the patient's baseline/nadir) together with protocol-defined radiographic progression and symptomatic or clinical progression. Patients with prior ketoconazole treatment for prostate cancer were excluded from this study. The primary efficacy endpoint was overall survival.

The median age of enrolled patients was 69 years (range 39-95). The number of patients treated with abiraterone acetate by racial group was Caucasian 737 (93.2%), Black 28 (3.5%), Asian 11 (1.4%) and

other 14 (1.8%). Eleven percent of patients enrolled had an ECOG performance score of 2; 70% had radiographic evidence of disease progression with or without PSA progression; 70% had received one prior cytotoxic chemotherapy and 30% received two. Liver metastasis was present in 11% of patients treated with abiraterone acetate.

In a planned analysis conducted after 552 deaths were observed, 42% (333 of 797) of patients treated with abiraterone acetate compared with 55% (219 of 398) of patients treated with placebo, had died. A statistically significant improvement in median overall survival was seen in patients treated with abiraterone acetate (see Table 12 and Figure 12). An updated survival analysis was conducted when 775 deaths (97% of the planned number of deaths for the final analysis) were observed. Results from this updated survival analysis were consistent with those in the primary survival analysis (see Table 3).

Table 12: Study 301: Overall survival of patients treated with either abiraterone acetate or placebo in combination with prednisone or prednisolone plus LHRH analogues or prior orchiectomy


| | Abiraterone acetate (N=797) | PLACEBO (N=398) |
|--------------------------------------|--|----------------------------|
| Primary Survival Analysis | | |
| Deaths (%) | 333 (42%) | 219 (55%) |
| Median survival (months) (95% CI) | 14.8 (14.1, 15.4) | 10.9 (10.2, 12.0) |
| p value ^a | <0.0001 | |
| Hazard ratio (95% CI) ^b | 0.646 (0.543, 0.768) | |
| Updated Survival Analysis | | |
| Deaths (%) | 501 (63%) | 274 (69%) |
| Median survival (months) (95% CI) | 15.8 (14.8, 17.0) | 11.2 (10.4, 13.1) |
| Hazard ratio (95% CI) ^b | 0.740 (0.638, 0.859) | |

^a p-value is derived from a log-rank test stratified by ECOG performance status score (0-1 vs. 2), pain score (absent vs. present), number of prior chemotherapy regimens (1 vs. 2), and type of disease progression (PSA only vs. radiographic).

^b Hazard ratio is derived from a stratified proportional hazards model. Hazard ratio < 1 favours abiraterone acetate

At all evaluation time points after the initial few months of treatment, a higher proportion of patients treated with abiraterone acetate remained alive, compared with the proportion of patients treated with placebo (see Figure 12).

Figure 12: Kaplan Meier survival curves of patients treated with either abiraterone acetate or placebo in combination with prednisone or prednisolone plus LHRH analogues or prior orchiectomy



AA= abiraterone acetate

Subgroup survival analyses showed a consistent survival benefit for treatment with abiraterone acetate (see Figure 13).

Figure 13: Overall survival by subgroup: hazard ratio and 95% confidence interval

| Variable | Subgroup | Median (months) | | | HR | 95% C.I. |
|-----------------------------|--------------|-----------------|---------|--|------|--------------|
| | | AA | Placebo | | | |
| All subjects | ALL | 14.8 | 10.9 | | 0.66 | (0.56, 0.79) |
| Baseline ECOG | 0-1 | 15.3 | 11.7 | | 0.64 | (0.53, 0.78) |
| | 2 | 7.3 | 7 | | 0.81 | (0.53, 1.24) |
| Baseline BPI | <4 | 16.2 | 13 | | 0.64 | (0.50, 0.82) |
| | >=4 | 12.6 | 8.9 | | 0.68 | (0.53, 0.85) |
| No. prior chemo regimens | 1 | 15.4 | 11.5 | | 0.63 | (0.51, 0.78) |
| | 2 | 14 | 10.3 | | 0.74 | (0.55, 0.99) |
| Type of progression | PSA only | NE | 12.3 | | 0.59 | (0.42, 0.82) |
| | Radiographic | 14.2 | 10.4 | | 0.69 | (0.56, 0.84) |
| Age | <65 | 14.4 | 11.2 | | 0.66 | (0.48, 0.91) |
| | >=65 | 14.8 | 10.7 | | 0.67 | (0.55, 0.82) |
| | >=75 | 14.9 | 9.3 | | 0.52 | (0.38, 0.71) |
| Visceral disease at entry | YES | 12.6 | 8.4 | | 0.70 | (0.52, 0.94) |
| | NO | 15.4 | 11.2 | | 0.62 | (0.50, 0.76) |
| Baseline PSA above median | YES | 12.8 | 8.8 | | 0.65 | (0.52, 0.81) |
| | NO | 16.2 | 13.2 | | 0.69 | (0.53, 0.90) |
| Baseline LDH above median | YES | 10.4 | 8 | | 0.71 | (0.58, 0.88) |
| | NO | NE | 16.4 | | 0.64 | (0.47, 0.87) |
| Baseline ALK-P above median | YES | 11.6 | 8.1 | | 0.60 | (0.48, 0.74) |
| | NO | NE | 16.4 | | 0.73 | (0.54, 0.97) |
| Region | N.A. | 15.1 | 10.7 | | 0.64 | (0.51, 0.80) |
| | Other | 14.8 | 11.5 | | 0.69 | (0.54, 0.90) |

0.5 0.75 1 1.5
 Favors AA ← ————— → Favors Placebo

AA= abiraterone acetate; ALK-P=alkaline phosphatase; BPI=Brief Pain Inventory; C.I.=confidence interval; ECOG=Eastern Cooperative Oncology Group performance score; HR=hazard ratio;

LDH=lactic dehydrogenase; N.A.=North America, NE=not evaluable

In addition to the observed improvement in overall survival, all secondary study endpoints favoured abiraterone acetate and were statistically significant after adjusting for multiple testing as follows:

Patients receiving abiraterone acetate demonstrated a significantly higher total PSA response rate (defined as a $\geq 50\%$ reduction from baseline), compared with patients receiving placebo, 38% versus 10%, $p < 0.0001$.

The median time to PSA progression was 10.2 months for patients treated with abiraterone acetate and 6.6 months for patients treated with placebo (HR=0.580; 95% CI: [0.462; 0.728], $p < 0.0001$).

The median radiographic progression-free survival was 5.6 months for patients treated with abiraterone acetate and 3.6 months for patients who received placebo (HR=0.673; 95% CI: [0.585; 0.776], $p < 0.0001$).

Pain

The proportion of patients with pain palliation was statistically significantly higher in the abiraterone acetate group than in the placebo group (44% versus 27%, $p=0.0002$). A responder for pain palliation was defined as a patient who experienced at least a 30% reduction from baseline in the BPI-SF worst pain intensity score over the last 24 hours without any increase in analgesic usage score observed at two consecutive evaluations four weeks apart. Only patients with a baseline pain score of ≥ 4 and at least one post-baseline pain score were analysed (N=512) for pain palliation.

A lower proportion of patients treated with abiraterone acetate had pain progression compared to patients taking placebo at 6 (22% versus 28%), 12 (30% versus 38%) and 18 months (35% versus 46%). Pain progression was defined as an increase from baseline of $\geq 30\%$ in the BPI-SF worst pain intensity score over the previous 24 hours without a decrease in analgesic usage score observed at two consecutive visits, or an increase of $\geq 30\%$ in analgesic usage score observed at two consecutive visits. The time to pain progression at the 25th percentile was 7.4 months in the abiraterone acetate group, versus 4.7 months in the placebo group.

Skeletal-related events

A lower proportion of patients in the abiraterone acetate group had skeletal-related events compared with the placebo group at 6 months (18% versus 28%), 12 months (30% versus 40%), and 18 months (35% versus 40%). The time to first skeletal-related event at the 25th percentile in the abiraterone acetate group was twice that of the control group at 9.9 months versus 4.9 months. A skeletal-related event was defined as a pathological fracture, spinal cord compression, palliative radiation to bone, or surgery to bone.

5.2 Pharmacokinetic properties

General introduction

Following administration of abiraterone acetate, the pharmacokinetics of abiraterone and abiraterone acetate have been studied in healthy subjects, patients with metastatic advanced prostate cancer and subjects without cancer with hepatic or renal impairment. Abiraterone acetate is rapidly converted *in vivo* to abiraterone, an androgen biosynthesis inhibitor (see section 5 – Mechanism of action).

Absorption

Following oral administration of abiraterone acetate in the fasting state, the time to reach maximum plasma abiraterone concentration is approximately 2 hours.

Administration of abiraterone acetate with food, compared with administration in a fasted state, results in up to a 10-fold (AUC) and up to a 17-fold (C_{max}) increase in mean systemic exposure of abiraterone, depending on the fat content of the meal. Given the normal variation in the content and composition of meals, taking abiraterone acetate with meals has the potential to result in highly variable exposures. Therefore, abiraterone acetate must not be taken with food. Abiraterone acetate must be taken on an empty stomach, at least one hour before or at least two hours after a meal. The

tablets should be swallowed with water (see section 4.2).

Distribution and protein binding

The plasma protein binding of ^{14}C -abiraterone in human plasma is 99.8%. The apparent volume of distribution is approximately 5630 L, suggesting that abiraterone extensively distributes to peripheral tissues.

Metabolism#

Following oral administration of ^{14}C -abiraterone acetate as capsules, abiraterone acetate is hydrolysed to abiraterone, which then undergoes metabolism including sulphation, hydroxylation and oxidation primarily in the liver. The majority of circulating radioactivity (approximately 92%) is found in the form of metabolites of abiraterone. Of 15 detectable metabolites, 2 main metabolites, abiraterone sulphate and N-oxide abiraterone sulphate, each represents approximately 43% of total radioactivity.

Elimination

The mean half-life of abiraterone in plasma is approximately 15 hours based on data from healthy subjects. Following oral administration of ^{14}C -abiraterone acetate 1,000 mg, approximately 88% of the radioactive dose is recovered in faeces and approximately 5% in urine. The major compounds present in faeces are unchanged abiraterone acetate and abiraterone (approximately 55% and 22% of the administered dose, respectively).

Special Populations

Renal impairment

The pharmacokinetics of abiraterone acetate was compared in patients with end-stage renal disease on a stable haemodialysis schedule versus matched control subjects with normal renal function. Systemic exposure to abiraterone after a single oral 1,000 mg dose did not increase in subjects with end-stage renal disease on dialysis.

Administration in patients with renal impairment, including severe renal impairment, does not require dose reduction (see section 4.2 – Renal impairment). However, there is no clinical experience in patients with prostate cancer and severe renal impairment. Caution is advised in these patients.

Hepatic impairment

The pharmacokinetics of abiraterone acetate was examined in subjects with pre-existing mild or moderate hepatic impairment (Child-Pugh Class A and B, respectively) and in healthy control subjects. Systemic exposure to abiraterone after a single oral 1,000 mg dose increased by approximately 11% and 260% in subjects with mild and moderate pre-existing hepatic impairment, respectively. The mean half-life of abiraterone is prolonged to approximately 18 hours in subjects with mild hepatic impairment and to approximately 19 hours in subjects with moderate hepatic impairment. No dose adjustment is necessary for patients with pre-existing mild hepatic impairment. There are no data on the clinical safety and efficacy of multiple doses of abiraterone acetate when administered to patients with moderate or severe hepatic impairment (Child Pugh Class B or C). No dose adjustment can be predicted. Abiraterone acetate should be used with caution in patients with moderate hepatic impairment only if the benefit clearly outweighs the possible risk (see section 4.2 – Hepatic impairment and 4.4 – Hepatotoxicity and Hepatic impairment). abiraterone acetate should not be used in patients with severe hepatic impairment. For patients who develop hepatotoxicity during treatment, suspension of treatment and dose adjustment may be required (see section 4.2 – Hepatic impairment, 4.3 and 4.4 – Hepatotoxicity and Hepatic impairment).

5.3 Preclinical safety data

Carcinogenicity and Mutagenicity

Abiraterone acetate was not carcinogenic in a 6-month study in the transgenic (Tg.rasH2) mouse. In a 24-month carcinogenicity study in the rat, abiraterone acetate increased the incidence of interstitial cell neoplasms in the testes. This finding is considered related to the pharmacological action of abiraterone and rat specific. Abiraterone acetate was not carcinogenic in female rats.

Abiraterone acetate and abiraterone were devoid of genotoxic potential in the standard panel of

genotoxicity tests, including an *in vitro* bacterial reverse mutation assay (the Ames test), an *in vitro* mammalian chromosome aberration test (using human lymphocytes) and an *in vivo* rat micronucleus assay.

Reproductive Toxicology

In fertility studies in both male and female rats, abiraterone acetate reduced fertility, which was completely reversible in 4 to 16 weeks after abiraterone acetate was stopped.

In a developmental toxicity study in the rat, abiraterone acetate affected pregnancy including reduced fetal weight and survival. Effects on the external genitalia were observed though abiraterone acetate was not teratogenic.

In these fertility and developmental toxicity studies performed in the rat, all effects were related to the pharmacological activity of abiraterone.

Abiraterone acetate is contraindicated in pregnancy (see sections 4.3 and 4.6 – Pregnancy).

Animal Toxicology

In all animal toxicity studies, circulating testosterone levels were significantly reduced. As a result, reduction in organ weights and morphological and/or histopathological changes in the reproductive organs, and the adrenal, pituitary and mammary glands were observed. All changes showed complete or partial reversibility. The changes in the reproductive organs and androgen-sensitive organs are consistent with the pharmacology of abiraterone. All treatment-related hormonal changes reversed or were shown to be resolving after a 4-week recovery period.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Croscarmellose sodium
Sodium laurilsulfate
Povidone
Cellulose, microcrystalline
Lactose monohydrate
Silica, colloidal anhydrous
Magnesium stearate

Coating

Poly(vinyl alcohol)
Titanium dioxide
Macrogol 3350
Talc

For 500 mg only

Iron oxide red
Iron oxide black

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Please refer to outer carton.

6.4 Special precautions for storage

Store at or below 30°C.

Keep out of reach of children.

6.5 Nature and contents of container

TEROPROS FILM COATED TABLET 500MG, TEROPROS FILM COATED TABLET 250MG

Aluminium-OPA/Alu/PVC blisters or Aluminium-PVC/PE/PVDC blisters:

250 mg: pack sizes of 84, 84x1, 112, 112x1, 120 & 120x1 tablets

500mg: pack sizes of 14, 28, 56, 56x1, 60, 60x1, 84, 84x1, 112 & 112x1 tablets

High density polyethylene (HDPE) bottles closed with a polypropylene (PP) child resistant closure containing:

250 mg: 120 tabs

500mg: 60 tabs

TEROPROS FILM COATED TABLET 1000MG

- Aluminium-OPA/Alu/PVC blisters containing 28, 30 and 60 (as bundle pack: 2 packs of 30) tablets
- Aluminium-OPA/Alu/PVC perforated unit dose blisters containing 28x1, 30x1 and 60x1 (as bundle pack: 2 packs of 30x1) tablets
- Aluminium-OPA/Alu/PVC calendar blisters containing 28, 30 and 84 (as bundle pack: 3 packs of 28) tablets
- High density polyethylene (HDPE) bottles closed with a polypropylene (PP) child resistant closure.
- High density polyethylene (HDPE) bottles closed with a polypropylene (PP) child resistant closure, containing an oxygen absorbing canister.
- High density polyethylene (HDPE) bottles closed with a polypropylene (PP) child resistant closure, containing a moisture absorbing canister.

Bottles of 28 or 30 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Based on its mechanism of action, abiraterone acetate may harm a developing fetus; therefore, to avoid inadvertent exposure women who are pregnant or women whom may be pregnant should not handle abiraterone acetate without protection, e.g., gloves.

Any unused product or waste material should be disposed of in accordance with local requirements. This product may pose a risk to the aquatic environment.

7. PRODUCT REGISTRANT

Novartis (Singapore) Pte. Ltd.
20 Pasir Panjang Road,
#10-25/28, Mapletree Business City,
Singapore 117439

8. DATE OF REVISION

Feb 2022