

SIZE: 155 x 595 mm



1. NAME OF THE MEDICINAL PRODUCT
Siranalen 75 mg hard capsules
Siranalen 150 mg hard capsules
Siranalen 300 mg hard capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION
Each hard capsule contains 75 mg, 150 mg or 300 mg of pregabalin.
Excipient with known effect: lactose monohydrate. Siranalen 300mg also contains carmoisine (E122).
Each 75 mg hard capsule contains 8.25 mg lactose monohydrate.
Each 150 mg hard capsule contains 16.50 mg lactose monohydrate.
Each 300 mg hard capsule contains 33 mg lactose monohydrate and 0.0881mg carmoisine (E122).
For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM
Hard capsule.
Siranalen 75 mg hard capsules: Blue-white hard gelatin capsules size “4” (approximately 14.3mm in length) filled with off white powder.
Siranalen 150 mg hard capsules: White-white hard gelatin capsules size “2” (approximately 18.0 mm in length) filled with off white powder.
Siranalen 300 mg hard capsules: Maroon-white hard gelatin capsules size “0” (approximately 21.4 mm in length) filled with off white powder.

4.1 CLINICAL PARTICULARS
4.1 Therapeutic indications
Neuropathic pain
Siranalen is indicated for the treatment of neuropathic pain, which includes diabetic peripheral neuropathy and post-herpetic neuralgia in adults.
Epilepsy
Siranalen is indicated as adjunctive therapy of partial seizures, with or without secondary generalization, in adults.
Generalised Anxiety Disorder
Siranalen is indicated for the treatment of Generalised Anxiety Disorder (GAD) in adults.
Fibromyalgia
Pregabalin is indicated for the management of fibromyalgia.

4.2 Posology and method of administration
Posology
The dose range is 150 to 600 mg per day given in either two or three divided doses.
Neuropathic pain
Pregabalin treatment can be started at a dose of 150 mg per day. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after an interval of 3 to 7 days, and if needed, to a maximum dose of 600 mg per day after an additional 7-day interval.
Epilepsy
Pregabalin treatment can be started with a dose of 150 mg per day. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after 1 week. The maximum dose of 600 mg per day may be achieved after an additional week.
Generalised Anxiety Disorder
The dose range is 150 to 600 mg per day given as two or three divided doses. The need for treatment should be reassessed regularly.
Fibromyalgia
Pregabalin treatment can be started with a dose of 150 mg per day. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after 1 week. Following an additional week the dose may be increased to 450 mg per day. The maximum dosage of 600 mg per day may be achieved after an additional week.

Pharmacokinetics
The recommended dose of pregabalin is 300 to 450 mg per day. Dosing should begin at 75 mg two times a day (150 mg per day) and may be increased to 150 mg two times a day (300 mg per day) within 1 week based on efficacy and tolerability. Patients who do not experience sufficient benefit with 300 mg per day may be further increased to 225 mg two times a day (450 mg per day). Although pregabalin was also studied at 600 mg per day, there is no evidence that this dose confers additional benefit and that this dose was less tolerated.
In view of the dose-dependent adverse reactions, treatment with doses above 450 mg per day is not recommended.
Discontinuation of pregabalin
If pregabalin has to be discontinued it is recommended this should be done gradually over a minimum of 1 week independent of the indication.
Patients with renal impairment
Dose reduction in patients with compromised renal function must be individualised according to creatinine clearance (see section 5.2), as indicated in Table 1 determined using the following formula:
$$CL_{cr}(ml/min) = \frac{[140 - age (years) \times weight (kg)]}{72 \times serum creatinine (mg / dl)} \times (0.85 \text{ for female patients})$$

For patients receiving hemodialysis, the pregabalin daily dose should be adjusted according to renal function. In addition to the daily dose, a supplementary dose should be given immediately following every 4-hour hemodialysis treatment (see Table 1).
Table 1. Pregabalin dosage adjustment based on renal function

Creatinine clearance (<i>CL_{cr}</i>)(mL/min)	Total pregabalin daily dose*		Dose regimen
	Starting dose (mg/day)	Maximum dose(mg/day)	
≥ 60	150	600	BID or TID
≥30 < 60	75	300	BID or TID
≥15 < 30	25-50**	150	QD or BID
< 15	25**	75	QD
Supplementary dosage following haemodialysis (mg)			
	25**	100	Single dose+

TID = Three divided doses
BID = Two divided doses
QD = Single daily dose.
*Total daily dose (mg/day) should be divided as indicated by dose regimen to provide mg/dose
**These doses cannot be executed with the strengths available for Siranalen. Other products are available in strengths lower than 75 mg.
+Supplementary dose is a single additional dose
Patients with hepatic impairment
No dose adjustment is required for patients with hepatic impairment (see section 5.2).
Use in children and adolescents (12 to 17 years of age)
The safety and effectiveness of pregabalin in pediatric patients below the age of 12 years and adolescents have not been established. The use in children and adolescents is not recommended (see section 5.3).
Elderly (over 65 years of age)
Elderly patients may require a dose reduction of pregabalin due to a decreased renal function (see section 5.2).
Method of administration
Siranalen may be taken with or without food.
Siranalen is for oral use only.

4.3 Contraindications
Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use
Some diabetic patients who gain weight on pregabalin treatment may need to adjust hypoglycaemic medicinal products.
There have been reports in the post-marketing experience of hypersensitivity reactions, including cases of angioedema. Pregabalin should be discontinued immediately if symptoms of angioedema, such as facial, perioral, or upper airway swelling occur.
Pregabalin treatment has been associated with dizziness and somnolence, which could increase the occurrence of accidental injury (fall) in the elderly population. There have also been post-marketing reports of loss of consciousness, confusion and mental impairment. Therefore, patients should be advised to exercise caution until they are familiar with the potential effects of the medication.
In the post-marketing experience, transient visual blurring and other changes in visual acuity have been reported in patients treated with pregabalin. Discontinuation of pregabalin may result in resolution or improvement of these visual symptoms.
There are insufficient data for the withdrawal of concomitant antiepileptic medicinal products, once seizure control with pregabalin in the add-on situation has been reached, in order to reach monotherapy on pregabalin.
After discontinuation of short-term and long-term treatment with pregabalin withdrawal symptoms have been observed in some patients. The following events have been mentioned: insomnia, headache, nausea, anxiety, hyperhidrosis, diarrhoea, flu syndrome, nervousness, depression, pain, sweating and dizziness. The patient should be informed about this at the start of the treatment.
Pregabalin is not known to be active at receptor sites associated with drugs of abuse. Cases of misuse and abuse have been reported in the post-marketing database. As with any CNS active drug, carefully evaluate patients for history of drug abuse and observe them for signs of pregabalin misuse or abuse (e.g., development of tolerance, dose escalation, drug-seeking behavior).
Concerning discontinuation of long-term treatment of pregabalin there are no data of the incidence and severity of withdrawal symptoms in relation to duration of use and dosage of pregabalin. Although the effects of discontinuation on the reversibility of renal failure have not been systematically studied, improved renal function following discontinuation or dose reduction of pregabalin has been reported.
There have been post-marketing reports of congestive heart failure in some patients receiving pregabalin. In short-term trials of patients without clinically significant heart or peripheral vascular disease, there was no apparent association between peripheral edema and cardiovascular complications such as hypertension or congestive heart failure. Because there are limited data on severe congestive heart failure patients, pregabalin should be used with caution in these patients (see section 4.8).
Treatment with pregabalin was associated with creatine kinase elevations. Mean changes in creatine kinase from baseline to the maximum values were 60 U/L for pregabalin-treated patients and 28 U/L for the placebo patients. In all controlled trials across multiple patient populations, 2% of patients on pregabalin and

1% of placebo patients had a value of creatine kinase at least three times the upper limit of normal. Three pregabalin-treated patients had events reported as rhabdomyolysis in pre-marketing clinical trials. The relationship between these myopathy events and pregabalin is not completely understood because the cases had documented factors that may have caused or contributed to these events. Pregabalin should be discontinued if myopathy is diagnosed or suspected or if markedly elevated creatine kinase levels occur in the context of symptoms of myopathy.
Suicidal ideation and behaviour have been reported in patients treated with anti-epileptic agents in several indications. A meta-analysis of randomised placebo-controlled trials of anti-epileptic drugs has also shown a small increased risk of suicidal ideation and behaviour. The mechanism of this risk is not known and the available data do not exclude the possibility of an increased risk for pregabalin.
Therefore, patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.
Siranalen contains lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.
Siranalen contains carmoisine (E122) that may cause allergic reactions.
There is evidence from case reports, human studies, and animal studies associating pregabalin with serious, life-threatening, or fatal respiratory depression when co-administered with CNS depressants, including opioids, or in the setting of underlying respiratory impairment. When the decision is made to co-prescribe pregabalin with another CNS depressant, particularly an opioid, or to prescribe pregabalin to patients with underlying respiratory impairment, monitor patients for symptoms of respiratory depression and sedation, and consider initiating pregabalin at a low dose. The management of respiratory depression may include close observation, supportive measures, and reduction or withdrawal of CNS depressants (including pregabalin).
There is more limited evidence from case reports, animal studies, and human studies associating pregabalin with serious respiratory depression, without co-administered CNS depressants or without underlying respiratory impairment. Patients with renal impairment might be at higher risk of experiencing this severe adverse reaction.

Women of child-bearing potential/Contraception
Pregabalin use in the first trimester of pregnancy may cause major birth defects in the unborn child. Pregabalin should not be used during pregnancy unless the benefit to the mother clearly outweighs the potential risk to the fetus. Women of child-bearing potential must use effective contraception during treatment (see Section 4.6 Fertility, pregnancy and lactation).

4.5 Interaction with other medicinal products and other forms of interaction
Since pregabalin is predominantly excreted unchanged in the urine, undergoes negligible metabolism in humans (<2% of a dose recovered in urine as metabolites), does not inhibit drug metabolism in vitro, and is not bound to plasma proteins, it is unlikely to produce, or be subject to, pharmacokinetic interactions. Accordingly, in in vivo studies no clinically relevant pharmacokinetic interactions were observed between pregabalin and phenytoin, carbamazepine, valproic acid, lamotrigine, gabapentin, lorazepam, oxycodone or ethanol. Population pharmacokinetic analysis indicated that oral antidiabetics, diuretics, insulin, phenobarbital, tiagabine and topiramate had no clinically significant effect on pregabalin clearance.
Co-administration of pregabalin with the oral contraceptives norethisterone and/or ethinyl oestradiol does not influence the steady-state pharmacokinetics of either substance.
Pregabalin may potentiate the effects of ethanol and lorazepam. In controlled clinical trials, multiple oral doses of pregabalin co-administered with oxycodone, lorazepam, or ethanol did not result in clinically important effects on respiration. Pregabalin appears to be additive in the impairment of cognitive and gross motor function caused by oxycodone.
In the post-marketing experience, there are reports of respiratory failure and coma in patients taking pregabalin and other CNS depressant medications. There are post-marketing reports of events related to reduced lower gastrointestinal tract function (e.g., intestinal obstruction, paralytic ileus, constipation) when pregabalin was co-administered with medications that have the potential to produce constipation, such as opioid analgesics.
No specific pharmacodynamic interaction studies were conducted in elderly volunteers.
In the post-marketing experience, there are reports of respiratory failure, coma and deaths in patients taking pregabalin and other CNS depressant medications, including in patients who are substance abusers.

4.6 Fertility, pregnancy and lactation
Pregnancy
There is a limited amount of data on the use of pregabalin in pregnant women. Data from an observational study, which included more than 2,700 pregnancies exposed to pregabalin based on routinely collected data from administrative and medical registers in Denmark, Finland, Norway, and Sweden, is as follows:
Major congenital malformations (MCM)
The risk of MCM among the paediatric population exposed to pregabalin in the first trimester was slightly higher compared to unexposed population (adjusted prevalence ratio and 95% confidence interval: 1.14 (0.96-1.35)), and compared to population exposed to lamotrigine (1.29 (1.01-1.65)) or to duloxetine (1.39 (1.07-1.82)).
Birth and post-natal neurodevelopmental outcomes
There were no statistically significant findings for stillbirth, low birth weight, preterm birth, small for gestational age (SGA), low Apgar score at 5 minutes, and microcephaly. Adjusted prevalence ratios (aPRs, and 95% confidence intervals) results for the meta-analysis for stillbirth, low birth weight, preterm birth, SGA, low Apgar score at 5 minutes, and microcephaly for pregabalin-exposed compared to unexposed to antiepileptic drugs (AEDs) of 1.72 (1.02-2.91), 1.05 (0.91-1.21), 1.13 (0.99-1.29), 1.21 (1.01-1.44), 1.18 (0.95-1.48), and 1.09 (0.88-1.36) respectively.
In paediatric population exposed in utero, the study did not provide evidence of an increased risk for attention deficit hyperactivity disorder (ADHD), autism spectrum disorders (ASD), and intellectual disabilities (ID). In the meta-analyses of the neurodevelopmental outcomes (ADHD, ASD, and ID), results for ADHD, ASD, and ID for pregabalin-exposed compared to unexposed to AEDs were 1.32 (1.04-1.67), 1.00 (0.68-1.47), and 1.03 (0.80-1.32) respectively.
Studies in animals have shown reproductive toxicity. Pregabalin should not be used during pregnancy unless the benefit to the mother clearly outweighs the potential risk to the fetus. Effective contraception must be used in women of child-bearing potential.

Breastfeeding
Pregabalin is excreted in the milk of lactating women (see section 5.2). As the safety of pregabalin in infants is not known, breast-feeding is not recommended during treatment with pregabalin. A decision must be made whether to discontinue breast-feeding or to discontinue from pregabalin therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.
4.7 Effects on ability to drive and use machines
Pregabalin may cause dizziness and somnolence and therefore may influence the ability to drive or use machines. Patients are advised not to drive, operate complex machinery or engage in other potentially hazardous activities until it is known whether this medicinal product affects their ability to perform these activities.
4.8 Undesirable effects
The pregabalin clinical program involved over 12,000 patients who were exposed to pregabalin, of whom over 7000 were in double-blind placebo controlled trials. The most commonly reported adverse reactions were dizziness and somnolence. Adverse reactions were usually mild to moderate in intensity. In all controlled studies, the discontinuation rate due to adverse reactions was 14% for patients receiving pregabalin and 5% for patients receiving placebo. The most common adverse reactions resulting in discontinuation from pregabalin treatment groups were dizziness and somnolence.
Selected adverse drug reactions that were treatment related in the pooled analysis of clinical trials, are listed in the table below by System Organ Class (SOC). The frequency of these terms has been based on all-causality adverse drug reactions in the clinical trial data set (very common (≥1/10), common (≥1/100, <1/10), uncommon (≥1/1000, <1/100) and rare (<1/1000)).
The adverse reactions listed may also be associated with the underlying disease and/or concomitant medications.

System Organ Class	Adverse drug reactions
Infections and infestations	
Common	Nasopharyngitis
Blood and lymphatic system disorders	
Uncommon	Neutropenia
Metabolism and nutrition disorders	
Common	Appetite increased
Uncommon	Anorexia, hypoglycaemia
Psychiatric disorders	
Common	Euphoric mood, confusion, irritability, depression, disorientation, insomnia, libido decreased
Uncommon	Hallucination, restlessness, agitation, depressed mood, elevated mood, mood swings, depersonalization, abnormal dreams, word finding difficulty, libido increased, anorexia
Rare	Panic attack, disinhibition, apathy
Nervous system disorders	
Very Common	Dizziness, somnolence
Common	Ataxia, co-ordination abnormal, tremor, dysarthria, amnesia, memory impairment, disturbance in attention, paraesthesia, hypoesthesia, sedation, balance disorder, lethargy

Uncommon	Syncope, myoclonus, psychomotor hyperactivity, dyskinesia, dizziness postural, intention tremor, nystagmus, cognitive disorder, speech disorder, hyperreflexia, hyperaesthesia, burning sensation
Rare	Stupor, parosmia, hypokinesia, ageusia, dysgraphia
Eye disorders	
Common	Vision blurred, diplopia
Uncommon	Peripheral vision loss, visual disturbance, eye swelling, visual field defect, visual acuity reduced, eye pain, asthenopia, photopsia, dry eye, lacrimation increased, eye irritation
Rare	Oscillopsia, altered visual depth perception, mydriasis, strabismus, visual brightness
Ear and labyrinth disorders	
Common	Vertigo
Uncommon	Hyperacusis
Cardiac disorders	
Uncommon	Tachycardia, atrioventricular block first degree, sinus bradycardia
Rare	Sinus tachycardia, sinus arrhythmia
Vascular disorders	
Uncommon	Hypotension, hypertension, hot flushes, flushing, peripheral coldness
Respiratory, thoracic and mediastinal disorders	
Uncommon	Dyspnoea, epistaxis, cough, nasal congestion, rhinitis, snoring
Rare	Throat tightness, nasal dryness
Gastrointestinal disorders	
Common	Vomiting, constipation, flatulence, abdominal distension, dry mouth
Uncommon	Gastroesophageal reflux disease, salivary hypersecretion, hypoaesthesia oral
Rare	Ascites, pancreatitis, dysphagia
Skin and subcutaneous tissue disorders	
Uncommon	Rash papular, urticaria, sweating
Rare	Cold sweat
Musculoskeletal and connective tissue disorders	
Common	Muscle cramp, arthralgia, back pain, pain in limb, cervical spasm
Uncommon	Joint swelling, myalgia, muscle twitching, neck pain, muscle stiffness
Rare	Rhabdomyolysis
Renal and urinary disorders	
Uncommon	Urinary incontinence, dysuria
Rare	Renal failure, oliguria
Reproductive system and breast disorders	
Uncommon	Erectile dysfunction, sexual dysfunction, ejaculation delayed, dysmenorrhoea
Rare	Breast pain, amenorrhoea, breast discharge, breast enlargement
General disorders and administration site conditions	
Common	Oedema peripheral, oedema, gait abnormal, fall, feeling drunk, feeling abnormal, fatigue
Uncommon	Generalised oedema, chest tightness, pain, pyrexia, thirst, chills asthenia
Investigations	
Common	Weight increased
Uncommon	Blood creatine phosphokinase increased, alanine aminotransferase increased, aspartate aminotransferase increased, blood glucose increased, platelet count decreased, blood potassium decreased, weight decreased
Rare	White blood cell count decreased, blood creatinine increased

The following adverse drug reactions were reported during post-marketing surveillance:
Immune system disorder
Uncommon: Hypersensitivity.
Rare: Angioedema, allergic reaction.
Nervous system disorders
Very Common: Headache.
Uncommon: Loss of consciousness, mental impairment.
Eye disorders
Rare: Keratitis\$.
Cardiac disorders
Rare: Congestive heart failure.
Respiratory, thoracic and mediastinal disorders
Rare: Pulmonary oedema\$.
Gastrointestinal disorders
Common: Nausea, diarrhea.
Rare: Swollen tongue.
Skin and subcutaneous tissue disorders
Uncommon: Face swelling, pruritus.
Rare: Stevens-Johnson syndrome
Renal and urinary disorders
Rare: Urinary retention.
Reproductive system and breast disorders
Rare: Gynaecomastia\$.
General disorders and administration site conditions
Uncommon: Malaise.
\$ Adverse drug reaction frequency estimated using “The Rule of 3”.

4.9 Overdose
In overdoses up to 15 g, no unexpected adverse reactions were reported.
In the post-marketing experience, the most commonly reported adverse events observed when pregabalin was taken in overdose included affective disorder, somnolence, confusional state, depression, agitation, and restlessness. Seizures were also reported. Treatment of pregabalin overdose should include general supportive measures and may include hemodialysis if necessary (see Section 4.2).

5. PHARMACOLOGICAL PROPERTIES
5.1 Pharmacodynamic properties
Pharmacotheapeutic group: Antiepileptics, other antiepileptics
ATC code: N03AX16
The active substance, pregabalin, is a gamma-aminobutyric acid analogue ((S)-3-(aminomethyl)-5-methylhexanoic acid).
Mechanism of action
Pregabalin binds to an auxiliary subunit (α2-δ protein) of voltage-gated calcium channels in the neuronal system.
Evidence from animal models with nerve damage has shown that pregabalin reduces calcium dependent release of pronociceptive neurotransmitters in the spinal cord possibly by disrupting calcium trafficking and/or reducing calcium currents. Evidence from other animal models of nerve damage suggests the antinociceptive activities of pregabalin may also be mediated through interactions with the descending noradrenergic and serotonergic pathways.
Clinical efficacy and safety
Neuropathic pain
Efficacy has been shown in studies in diabetic neuropathy and post-herpetic neuralgia. Efficacy has not been studied in other models of neuropathic pain.
Pregabalin has been studied in 9 controlled clinical studies of up to 13 weeks with twice a day dosing and up to 8 weeks with three times a day dosing. Overall, the safety and efficacy profiles for twice a day and three times a day dosing regimens were similar.
In clinical trials up to 13 weeks, a reduction in pain was seen by Week 1 and was maintained throughout the treatment period.
In controlled clinical trials 35% of the pregabalin treated patients and 18% of the patients on placebo had a 50% improvement in pain score. For patients, not experiencing somnolence, such an improvement was observed in 33% of patients treated with pregabalin and 18% of patients on placebo. For patients who experienced somnolence the responder rates were 48% on pregabalin and 16% on placebo.
Epilepsy
Pregabalin has been studied in 3 controlled clinical trials of 12-week duration with either twice a day dosing or three times a day dosing. Overall, the safety and efficacy profiles for twice a day and three times a day dosing regimens were similar.
A reduction in seizure frequency was observed by Week 1.
Generalised Anxiety Disorder
Pregabalin has been studied in 6 controlled studies of 4 to 6 weeks duration, an elderly study of 8 weeks duration and a long-term relapse prevention study with a double-blind relapse prevention phase of 6 months duration.
Relief of the symptoms of GAD as reflected by the Hamilton Anxiety Rating Scale (HAM-A) was observed by Week 1.
In controlled clinical trials (4-8 weeks duration), 52% of the pregabalin-treated patients and 38% of the patients on placebo had at least a 50% improvement in HAM-A total score from baseline to endpoint.

Fibromyalgia
Pregabalin as monotherapy has been studied in 5 placebo-controlled studies, three of 12 weeks fixed-dose duration, one of 7 weeks fixed-dose duration, and a 6-month study demonstrating long-term efficacy. Pregabalin treatment in all fixed-dose studies produced a significant reduction in pain associated with fibromyalgia at doses from 300 to 600 mg per day (BID).
In the three 12-week fixed-dose studies, 40% of pregabalin-treated patients experienced a 30% or more improvement in pain score versus 28% of the patients on placebo; 23% of treated patients experienced a 50% or more improvement in pain score versus 15% of the patients on placebo.
Pregabalin produced significantly superior global assessment scores via the Patient Global Impression of Change (PGIC) in the three 12-week fixed-dose studies as compared to placebo treatment (41% patients feeling very much or much improved on pregabalin versus 29% on placebo). As measured by Fibromyalgia Impact Questionnaire (FIQ), pregabalin produced a statistically

significant improvement in function versus placebo treatment in 2 out of the 3 fixed-dose studies in which it was evaluated.
Pregabalin treatment produced significant improvements in patient-reported sleep outcomes in the 4 fixed-dose studies as measured by Medical Outcomes Study Sleep Scale (MOS-SS) Sleep disturbance subscale, MOS-SS overall sleep problem index, and the daily sleep quality diary.
In the 6-month study, improvement in pain, global assessment (PGIC), function (FIQ total score) and sleep (MOS-SS Sleep disturbance subscale) were maintained for pregabalin-treated patients for a significantly longer period compared to placebo.
Pregabalin 600 mg per day showed an additional improvement in patient-reported sleep outcomes as compared to 300 and 450 mg per day; mean effects on pain, global assessment, and FIQ were similar at 450 and 600 mg per day, although the 600 mg per day dose was less well tolerated.

5.2 Pharmacokinetic properties
Pregabalin steady-state pharmacokinetics are similar in healthy volunteers, patients with epilepsy receiving anti-epileptic drugs and patients with chronic pain.
Absorption
Pregabalin is rapidly absorbed when administered in the fasted state, with peak plasma concentrations occurring within 1 hour following both single and multiple dose administration. Pregabalin oral bioavailability is estimated to be ≥ 90% and is independent of dose. Following repeated administration, steady state is achieved within 24 to 48 hours. The rate of pregabalin absorption is decreased when given with food resulting in a decrease in C_{max} by approximately 25-30% and a delay in t_{max} to approximately 2.5 hours. However, administration of pregabalin with food has no clinically significant effect on the extent of pregabalin absorption.
Distribution
In preclinical studies, pregabalin has been shown to cross the blood brain barrier in mice, rats, and monkeys. Pregabalin has been shown to cross the placenta in rats and is present in the milk of lactating rats. In humans, the apparent volume of distribution of pregabalin following oral administration is approximately 0.56 l/kg. Pregabalin is not bound to plasma proteins.
Biotransformation
Pregabalin undergoes negligible metabolism in humans. Following a dose of radiolabelled pregabalin, approximately 98% of the radioactivity recovered in the urine was unchanged pregabalin. The N-methylated derivative of pregabalin, the major metabolite of pregabalin found in urine, accounted for 0.9% of the dose. In preclinical studies, there was no indication of racemisation of pregabalin S-enantiomer to the R-enantiomer.
Elimination
Pregabalin is eliminated from the systemic circulation primarily by renal excretion as unchanged drug.
Pregabalin mean elimination half-life is 6.3 hours. Pregabalin plasma clearance and renal clearance are directly proportional to creatinine clearance (see section 5.2).

Dose adjustment in patients with reduced renal function or undergoing haemodialysis is necessary (see section 4.2 Table 1).
Linearity/non-linearity
Pregabalin pharmacokinetics are linear over the recommended daily dose range. Inter-subject pharmacokinetic variability for pregabalin is low (<20%). Multiple dose pharmacokinetics are predictable from single-dose data. Therefore, there is no need for routine monitoring of plasma concentrations of pregabalin.

Gender
Clinical trials indicate that gender does not have a clinically significant influence on the plasma concentrations of pregabalin.
Renal impairment
Pregabalin clearance is directly proportional to creatinine clearance. In addition, pregabalin is effectively removed from plasma by haemodialysis (following a 4 hour haemodialysis treatment plasma pregabalin concentrations are reduced by approximately 50%). Because renal elimination is the major elimination pathway, dose reduction in patients with renal impairment and dose supplementation following haemodialysis is necessary (see section 4.2 Table 1).
Hepatic impairment
No specific pharmacokinetic studies were carried out in patients with impaired liver function. Since pregabalin does not undergo significant metabolism and is excreted predominantly as unchanged drug in the urine, impaired liver function would not be expected to significantly alter pregabalin plasma concentrations.
Elderly (over 65 years of age)
Pregabalin clearance tends to decrease with increasing age. This decrease in pregabalin oral clearance is consistent with decreases in creatinine clearance associated with increasing age. Reduction of pregabalin dose may be required in patients who have age related compromised renal function (see section 4.2 Table 1).
Breast-feeding mothers
The pharmacokinetics of 150 mg pregabalin given every 12 hours (300 mg daily dose) was evaluated in 10 lactating women who were at least 12 weeks postpartum. Lactation had little to no influence on pregabalin pharmacokinetics. Pregabalin was excreted into breast milk with average steady-state concentrations approximately 76% of those in maternal plasma. The estimated average daily infant dose of pregabalin from breast milk (assuming mean milk consumption of 150 mL/kg/day) was 0.31 mg/kg/day, which on a mg/kg basis would be approximately 7% of the maternal dose.

5.3 Preclinical safety data
In conventional safety pharmacology studies in animals, pregabalin was well-tolerated at clinically relevant doses. In repeated dose toxicity studies in rats and monkeys CNS effects were observed, including hypoactivity, hyperactivity and ataxia. An increased incidence of retinal atrophy commonly observed in aged albino rats was seen after long term exposure to pregabalin at exposures ≥ 5 times the mean human exposure at the maximum recommended clinical dose.
Teratogenicity
Pregabalin was not teratogenic in mice, rats or rabbits. Foetal toxicity in rats and rabbits occurred only at exposures sufficiently above human exposure. In prenatal/postnatal toxicity studies, pregabalin induced offspring developmental toxicity in rats at exposures >2 times the maximum recommended human exposure.
Mutagenicity
Pregabalin is not genotoxic based on results of a battery of *in vitro* and *in vivo* tests.
Carcinogenicity
Two-year carcinogenicity studies with pregabalin were conducted in rats and mice. No tumours were observed in rats at exposures up to 24 times the mean human exposure at the maximum recommended clinical dose of 600 mg/day. In mice, no increased incidence of tumours was found at exposures similar to the mean human exposure, but an increased incidence of haemangiosarcoma was observed at higher exposures. The non-genotoxic mechanism of pregabalin-induced tumour formation in mice involves platelet changes and associated endothelial cell proliferation. These platelet changes and endothelial cell proliferation were observed in mice not present in rats or in humans based on short term and limited long term clinical data. There is no evidence to suggest an associated risk to humans.
In juvenile rats the types of toxicity do not differ qualitatively from those observed in adult rats. However, juvenile rats are more sensitive. At therapeutic exposures, there was evidence of CNS clinical signs of hyperactivity and bruxism and some changes in growth (transient body weight gain suppression). Effects on the oestrus cycle were observed at 5-fold the human therapeutic exposure. Reduced acoustic startle response was observed in juvenile rats 1-2 weeks after exposure at >2 times (acoustic startle response) or >5 times (learning/memory) the human therapeutic exposure. Reduced acoustic startle response was observed in juvenile rats 1-2 weeks after exposure at >2 times the human therapeutic exposure. Nine weeks after exposure, this effect was no longer observable.

6. PHARMACEUTICAL PARTICULARS
6.1 List of excipients
Capule content:
Maize starch
Lactose monohydrate
Talc
Hard Capsule Shell:
Capsule body:
Gelatin
Titanium dioxide (E171)
Sodium lauryl sulphate
Capsule cap of 75 mg hard capsule:
Gelatin
Titanium dioxide (E171)
Patent blue V (E131)
Sodium lauryl sulphate
Capsule cap of 150 mg hard capsule:
Gelatin
Titanium dioxide (E171)
Sodium lauryl sulphate
Capsule cap of 300 mg hard capsule:
Gelatin
Titanium dioxide (E 171)
Erythrosine (E 127)
Carmoisine (E122)
Brilliant blue FCF (E 133)
Sodium lauryl sulphate
6.2 Incompatibilities
Not applicable.
6.3 Shelf life
36 months
6.4 Special precautions for storage
Store below 30°C in the original package
6.5 Nature and contents of container
PVC/Al blisters and PVC/PVDC/Al blisters containing 14, 21, 30, 50, 56, 84, 100 or 112 capsules.
Not all strengths and pack sizes may be marketed.
6.6 Special precautions for disposal and other handling
No special requirements for disposal.

7. MARKETING AUTHORISATION HOLDER
Medochemie Singapore Pte Ltd,
221 Henderson Road, #06-04 Henderson Building,
Singapore 159557

8. DATE OF REVISION OF THE TEXT
12/2022

CODE:.....
SINGAPORE

PHARMACODE

PHARMACODE