

FEDAC COMPOUND LINCTUS

COMPOSITION:

| | |
|------------------------|-------------|
| Tripolidine HCl BP | 1.25 mg/5ml |
| Pseudoephedrine HCl BP | 30 mg/5ml |
| Codeine Phosphate BP | 10 mg/5ml |

Excipients:

Propyl Hydroxybenzoate, Methyl Hydroxybenzoate, sugar, sodium saccharin, citric acid, sodium citrate dihydrate, strawberry flavour, glycerin, sodium carboxymethylcellulose, carmoisine water soluble and purified water.

INDICATIONS:

Temporary relief of coughs and upper respiratory symptoms including nasal congestion associated with allergy or the common cold.

RECOMMENDED DOSAGE:

Adults and children over 12 years: 2 teaspoonfuls 3 times a day

Children and Adolescents: Codeine should not be used in children below 12 years old. In those 12 years old and above, codeine may be used for the treatment of unproductive cough which is not relieved by other anti-tussives alone. The lowest effective dose should be used for the shortest possible duration.

SIDE EFFECTS:

Drowsiness, disturbed coordination, dryness of mouth, nose and throat.

CONTRA-INDICATIONS:

In patients intolerant to Pseudoephedrine HCl or Tripolidine HCl or Codeine Phosphate. Patients with severe hypertension, severe coronary artery disease and in patients on MAO inhibitors. FEDAC COMPOUND LINCTUS should not be used in newborn or premature infants and should not be used to treat lower respiratory tract symptoms including asthma.

SPECIAL WARNINGS:

Caution is advised when codeine is used in patients with underlying respiratory conditions, including those with asthma and other chronic breathing problems.

Parents and caregivers of paediatric patients should be advised on the possible signs and symptoms of respiratory depression in their children, such as unusual sleepiness, confusion and difficult or noisy breathing, and to seek immediate medical attention if these are observed.

Nursing mothers should also be advised to exercise caution when taking codeine, since codeine's metabolite (morphine) may subsequently be found in the breast milk. If the infant shows signs of increased sleepiness, difficulty breastfeeding, breathing difficulties or limpness, immediate medical attention should be sought.

CAUTION:

In elderly or debilitated patients, in severe impairment of renal or hepatic function. This preparation may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcohol. Codeine may be habit forming when used over long periods or in high doses and may cause psychological dependence, physical dependence and tolerance.

STORAGE:

In airtight containers and store below 25°C, protect from light.

PACKING:

3.8L

DATE OF REVISION:

April 2022

For further information, please consult your doctor or pharmacist.

Manufactured by: ICM Pharma Pte Ltd
Blk 26, Kallang Place #05-08/12
Singapore 339157