

## **PACKAGE INSERT**

### **1. NAME OF THE MEDICINAL PRODUCT**

Tyroka 200 mg film-coated tablets

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film-coated tablet contains 200 mg pazopanib (as hydrochloride).

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Film-coated tablet.

Capsule-shaped, pink, film-coated tablet (14.5 mm length and 5.6 mm width) with “173” debossed on one side and plain on the other side.

### **4. CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

##### Renal cell carcinoma (RCC)

Pazopanib is indicated in adults for the first-line treatment of advanced renal cell carcinoma (RCC) and for patients who have received prior cytokine therapy for advanced disease.

##### Soft-tissue sarcoma (STS)

Pazopanib is indicated for the treatment of adult patients with selective subtypes of soft-tissue sarcoma (STS) who have received prior chemotherapy for metastatic disease or who have progressed within 12 months after (neo) adjuvant therapy.

Efficacy and safety has only been established in certain STS histological tumour subtypes (see section 5.1).

#### **4.2 Posology and method of administration**

Pazopanib treatment should only be initiated by a physician experienced in the administration of anti-cancer medicinal products.

##### Posology

###### *Adults*

The recommended dose of pazopanib for the treatment of RCC or STS is 800 mg once daily.

###### *Dose modifications*

Initial dose reduction should be from 800 mg to 400 mg daily. Subsequent dose modification (decrease or increase) should be in 200 mg decrements or increments in a stepwise fashion based on individual tolerability in order to manage adverse reactions. The daily dose of pazopanib should not exceed 800 mg.

###### *Paediatric population*

Pazopanib is not recommended for use in children and adolescents under 18 years (see sections 4.4, 5.1 and 5.3).

### *Elderly*

There are limited data on the use of pazopanib in patients aged 65 years and older. In the RCC studies of pazopanib, overall no clinically significant differences in safety of pazopanib were observed between subjects aged at least 65 years and younger subjects. Clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some elderly patients cannot be ruled out.

No alteration of dosage, dosing frequency or route of administration is required in patients over 65 years.

### *Renal impairment*

Renal impairment is unlikely to have a clinically relevant effect on pazopanib pharmacokinetics given the low renal excretion of pazopanib and metabolites (see section 5.2). Therefore, no dose adjustment is required in patients with creatinine clearance 30 ml/min and above. There is no experience of pazopanib in patients with severe renal impairment or in patients undergoing peritoneal dialysis or hemodialysis; therefore, use of pazopanib is not recommended in these patients.

### *Hepatic impairment*

Dosing recommendations in hepatically impaired patients are based on pharmacokinetic studies of pazopanib in patients with varying degrees of hepatic dysfunction (see section 5.2). All patients should have liver function tests to determine whether they have hepatic impairment before starting and during pazopanib therapy (see section 4.4). Administration of pazopanib to patients with mild or moderate hepatic impairment should be undertaken with caution and close monitoring of tolerability. 800 mg pazopanib once daily is the recommended dose in patients with mild abnormalities in serum liver tests (defined either as normal bilirubin and any degree of alanine aminotransferase (ALAT) elevation or as an elevation of bilirubin up to 1.5 x upper limit of normal (ULN) regardless of the ALAT value). A reduced pazopanib dose of 200 mg once daily is recommended in patients with moderate hepatic impairment (defined as an elevation of bilirubin >1.5 to 3 x ULN regardless of the ALAT value) (see section 5.2).

Pazopanib is not recommended in patients with severe hepatic impairment (defined as total bilirubin >3 x ULN regardless of the ALAT value).

See section 4.4 for liver monitoring and dose modification for patients with drug-induced hepatotoxicity.

### Method of administration

Pazopanib is for oral use. It should be taken without food, at least one hour before or two hours after a meal (see section 5.2). The film-coated tablets should be taken whole with water and not broken or crushed (see section 5.2).

If a dose is missed, it should not be taken if it is less than 12 hours until the next dose.

## **4.3 Contraindications**

Pazopanib is contraindicated in patients with severe hepatic impairment and hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### Hepatic effects

Cases of hepatic failure (including fatalities) have been reported during use of pazopanib.

Administration of pazopanib to patients with mild or moderate hepatic impairment should be undertaken with caution and close monitoring. 800 mg pazopanib once daily is the recommended dose in patients with mild abnormalities in serum liver tests (either normal bilirubin and any degree of ALAT elevation or elevation of bilirubin up to 1.5 x ULN regardless of the ALAT value). A reduced pazopanib dose of 200 mg once daily is recommended in patients with moderate hepatic impairment

(elevation of bilirubin >1.5 to 3 x ULN regardless of the ALAT value) (see sections 4.2 and 5.2). Pazopanib is not recommended in patients with severe hepatic impairment (total bilirubin >3 x ULN regardless of the ALAT value) (see sections 4.2 and 5.2). Exposure at a 200 mg dose is markedly reduced, though highly variable, in these patients, with values considered insufficient to obtain a clinically relevant effect.

In clinical studies with pazopanib, increase in serum transaminases (ALAT, aspartate aminotransferase [ASAT]) and bilirubin were observed (see section 4.8). In the majority of the cases, isolated increases in ALAT and ASAT have been reported, without concomitant elevations of alkaline phosphatase or bilirubin. Patients over 60 years of age may be at greater risk for mild (>3 x ULN) to severe (>8 x ULN) elevation of ALAT. Patients who carry the HLA-B\*57:01 allele have an increased risk of pazopanib-associated ALAT elevations. Liver function should be monitored in all subjects receiving pazopanib, regardless of genotype or age (see section 5.1).

Serum liver tests should be performed before initiation of treatment with pazopanib, at weeks 3, 5, 7 and 9, then at months 3 and 4, with additional tests as clinically indicated. Periodic testing should then continue after month 4.

See Table 1 for dose modification guidance for patients with baseline values of total bilirubin  $\leq$  1.5 x ULN and ALAT  $\leq$  2 x ULN:

**Table 1 Dose modifications for drug-induced hepatotoxicity**

Liver test values	Dose modification
ALAT elevation between 3 and 8 x ULN	Continue on pazopanib with weekly monitoring of liver function until ALAT return to Grade 1 or baseline.
ALAT elevation of >8 x ULN	Interrupt pazopanib until ALAT return to Grade 1 or baseline. If the potential benefit of reinitiating pazopanib treatment is considered to outweigh the risk for hepatotoxicity, then reintroduce pazopanib at a reduced dose of 400 mg daily and perform serum liver tests weekly for 8 weeks. Following reintroduction of pazopanib, if ALAT elevations > 3 x ULN recur, then pazopanib should be permanently discontinued.
ALAT elevations >3 x ULN concurrently with bilirubin elevations >2 x ULN	Permanently discontinue pazopanib. Patients should be monitored until return to Grade 1 or baseline. Pazopanib is a UGT1A1 inhibitor. Mild, indirect (unconjugated) hyperbilirubinaemia may occur in patients with Gilbert's syndrome. Patients with only a mild indirect hyperbilirubinaemia, known or suspected Gilbert's syndrome, and elevation in ALAT >3 x ULN should be managed as per the recommendations outlined for isolated ALAT elevations.

Concomitant use of pazopanib and simvastatin increases the risk of ALAT elevations (see section 4.5) and should be undertaken with caution and close monitoring.

### Hypertension

In clinical studies with pazopanib, events of hypertension including newly diagnosed symptomatic episodes of elevated blood pressure (hypertensive crisis) have occurred. Blood pressure should be well controlled prior to initiating pazopanib. Patients should be monitored for hypertension early after starting treatment (no longer than one week after starting pazopanib) and frequently thereafter to ensure blood pressure control. Elevated blood pressure levels (systolic blood pressure  $\geq 150$  mm Hg or diastolic blood pressure  $\geq 100$  mm Hg) occurred early in the course of treatment (approximately 40% of cases occurred by day 9 and approximately 90% of cases occurred in the first 18 weeks). Blood pressure should be monitored and managed promptly using a combination of anti-hypertensive therapy and dose modification of pazopanib (interruption and re-initiation at a reduced dose based on clinical judgement) (see sections 4.2 and 4.8). Pazopanib should be discontinued if there is evidence of hypertensive crisis or if hypertension is severe and persists despite anti-hypertensive therapy and pazopanib dose reduction. Serious cases of artery dissection have been reported in patients using VEGFR TKIs, including pazopanib, with or without hypertension.

### Posterior reversible encephalopathy syndrome (PRES)/Reversible posterior leukoencephalopathy syndrome (RPLS)

PRES/RPLS has been reported in association with pazopanib. PRES/RPLS can present with headache, hypertension, seizure, lethargy, confusion, blindness and other visual and neurological disturbances, and can be fatal. Patients developing PRES/RPLS should permanently discontinue treatment with pazopanib.

### Interstitial lung disease (ILD)/Pneumonitis

ILD, which can be fatal, has been reported in association with pazopanib (see section 4.8). Monitor patients for pulmonary symptoms indicative of ILD/pneumonitis and discontinue pazopanib in patients developing ILD or pneumonitis.

### Cardiac dysfunction/Heart failure

The risks and benefits of pazopanib should be considered before beginning therapy in patients who have pre-existing cardiac dysfunction. The safety and pharmacokinetics of pazopanib in patients with moderate to severe heart failure or those with a below normal left ventricular ejection fraction (LVEF) have not been studied.

In clinical studies with pazopanib, events of cardiac dysfunction such as congestive heart failure and decreased LVEF have occurred (see section 4.8). In a randomised study comparing pazopanib and sunitinib in RCC (VEG108844), subjects had baseline and follow up LVEF measurements. Myocardial dysfunction occurred in 13% (47/362) of subjects in the pazopanib arm compared to 11% (42/369) of subjects in the sunitinib arm. Congestive heart failure was observed in 0.5% of subjects in each treatment arm. Congestive heart failure was reported in 3 out of 240 subjects (1%) in the Phase III VEG110727 STS study. Decreases in LVEF in subjects who had post-baseline and follow-up LVEF measurement were detected in 11% (15/140) in the pazopanib arm, compared with 3% (1/39) in the placebo arm.

### *Risk factors*

Thirteen of the 15 subjects in the pazopanib arm of the STS Phase III study had concurrent hypertension which may have exacerbated cardiac dysfunction in patients at risk by increasing cardiac after-load. 99% of patients (243/246) enrolled in the STS Phase III study, including the 15 subjects, received anthracycline. Prior anthracycline therapy may be a risk factor for cardiac dysfunction.

### *Outcome*

Four of the 15 subjects had full recovery (within 5% of baseline) and 5 had partial recovery (within the normal range, but  $>5\%$  below baseline). One subject did not recover and follow-up data were not available for the other 5 subjects.

### Management

Interruption of pazopanib and/or dose reduction should be combined with treatment of hypertension (if present, refer to hypertension warning section above) in patients with significant reductions in LVEF, as clinically indicated.

Patients should be carefully monitored for clinical signs or symptoms of congestive heart failure. Baseline and periodic evaluation of LVEF is recommended in patients at risk of cardiac dysfunction.

### QT prolongation and torsade de pointes

In clinical studies with pazopanib, events of QT prolongation and torsade de pointes have occurred (see section 4.8). Pazopanib should be used with caution in patients with a history of QT interval prolongation, in patients taking antiarrhythmics or other medicinal products that may prolong QT interval and in patients with relevant pre-existing cardiac disease. When using pazopanib, baseline and periodic monitoring of electrocardiograms and maintenance of electrolytes (e.g. calcium, magnesium, potassium) within normal range is recommended.

### Arterial thrombotic events

In clinical studies with pazopanib, myocardial infarction, myocardial ischaemia, ischaemic stroke and transient ischaemic attack were observed (see section 4.8). Fatal events have been observed. Pazopanib should be used with caution in patients who are at increased risk of thrombotic events or who have had a history of thrombotic events. Pazopanib has not been studied in patients who have had an event within the previous 6 months. A treatment decision should be made based on the assessment of individual patient's benefit/risk.

### Venous thromboembolic events

In clinical studies with pazopanib, venous thromboembolic events including venous thrombosis and fatal pulmonary embolus have occurred. While observed in both RCC and STS studies, the incidence was higher in the STS population (5%) than in the RCC population (2%).

### Thrombotic microangiopathy (TMA)

TMA has been reported in clinical studies of pazopanib as monotherapy, in combination with bevacizumab, and in combination with topotecan (see section 4.8). Patients developing TMA should permanently discontinue treatment with pazopanib. Reversal of effects of TMA has been observed after treatment was discontinued. Pazopanib is not indicated for use in combination with other agents.

### Haemorrhagic events

In clinical studies with pazopanib haemorrhagic events have been reported (see section 4.8). Fatal haemorrhagic events have occurred. Pazopanib has not been studied in patients who had a history of haemoptysis, cerebral haemorrhage or clinically significant gastrointestinal (GI) haemorrhage in the past 6 months and should not be used in those patients. Pazopanib should be used with caution in patients with significant risk of haemorrhage.

### Aneurysms and artery dissections

The use of VEGF pathway inhibitors in patients with or without hypertension may promote the formation of aneurysm and/or artery dissections. Before initiating pazopanib, this risk should be carefully considered in patients with risk factors such as hypertension or history of aneurysm.

### Gastrointestinal (GI) perforations and fistula

In clinical studies with pazopanib, events of GI perforation or fistula have occurred (see section 4.8). Fatal perforation events have occurred. Pazopanib should be used with caution in patients at risk for GI perforation or fistula.

### Wound healing

No formal studies of the effect of pazopanib on wound healing have been conducted. Since vascular endothelial growth factor (VEGF) inhibitors may impair wound healing, treatment with pazopanib should be stopped at least 7 days prior to scheduled surgery. The decision to resume pazopanib after

surgery should be based on clinical judgement of adequate wound healing. Pazopanib should be discontinued in patients with wound dehiscence.

#### Hypothyroidism

In clinical studies with pazopanib, events of hypothyroidism have occurred (see section 4.8). Baseline laboratory measurement of thyroid function is recommended and patients with hypothyroidism should be treated as per standard medical practice prior to the start of pazopanib treatment. All patients should be observed closely for signs and symptoms of thyroid dysfunction on pazopanib treatment.

Laboratory monitoring of thyroid function should be performed periodically and managed as per standard medical practice.

#### Proteinuria

In clinical studies with pazopanib, proteinuria has been reported. Baseline and periodic urinalysis during treatment is recommended and patients should be monitored for worsening proteinuria.

Pazopanib should be discontinued if the patient develops nephrotic syndrome.

#### Tumour lysis syndrome (TLS)

The occurrence of TLS, including fatal TLS, has been associated with the use of pazopanib (see section 4.8). Patients at increased risk of TLS are those with rapidly growing tumours, a high tumour burden, renal dysfunction, or dehydration. Preventative measures, such as treatment of high uric acid levels and intravenous hydration, should be considered prior to initiation of pazopanib. Patients at risk should be closely monitored and treated as clinically indicated.

#### Pneumothorax

In clinical studies with pazopanib in advanced soft tissue sarcoma, events of pneumothorax have occurred (see section 4.8). Patients on pazopanib treatment should be observed closely for signs and symptoms of pneumothorax.

#### Paediatric population

Because the mechanism of action of pazopanib can severely affect organ growth and maturation during early post-natal development (see section 5.3), pazopanib should not be given to human paediatric patients.

#### Infections

Cases of serious infections (with or without neutropenia), in some cases with fatal outcome, have been reported.

#### Combination with other systemic anti-cancer therapies

Clinical studies of pazopanib in combination with pemetrexed (non-small cell lung cancer [NSCLC]), lapatinib (cervical cancer) or pembrolizumab (advanced renal cell carcinoma) were terminated early due to concerns over increased toxicity and/or mortality, and a safe and effective combination dose has not been established with these regimens. Pazopanib is not indicated for use in combination with other anti-cancer agents.

#### Pregnancy

Pre-clinical studies in animals have shown reproductive toxicity (see section 5.3). If pazopanib is used during pregnancy, or if the patient becomes pregnant whilst receiving pazopanib, the potential hazard to the foetus should be explained to the patient. Women of childbearing potential should be advised to avoid becoming pregnant while receiving treatment with pazopanib (see section 4.6).

#### Interactions

Concomitant treatment with strong inhibitors of CYP3A4, P-glycoprotein (P-gp) or breast cancer resistance protein (BCRP) should be avoided due to risk of increased exposure to pazopanib (see section 4.5). Selection of alternative concomitant medicinal products with no or minimal potential to inhibit CYP3A4, P-gp or BCRP should be considered.

Concomitant treatment with inducers of CYP3A4 should be avoided due to risk of decreased exposure to pazopanib (see section 4.5).

Cases of hyperglycaemia have been observed during concomitant treatment with ketoconazole.

Concomitant administration of pazopanib with uridine diphosphate glucuronosyl transferase 1A1 (UGT1A1) substrates (e.g. irinotecan) should be undertaken with caution since pazopanib is an inhibitor of UGT1A1 (see section 4.5).

Grapefruit juice should be avoided during treatment with pazopanib (see section 4.5).

#### Excipients

This medicinal product contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

### **4.5 Interaction with other medicinal products and other forms of interaction**

#### Effects of other medicinal products on pazopanib

*In vitro* studies suggested that the oxidative metabolism of pazopanib in human liver microsomes is mediated primarily by CYP3A4, with minor contributions from CYP1A2 and CYP2C8. Therefore, inhibitors and inducers of CYP3A4 may alter the metabolism of pazopanib.

#### *CYP3A4, P-gp, BCRP inhibitors*

Pazopanib is a substrate for CYP3A4, P-gp and BCRP.

Concurrent administration of pazopanib (400 mg once daily) with the strong CYP3A4 and P-gp inhibitor ketoconazole (400 mg once daily) for 5 consecutive days resulted in a 66% and 45% increase in mean pazopanib  $AUC_{(0-24)}$  and  $C_{max}$ , respectively, relative to administration of pazopanib alone (400 mg once daily for 7 days). Pharmacokinetic parameter comparisons of pazopanib  $C_{max}$  (range of means 27.5 to 58.1 µg/ml) and  $AUC_{(0-24)}$  (range of means 48.7 to 1040 µg\*h/ml) after administration of pazopanib 800 mg alone and after administration of pazopanib 400 mg plus ketoconazole 400 mg (mean  $C_{max}$  59.2 µg/ml, mean  $AUC_{(0-24)}$  1300 µg\*h/ml) indicated that, in the presence of a strong CYP3A4 and P-gp inhibitor a dose reduction to pazopanib 400 mg once daily will, in the majority of patients, result in systemic exposure similar to that observed after administration of 800 mg pazopanib once daily alone. Some patients however may have systemic pazopanib exposure greater than what has been observed after administration of 800 mg pazopanib alone.

Co-administration of pazopanib with other strong inhibitors of the CYP3A4 family (e.g. itraconazole, clarithromycin, atazanavir, indinavir, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin, voriconazole) may increase pazopanib concentrations. Grapefruit juice contains an inhibitor of CYP3A4 and may also increase plasma concentrations of pazopanib.

Administration of 1500 mg lapatinib (a substrate for and weak inhibitor of CYP3A4 and P-gp and a potent inhibitor of BCRP) with 800 mg pazopanib resulted in an approximately 50% to 60% increase in mean pazopanib  $AUC_{(0-24)}$  and  $C_{max}$  compared to administration of 800 mg pazopanib alone. Inhibition of P-gp and/or BCRP by lapatinib likely contributed to the increased exposure to pazopanib.

Co-administration of pazopanib with a CYP3A4, P-gp, and BCRP inhibitor, such as lapatinib, will result in an increase in plasma pazopanib concentrations. Co-administration with potent P-gp or BCRP inhibitors may also alter the exposure and distribution of pazopanib, including distribution into the central nervous systems (CNS).

Concomitant use of pazopanib with a strong CYP3A4 inhibitor should be avoided (see section 4.4). If no medically acceptable alternative to a strong CYP3A4 inhibitor is available, the dose of pazopanib should be reduced to 400 mg daily during concomitant administration. In such cases there should be close attention to adverse drug reaction, and further dose reduction may be considered if possible drug-related adverse events are observed.

Combination with strong P-gp or BCRP inhibitors should be avoided, or selection of an alternate concomitant medicinal product with no or minimal potential to inhibit P-gp or BCRP is recommended.

#### *CYP3A4, P-gp, BCRP inducers*

CYP3A4 inducers such as rifampin may decrease plasma pazopanib concentrations. Co-administration of pazopanib with potent P-gp or BCRP inducers may alter the exposure and distribution of pazopanib, including distribution into the CNS. Selection of an alternative concomitant medication with no or minimal enzyme or transporter induction potential is recommended.

#### Effects of pazopanib on other medicinal products

*In vitro* studies with human liver microsomes showed that pazopanib inhibited CYP enzymes 1A2, 3A4, 2B6, 2C8, 2C9, 2C19, and 2E1. Potential induction of human CYP3A4 was demonstrated in an *in vitro* human PXR assay. Clinical pharmacology studies, using pazopanib 800 mg once daily, have demonstrated that pazopanib does not have a clinically relevant effect on the pharmacokinetics of caffeine (CYP1A2 probe substrate), warfarin (CYP2C9 probe substrate), or omeprazole (CYP2C19 probe substrate) in cancer patients. Pazopanib resulted in an increase of approximately 30% in the mean AUC and  $C_{max}$  of midazolam (CYP3A4 probe substrate) and increases of 33% to 64% in the ratio of dextrometorphane to dextrophan concentrations in the urine after oral administration of dextromethorphan (CYP2D6 probe substrate). Co-administration of pazopanib 800 mg once daily and paclitaxel 80 mg/m<sup>2</sup> (CYP3A4 and CYP2C8 substrate) once weekly resulted in a mean increase of 26% and 31% in paclitaxel AUC and  $C_{max}$ , respectively.

Based on *in vitro* IC<sub>50</sub> and *in vivo* plasma  $C_{max}$  values, pazopanib metabolites GSK1268992 and GSK1268997 may contribute to the net inhibitory effect of pazopanib towards BCRP. Furthermore, inhibition of BCRP and P-gp by pazopanib in the gastrointestinal tract cannot be excluded. Care should be taken when pazopanib is co-administered with other oral BCRP and P-gp substrates.

*In vitro*, pazopanib inhibited human organic anion transporting polypeptide (OATP1B1). It cannot be excluded that pazopanib will affect the pharmacokinetics of substrates of OATP1B1 (e.g. statins, see “Effect of concomitant use of pazopanib and simvastatin” below).

Pazopanib is an inhibitor of the uridine diphosphoglucuronosyl-transferase 1A1 (UGT1A1) enzyme *in vitro*. The active metabolite of irinotecan, SN-38, is a substrate for OATP1B1 and UGT1A1. Co-administration of pazopanib 400 mg once daily with cetuximab 250 mg/m<sup>2</sup> and irinotecan 150 mg/m<sup>2</sup> resulted in an approximately 20% increase in systemic exposure to SN-38. Pazopanib may have a greater impact on SN-38 disposition in subjects with the UGT1A1\*28 polymorphism relative to subjects with the wild-type allele. However, the UGT1A1 genotype was not always predictive of the effect of pazopanib on SN-38 disposition. Care should be taken when pazopanib is co-administered with substrates of UGT1A1.

#### Effect of concomitant use of pazopanib and simvastatin

Concomitant use of pazopanib and simvastatin increases the incidence of ALAT elevations. Results from a meta-analysis using pooled data from clinical studies with pazopanib show that ALAT >3x ULN was reported in 126/895 (14%) of patients who did not use statins, compared with 11/41 (27%) of patients who had concomitant use of simvastatin (p=0.038). If a patient receiving concomitant simvastatin develops ALAT elevations, follow guidelines for pazopanib dosology and discontinue simvastatin (see section 4.4). In addition, concomitant use of pazopanib and other statins should be undertaken with caution as there are insufficient data available to assess their impact on ALAT levels. It cannot be excluded that pazopanib will affect the pharmacokinetics of other statins (e.g. atorvastatin, fluvastatin, pravastatin, rosuvastatin).

#### Effect of food on pazopanib

Administration of pazopanib with a high-fat or low-fat meal results in an approximately 2-fold increase in AUC and  $C_{max}$ . Therefore, pazopanib should be administered at least 1 hour before or 2 hours after a meal.



#### Medicinal products that raise gastric pH

Concomitant administration of pazopanib with esomeprazole decreases the bioavailability of pazopanib by approximately 40% (AUC and  $C_{max}$ ), and co-administration of pazopanib with medicines that increase gastric pH should be avoided. If the concomitant use of a proton-pump inhibitor (PPI) is medically necessary, it is recommended that the dose of pazopanib be taken without food once daily in the evening concomitantly with the PPI. If the concomitant administration of an  $H_2$ -receptor antagonist is medically necessary, pazopanib should be taken without food at least 2 hours before or at least 10 hours after a dose of an  $H_2$ -receptor antagonist. Pazopanib should be administered at least 1 hour before or 2 hours after administration of short-acting antacids. The recommendations for how PPIs and  $H_2$ -receptor antagonists are co-administered are based on physiological considerations.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy/Contraception in males and females

There are no adequate data from the use of pazopanib in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). The potential risk for humans is unknown.

Pazopanib should not be used during pregnancy unless the clinical condition of the woman requires treatment with pazopanib. If pazopanib is used during pregnancy, or if the patient becomes pregnant while receiving pazopanib, the potential hazard to the foetus should be explained to the patient.

Women of childbearing potential should be advised to use adequate contraception during treatment and for at least 2 weeks after the last dose of pazopanib and to avoid becoming pregnant while receiving treatment with pazopanib.

Male patients (including those who have had vasectomies) should use condoms during sexual intercourse while taking pazopanib and for at least 2 weeks after the last dose of pazopanib to avoid potential exposure to the medicinal product for pregnant partners and female partners of reproductive potential.

#### Breast-feeding

The safe use of pazopanib during breast-feeding has not been established. It is not known whether pazopanib or its metabolites are excreted in human milk. There are no animal data on the excretion of pazopanib in animal milk. A risk to the breast-fed child cannot be excluded. Breast-feeding should be discontinued during treatment with pazopanib.

#### Fertility

Animal studies indicate that male and female fertility may be affected by treatment with pazopanib (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

pazopanib has no or negligible influence on the ability to drive and use machines. A detrimental effect on such activities cannot be predicted from the pharmacology of pazopanib. The clinical status of the patient and the adverse event profile of pazopanib should be borne in mind when considering the patient's ability to perform tasks that require judgement, motor or cognitive skills. Patients should avoid driving or using machines if they feel dizzy, tired or weak.

### **4.8 Undesirable effects**

#### Summary of the safety profile

Pooled data from the pivotal RCC study (VEG105192, n=290), the extension study (VEG107769, n=71), the supportive Phase II study (VEG102616, n=225) and the randomised, open-label, parallel group Phase III non-inferiority study (VEG108844, n=557) were evaluated in the overall evaluation of safety and tolerability of pazopanib (total n=1149) in subjects with RCC (see section 5.1).

Pooled data from the pivotal STS study (VEG110727, n=369) and the supportive Phase II study (VEG20002, n=142) was evaluated in the overall evaluation of safety and tolerability of pazopanib (total safety population n=382) in subjects with STS (see section 5.1).

The most important serious adverse reactions identified in the RCC or STS studies were transient ischaemic attack, ischaemic stroke, myocardial ischaemia, myocardial and cerebral infarction, cardiac dysfunction, gastrointestinal perforation and fistula, QT prolongation, Torsade de Pointes and pulmonary, gastrointestinal and cerebral haemorrhage, all adverse reactions being reported in <1% of treated patients. Other important serious adverse reactions identified in STS studies included venous thromboembolic events, left ventricular dysfunction and pneumothorax.

Fatal events that were considered possibly related to pazopanib included gastrointestinal haemorrhage, pulmonary haemorrhage/haemoptysis, abnormal hepatic function, intestinal perforation and ischaemic stroke.

The most common adverse reactions (experienced by at least 10% of the patients) of any grade in the RCC and STS trials included: diarrhoea, hair colour change, skin hypopigmentation, exfoliative rash, hypertension, nausea, headache, fatigue, anorexia, vomiting, dysgeusia, stomatitis, weight decreased, pain, elevated alanine aminotransferase and elevated aspartate aminotransferase.

Adverse drug reactions, all grades, which were reported in RCC and STS subjects or during the post-marketing period are listed below by MedDRA body system organ class, frequency and grade of severity. The following convention has been utilised for the classification of frequency: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); and not known (cannot be estimated from the available data).

Categories have been assigned based on absolute frequencies in the clinical trial data. Post-marketing data on safety and tolerability across all pazopanib clinical studies and from spontaneous reports have also been evaluated. Within each system organ class, adverse reactions with the same frequency are presented in order of decreasing seriousness.

#### Tabulated list of adverse reactions

**Table 2 Treatment-related adverse reactions reported in RCC studies (n=1149) or during post-marketing period**

System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
<b>Infections and Infestations</b>	Common	Infections (with or without neutropenia) <sup>†</sup>	not known	not known	not known
	Uncommon	Gingival infection	1 (<1%)	0	0
		Infectious peritonitis	1 (<1%)	0	0
<b>Neoplasms benign, malignant and unspecified (incl. cysts and polyps)</b>	Uncommon	Tumour pain	1 (<1%)	1 (<1%)	0
<b>Blood and lymphatic system disorders</b>	Common	Thrombocytopenia	80 (7%)	10 (<1%)	5 (<1%)
		Neutropenia	79 (7%)	20 (2%)	4 (<1%)
		Leukopenia	63 (5%)	5 (<1%)	0
	Uncommon	Polycythaemia	6 (0.03%)	1	0

System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
	Rare	Thrombotic microangiopathy (including thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome)†	not known	not known	not known
<b>Endocrine disorders</b>	Common	Hypothyroidism	83 (7%)	1 (<1%)	0
<b>Metabolism and nutrition disorders</b>	Very common	Decreased appetite <sup>e</sup>	317 (28%)	14 (1%)	0
	Common	Hypophosphataemia	21 (2%)	7 (<1%)	0
		Dehydration	16 (1%)	5 (<1%)	0
	Uncommon	Hypomagnesaemia	10 (<1%)	0	0
	Not known	Tumour lysis syndrome*	not known	not known	not known
<b>Psychiatric disorders</b>	Common	Insomnia	30 (3%)	0	0
<b>Nervous system disorders</b>	Very common	Dysgeusia <sup>c</sup>	254 (22%)	1 (<1%)	0
		Headache	122 (11%)	11 (<1%)	0
	Common	Dizziness	55 (5%)	3 (<1%)	1 (<1%)
		Lethargy	30 (3%)	3 (<1%)	0
		Paraesthesia	20 (2%)	2 (<1%)	0
		Peripheral sensory neuropathy	17 (1%)	0	0
	Uncommon	Hypoaesthesia	8 (<1%)	0	0
		Transient ischaemic attack	7 (<1%)	4 (<1%)	0
		Somnolence	3 (<1%)	1 (<1%)	0
		Cerebrovascular accident	2 (<1%)	1 (<1%)	1 (<1%)
		Ischaemic stroke	2 (<1%)	0	1 (<1%)
	Rare	Posterior reversible encephalopathy/ reversible posterior leukoencephalopathy syndrome†	not known	not known	not known
<b>Eye disorders</b>	Common	Vision blurred	19 (2%)	1 (<1%)	0
	Uncommon	Retinal detachment†	1 (<1%)	1 (<1%)	0
		Retinal tear†	1 (<1%)	1 (<1%)	0
		Eyelash discolouration	4 (<1%)	0	0
<b>Cardiac disorders</b>	Uncommon	Bradycardia	6 (<1%)	0	0
		Myocardial infarction	5 (<1%)	1 (<1%)	4 (<1%)
		Cardiac dysfunction <sup>f</sup>	4 (<1%)	1 (<1%)	0
		Myocardial ischaemia	3 (<1%)	1 (<1%)	0
<b>Vascular disorders</b>	Very common	Hypertension	473 (41%)	115 (10%)	1 (<1%)
	Common	Hot flush	16 (1%)	0	0
		Venous thromboembolic event <sup>g</sup>	13 (1%)	6 (<1%)	7 (<1%)
		Flushing	12 (1%)	0	0

System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
	Uncommon	Hypertensive crisis	6 (<1%)	0	2 (<1%)
		Haemorrhage	1 (<1%)	0	0
	Not known	Aneurysms <sup>†</sup> and artery dissections <sup>†</sup>	not known	not known	not known
<b>Respiratory, thoracic and mediastinal disorders</b>	Common	Epistaxis	50 (4%)	1 (<1%)	0
		Dysphonia	48 (4%)	0	0
		Dyspnoea	42 (4%)	8 (<1%)	1 (<1%)
		Haemoptysis	15 (1%)	1 (<1%)	0
	Uncommon	Rhinorrhoea	8 (<1%)	0	0
		Pulmonary haemorrhage	2 (<1%)	0	0
		Pneumothorax	1 (<1%)	0	0
	Rare	Interstitial lung disease/pneumonitis <sup>†</sup>	not known	not known	not known
<b>Gastrointestinal disorders</b>	Very common	Diarrhoea	614 (53% <sub>-</sub> )	65 (6%)	2 (<1%)
		Nausea	386 (34%)	14 (1%)	0
		Vomiting	225 (20%)	18 (2%)	1 (<1%)
		Abdominal pain <sup>a</sup>	139 (12%)	15 (1%)	0
	Common	Stomatitis	96 (8%)	4 (<1%)	0
		Dyspepsia	83 (7%)	2 (<1%)	0
		Flatulence	43 (4%)	0	0
		Abdominal distension	36 (3%)	2 (<1%)	0
		Mouth ulceration	28 (2%)	3 (<1%)	0
		Dry mouth	27 (2%)	0	0
	Uncommon	Pancreatitis	8 (<1%)	4 (<1%)	0
		Rectal haemorrhage	8 (<1%)	2 (<1%)	0
		Haematochezia	6 (<1%)	0	0
		Gastrointestinal haemorrhage	4 (<1%)	2 (<1%)	0
		Melaena	4 (<1%)	1 (<1%)	0
		Frequent bowel movements	3 (<1%)	0	0
		Anal haemorrhage	2 (<1%)	0	0
		Large intestine perforation	2 (<1%)	1 (<1%)	0
		Mouth haemorrhage	2 (<1%)	0	0
		Upper gastrointestinal haemorrhage	2 (<1%)	1 (<1%)	0
		Enterocutaneous fistula	1 (<1%)	0	0
		Haematemesis	1 (<1%)	0	0
		Haemorrhoidal haemorrhage	1 (<1%)	0	0
		Ileal perforation	1 (<1%)	0	1 (<1%)
		Oesophageal haemorrhage	1 (<1%)	0	0
		Retroperitoneal haemorrhage	1 (<1%)	0	0
<b>Hepatobiliary disorders</b>	Common	Hyperbilirubinaemia	38 (3%)	2 (<1%)	1 (<1%)
		Hepatic function abnormal	29 (3%)	13 (1%)	2 (<1%)
		Hepatotoxicity	18 (2%)	11 (<1%)	2 (<1%)
	Uncommon	Jaundice	3 (<1%)	1 (<1%)	0

System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
		Drug induced liver injury	2 (<1%)	2 (<1%)	0
		Hepatic failure <sup>†</sup>	1 (<1%)	0	1 (<1%)
Skin and subcutaneous disorders	Very common	Hair colour change	404 (35%)	1 (<1%)	0
		Palmar-plantar erythrodysesthesia syndrome	206 (18%)	39 (3%)	0
		Alopecia	130 (11%)	0	0
		Rash	129 (11%)	7 (<1%)	0
	Common	Skin hypopigmentation	52 (5%)	0	0
		Dry skin	50 (4%)	0	0
		Pruritus	29 (3%)	0	0
		Erythema	25 (2%)	0	0
		Skin depigmentation	20 (2%)	0	0
		Hyperhidrosis	17 (1%)	0	0
	Uncommon	Nail disorders	11 (<1%)	0	0
		Skin exfoliation	10 (<1%)	0	0
		Photosensitivity reaction	7 (<1%)	0	0
		Rash erythematous	6 (<1%)	0	0
		Skin disorder	5 (<1%)	0	0
		Rash macular	4 (<1%)	0	0
		Rash pruritic	3 (<1%)	0	0
		Rash vesicular	3 (<1%)	0	0
		Pruritus generalised	2 (<1%)	1 (<1%)	0
		Rash generalised	2 (<1%)	0	0
		Rash papular	2 (<1%)	0	0
		Plantar erythema	1 (<1%)	0	0
Musculoskeletal and connective tissue disorders	Common	Arthralgia	48 (4%)	8 (<1%)	0
		Myalgia	35 (3%)	2 (<1%)	0
		Muscle spasms	25 (2%)	0	0
	Uncommon	Musculoskeletal pain	9 (<1%)	1 (<1%)	0
Renal and urinary disorders	Very Common	Proteinuria	135 (12%)	32 (3%)	0
	Uncommon	Haemorrhage urinary tract	1 (<1%)	0	0
Reproductive system and breast disorders	Uncommon	Menorrhagia	3 (<1%)	0	0
		Vaginal haemorrhage	3 (<1%)	0	0
		Metrorrhagia	1 (<1%)	0	0
General disorders and administration site conditions	Very common	Fatigue	415 (36%)	65 (6%)	1 (<1%)
	Common	Mucosal inflammation	86 (7%)	5 (<1%)	0
		Asthenia	82 (7%)	20 (2%)	1 (<1%)
		Oedema <sup>b</sup>	72 (6%)	1 (<1%)	0
		Chest pain	18 (2%)	2 (<1%)	0
	Uncommon	Chills	4 (<1%)	0	0
		Mucous membrane disorder	1 (<1%)	0	0

System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
<b>Investigations</b>	Very common	Alanine aminotransferase increased	246 (21%)	84 (7%)	14 (1%)
		Aspartate aminotransferase increased	211 (18%)	51 (4%)	10 (<1%)
	Common	Weight decreased	96 (8%)	7 (<1%)	0
		Blood bilirubin increased	61 (5%)	6 (<1%)	1 (<1%)
		Blood creatinine increased	55 (5%)	3 (<1%)	0
		Lipase increased	51 (4%)	21 (2%)	7 (<1%)
		White blood cell count decreased <sup>d</sup>	51 (4%)	3 (<1%)	0
		Blood thyroid stimulating hormone increased	36 (3%)	0	0
		Amylase increased	35 (3%)	7 (<1%)	0
		Gamma-glutamyltransferase increased	31 (3%)	9 (<1%)	4 (<1%)
		Blood pressure increased	15 (1%)	2 (<1%)	0
		Blood urea increased	12 (1%)	1 (<1%)	0
		Liver function test abnormal	12 (1%)	6 (<1%)	1 (<1%)
	Uncommon	Hepatic enzyme increased	11 (<1%)	4 (<1%)	3 (<1%)
		Blood glucose decreased	7 (<1%)	0	1 (<1%)
		Electrocardiogram QT prolonged	7 (<1%)	2 (<1%)	0
		Transaminase increased	7 (<1%)	1 (<1%)	0
		Thyroid function test abnormal	3 (<1%)	0	0
		Blood pressure diastolic increased	2 (<1%)	0	0
		Blood pressure systolic increased	1 (<1%)	0	0

† Treatment-related adverse reaction reported during post-marketing period (spontaneous case reports and serious adverse reactions from all pazopanib clinical studies).

\* Treatment-related adverse reaction reported only during the post-marketing period. Frequency cannot be estimated from the available data.

The following terms have been combined:

<sup>a</sup> Abdominal pain, abdominal pain upper and abdominal pain lower

<sup>b</sup> Oedema, oedema peripheral, eye oedema, localised oedema and face oedema

<sup>c</sup> Dysgeusia, ageusia and hypogeusia

<sup>d</sup> White cell count decreased, neutrophil count decreased and leukocyte count decreased

<sup>e</sup> Decreased appetite and anorexia

<sup>f</sup> Cardiac dysfunction, left ventricular dysfunction, cardiac failure and restrictive cardiomyopathy

<sup>g</sup> Venous thromboembolic event, deep vein thrombosis, pulmonary embolism and thrombosis

Neutropenia, thrombocytopenia and palmar-plantar erythrodysesthesia syndrome were observed more frequently in patients of East Asian descent.

**Table 3 Treatment-related adverse reactions reported in STS studies (n=382) or during post-marketing period**

<b>System Organ Class</b>	<b>Frequency (all grades)</b>	<b>Adverse reactions</b>	<b>All grades n (%)</b>	<b>Grade 3 n (%)</b>	<b>Grade 4 n (%)</b>
<b>Infections and infestations</b>	Common	Gingival infection	4 (1%)	0	0
<b>Neoplasms benign, malignant and unspecified (incl. cysts and polyps)</b>	Very common	Tumour pain	121 (32%)	32 (8%)	0
<b>Blood and lymphatic system disorders<sup>f</sup></b>	Very common	Leukopenia	106 (44%)	3 (1%)	0
		Thrombocytopenia	86 (36%)	7 (3%)	2 (<1%)
		Neutropenia	79 (33%)	10 (4%)	0
	Uncommon	Thrombotic microangiopathy (including thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome)	1 (<1%)	1 (<1%)	0
<b>Endocrine disorders</b>	Common	Hypothyroidism	18 (5%)	0	0
<b>Metabolism and nutrition disorders</b>	Very common	Decreased appetite	108 (28%)	12 (3%)	0
		Hypoalbuminemia <sup>f</sup>	81 (34%)	2 (<1%)	0
	Common	Dehydration	4 (1%)	2 (1%)	0
	Uncommon	Hypomagnesaemia	1 (<1%)	0	0
	Not known	Tumour lysis syndrome*	not known	not known	not known
<b>Psychiatric disorders</b>	Common	Insomnia	5 (1%)	1 (<1%)	0
<b>Nervous system disorders</b>	Very common	Dysgeusia <sup>c</sup>	79 (21%)	0	0
		Headache	54 (14%)	2 (<1%)	0
	Common	Peripheral sensory neuropathy	30 (8%)	1 (<1%)	0
		Dizziness	15 (4%)	0	0
	Uncommon	Somnolence	3 (<1%)	0	0
		Paresthesia	1 (<1%)	0	0
		Cerebral infarction	1 (<1%)	0	1 (<1%)
<b>Eye disorders</b>	Common	Vision blurred	15 (4%)	0	0
<b>Cardiac disorders</b>	Common	Cardiac dysfunction <sup>g</sup>	21 (5%)	3 (<1%)	1 (<1%)
		Left ventricular dysfunction	13 (3%)	3 (<1%)	0
		Bradycardia	4 (1%)	0	0
	Uncommon	Myocardial infarction	1 (<1%)	0	0
<b>Vascular disorders</b>	Very common	Hypertension	152 (40%)	26 (7%)	0
	Common	Venous thromboembolic event <sup>d</sup>	13 (3%)	4 (1%)	5 (1%)
		Hot flush	12 (3%)	0	0
		Flushing	4 (1%)	0	0
	Uncommon	Haemorrhage	2 (<1%)	1 (<1%)	0

<b>System Organ Class</b>	<b>Frequency (all grades)</b>	<b>Adverse reactions</b>	<b>All grades n (%)</b>	<b>Grade 3 n (%)</b>	<b>Grade 4 n (%)</b>
	Not known	Aneurysms and artery dissections	not known	not known	not known
<b>Respiratory, thoracic and mediastinal disorders</b>	Common	Epistaxis	22 (6%)	0	0
		Dysphonia	20 (5%)	0	0
		Dyspnoea	14 (4%)	3 (<1%)	0
		Cough	12 (3%)	0	0
		Pneumothorax	7 (2%)	2 (<1%)	1 (<1%)
		Hiccups	4 (1%)	0	0
		Pulmonary haemorrhage	4 (1%)	1 (<1%)	0
	Uncommon	Oropharyngeal pain	3 (<1%)	0	0
		Bronchial haemorrhage	2 (<1%)	0	0
		Rhinorrhoea	1 (<1%)	0	0
		Haemoptysis	1 (<1%)	0	0
	Rare	Interstitial lung disease/pneumonitis†	not known	not known	not known
<b>Gastrointestinal disorders</b>	Very common	Diarrhoea	174 (46%)	17 (4%)	0
		Nausea	167 (44%)	8 (2%)	0
		Vomiting	96 (25%)	7 (2%)	0
		Abdominal pain <sup>a</sup>	55 (14%)	4 (1%)	0
		Stomatitis	41 (11%)	1 (<1%)	0
	Common	Abdominal distension	16 (4%)	2 (1%)	0
		Dry mouth	14 (4%)	0	0
		Dyspepsia	12 (3%)	0	0
		Mouth haemorrhage	5 (1%)	0	0
		Flatulence	5 (1%)	0	0
		Anal haemorrhage	4 (1%)	0	0
	Uncommon	Gastrointestinal haemorrhage	2 (<1%)	0	0
		Rectal haemorrhage	2 (<1%)	0	0
		Enterocutaneous fistula	1 (<1%)	1 (<1%)	0
		Gastric haemorrhage	1 (<1%)	0	0
		Melaena	2 (<1%)	0	0
		Oesophageal haemorrhage	1 (<1%)	0	1 (<1%)
		Peritonitis	1 (<1%)	0	0
		Retroperitoneal haemorrhage	1 (<1%)	0	0
		Upper gastrointestinal haemorrhage	1 (<1%)	1 (<1%)	0
		Ileal perforation	1 (<1%)	0	1 (<1%)
<b>Hepatobiliary disorders</b>	Uncommon	Hepatic function abnormal	2 (<1%)	0	1 (<1%)



System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
<b>Skin and subcutaneous disorders</b>	Very common	Hair colour change	93 (24%)	0	0
		Skin hypopigmentation	80 (21%)	0	0
		Exfoliative rash	52 (14%)	2 (<1%)	0
	Common	Alopecia	30 (8%)	0	0
		Skin disorder <sup>c</sup>	26 (7%)	4 (1%)	0
		Dry skin	21 (5%)	0	0
		Hyperhidrosis	18 (5%)	0	0
		Nail disorder	13 (3%)	0	0
		Pruritus	11 (3%)	0	0
		Erythema	4 (1%)	0	0
	Uncommon	Skin ulcer	3 (<1%)	1 (<1%)	0
		Rash	1 (<1%)	0	0
		Rash papular	1 (<1%)	0	0
		Photosensitivity reaction	1 (<1%)	0	0
		Palmar-plantar erythrodysesthesia syndrome	2 (<1%)	0	0
<b>Musculoskeletal and connective tissue disorders</b>	Common	Musculoskeletal pain	35 (9%)	2 (<1%)	0
		Myalgia	28 (7%)	2 (<1%)	0
		Muscle spasms	8 (2%)	0	0
	Uncommon	Arthralgia	2 (<1%)	0	0
<b>Renal and urinary disorders</b>	Uncommon	Proteinuria	2 (<1%)	0	0
<b>Reproductive system and breast disorder</b>	Uncommon	Vaginal haemorrhage	3 (<1%)	0	0
		Menorrhagia	1 (<1%)	0	0
<b>General disorders and administration site conditions</b>	Very common	Fatigue	178 (47%)	34 (9%)	1 (<1%)
	Common	Oedema <sup>b</sup>	18 (5%)	1 (<1%)	0
		Chest pain	12 (3%)	4 (1%)	0
		Chills	10 (3%)	0	0
	Uncommon	Mucosal inflammation <sup>e</sup>	1 (<1%)	0	0
		Asthenia	1 (<1%)	0	0
<b>Investigations<sup>h</sup></b>	Very common	Weight decreased	86 (23%)	5 (1%)	0
	Common	Ear, nose and throat examination abnormal <sup>e</sup>	29 (8%)	4 (1%)	0
		Alanine aminotransferase increased	8 (2%)	4 (1%)	2 (<1%)
		Blood cholesterol abnormal	6 (2%)	0	0
		Aspartate aminotransferase increased	5 (1%)	2 (<1%)	2 (<1%)
		Gamma glutamyltransferase increased	4 (1%)	0	3 (<1%)
	Uncommon	Blood bilirubin increased	2 (<1%)	0	0
		Aspartate aminotransferase	2 (<1%)	0	2 (<1%)
		Alanine aminotransferase	1 (<1%)	0	1 (<1%)
		Platelet count decreased	1 (<1%)	0	1 (<1%)

System Organ Class	Frequency (all grades)	Adverse reactions	All grades n (%)	Grade 3 n (%)	Grade 4 n (%)
		Electrocardiogram QT prolonged	2 (<1%)	1 (<1%)	0

† Treatment-related adverse reaction reported during post-marketing period (spontaneous case reports and serious adverse reactions from all pazopanib clinical studies).

\* Treatment-related adverse reaction reported only during the post-marketing period. Frequency cannot be estimated from the available data.

The following terms have been combined:

<sup>a</sup> Abdominal pain, abdominal pain upper and gastrointestinal pain

<sup>b</sup> Oedema, oedema peripheral and eyelid oedema

<sup>c</sup> The majority of these cases were Palmar-plantar erythrodysesthesia syndrome

<sup>d</sup> Venous thromboembolic events – includes Deep vein thrombosis, Pulmonary embolism and Thrombosis terms

<sup>e</sup> The majority of these cases describe mucositis

<sup>f</sup> Frequency is based on laboratory value tables from VEG110727 (N=240). These were reported as adverse events less frequently by investigators than as indicated by laboratory value tables.

<sup>g</sup> Cardiac dysfunction events – includes Left ventricular dysfunction, Cardiac failure and Restrictive cardiomyopathy

<sup>h</sup> Frequency is based on adverse events reported by investigators. Laboratory abnormalities were reported as adverse events less frequently by investigators than as indicated by laboratory value tables.

Neutropenia, thrombocytopenia and palmar-plantar erythrodysesthesia syndrome were observed more frequently in patients of East Asian descent.

## 4.9 Overdose

Pazopanib doses up to 2000 mg have been evaluated in clinical studies. Grade 3 fatigue (dose-limiting toxicity) and Grade 3 hypertension were each observed in 1 of 3 patients dosed at 2000 mg and 1000 mg daily, respectively.

There is no specific antidote for overdose with pazopanib and treatment of overdose should consist of general supportive measures.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, other antineoplastic agents, protein kinase inhibitors, ATC code: L01EX03

#### Mechanism of action

Pazopanib is an orally administered, potent multi-target tyrosine kinase inhibitor (TKI) of vascular endothelial growth factor receptors (VEGFR) -1, -2, and -3, platelet-derived growth factor (PDGFR) - $\alpha$  and - $\beta$ , and stem cell factor receptor (c-KIT), with IC<sub>50</sub> values of 10, 30, 47, 71, 84 and 74 nM, respectively. In preclinical experiments, pazopanib dose-dependently inhibited ligand-induced auto-phosphorylation of VEGFR-2, c-Kit and PDGFR- receptors in cells. *In vivo*, pazopanib inhibited VEGF-induced VEGFR-2 phosphorylation in mouse lungs, angiogenesis in various animal models, and the growth of multiple human tumour xenografts in mice.

#### Pharmacogenomics

In a pharmacogenetic meta-analysis of data from 31 clinical studies of pazopanib administered either as monotherapy or in combination with other agents, ALAT >5 x ULN (NCI CTC Grade 3) occurred in 19% of HLA-B\*57:01 allele carriers and in 10% of non-carriers. In this dataset, 133/2235 (6%) of the patients carried the HLA-B\*57:01 allele (see section 4.4).

## Clinical studies

### *Renal cell carcinoma (RCC)*

The safety and efficacy of pazopanib in RCC were evaluated in a randomised, double-blind, placebo-controlled multicentre study. Patients (N=435) with locally advanced and/or metastatic RCC were randomised to receive pazopanib 800 mg once daily or placebo. The primary objective of the study was to evaluate and compare the two treatment arms for progression-free survival (PFS) and the principle secondary endpoint was overall survival (OS). The other objectives were to evaluate the overall response rate and duration of response.

From the total of 435 patients in this study, 233 patients were treatment-naïve and 202 were second-line patients who had received one prior IL-2 or INF-based therapy. The performance status (ECOG) was similar between the pazopanib and placebo groups (ECOG 0: 42% vs. 41%, ECOG 1: 58% vs. 59%). The majority of patients had either favourable (39%) or intermediate (54%), MSKCC (Memorial Sloan Kettering Cancer Centre) / Motzer prognostic factors. All patients had clear cell histology or predominantly clear cell histology. Approximately half of all patients had 3 or more organs involved in their disease and most patients had the lung (74%), and/or lymph nodes (54%) as a metastatic location for disease at baseline.

A similar proportion of patients in each arm were treatment-naïve and cytokine pre-treated (53% and 47% in pazopanib arm, 54% and 46% in placebo arm). In the cytokine pre-treated subgroup, the majority (75%) had received interferon-based treatment.


Similar proportions of patients in each arm had prior nephrectomy (89% and 88% in the pazopanib and placebo arms, respectively) and/or prior radiotherapy (22% and 15% in the pazopanib and placebo arms, respectively).

The primary analysis of the primary endpoint PFS is based on disease assessment by independent radiological review in the entire study population (treatment-naïve and cytokine pre-treated).

**Table 4 Overall efficacy results in RCC by independent assessment (VEG105192)**


Endpoints/Study	Pazopanib	Placebo	HR (95% CI)	P value (one-sided)
PFS				
Overall* ITT	N=290	N=145		
Median (months)	9.2	4.2	0.46 (0.34, 0.62)	<0.0000001
Response rate	N =290	N =145		
% (95% CI)	30 (25.1,35.6)	3 (0.5, 6.4)	—	<0.001
HR=hazard ratio; ITT=intent to treat; PFS=progression-free survival. * - treatment-naïve and cytokine pre-treated populations				

**Figure 1** Kaplan-Meier curve for progression-free survival by independent assessment for the overall population (treatment-naïve and cytokine pre-treated populations) (VEG105192)




x axis; Months, y axis; Proportion Progression Free, Pazopanib — (N=290) Median 9.2 months; Placebo - - - (N=145) Median 4.2 months; Hazard Ratio = 0.46, 95% CI (0.34, 0.62), P < 0.0000001

**Figure 2** Kaplan-Meier curve for progression-free survival by independent assessment for the treatment-naïve population (VEG105192)



x axis; Months, y axis; Proportion Progression Free, Pazopanib — (N = 155) Median 11.1 months; Placebo - - - (N = 78) Median 2.8 months; Hazard Ratio = 0.40, 95% CI (0.27, 0.60), P < 0.0000001

**Figure 3** Kaplan-Meier Curve for progression-free survival by independent assessment for the cytokine pre-treated population (VEG105192)



x axis; Months, y axis; Proportion Progression Free, Pazopanib — (N=135) Median 7.4 months; Placebo ----- (N=67) Median 4.2 months; Hazard Ratio =0.54, 95% CI (0.35, 0.84), P <0.001

For patients who responded to treatment, the median time to response was 11.9 weeks and the median duration of response was 58.7 weeks as per independent review (VEG105192).

The median overall survival (OS) data at the protocol-specified final survival analysis were 22.9 months and 20.5 months [HR=0.91 (95% CI: 0.71, 1.16; p=0.224)] for patients randomised to the pazopanib and placebo arms, respectively. The OS results are subject to potential bias as 54% of patients in the placebo arm also received pazopanib in the extension part of this study following disease progression. Sixty-six per cent of placebo patients received post-study therapy compared to 30% of pazopanib patients.

No statistical differences were observed between treatment groups for Global Quality of Life using EORTC QLQ-C30 and EuroQoL EQ-5D.

In a Phase II study of 225 patients with locally recurrent or metastatic clear cell renal cell carcinoma, objective response rate was 35% and median duration of response was 68 weeks, as per independent review. Median PFS was 11.9 months.

The safety, efficacy and quality of life of pazopanib versus sunitinib was evaluated in a randomised, open-label, parallel group Phase III non-inferiority study (VEG108844).


In VEG108844, patients (N=1110) with locally advanced and/or metastatic RCC who had not received prior systemic therapy, were randomised to receive either pazopanib 800 mg once daily continuously or sunitinib 50 mg once daily in 6-week cycles of dosing with 4 weeks on treatment followed by 2 weeks without treatment.

The primary objective of this study was to evaluate and compare PFS in patients treated with pazopanib to those treated with sunitinib. Demographic characteristics were similar between the treatment arms. Disease characteristics at initial diagnosis and at screening were balanced between the treatment arms with the majority of patients having clear cell histology and Stage IV disease.

VEG108844 achieved its primary endpoint of PFS and demonstrated that pazopanib was non-inferior to sunitinib, as the upper bound of the 95% CI for the hazard ratio was less than the protocol-specified non-inferiority margin of 1.25. Overall efficacy results are summarised in Table 5.

**Table 5 Overall efficacy results (VEG108844)**

Endpoint	Pazopanib N=557	Sunitinib N=553	HR (95% CI)
<b>PFS</b>			
Overall			
Median (months)	8.4	9.5	1.047
(95% CI)	(8.3, 10.9)	(8.3, 11.0)	(0.898, 1.220)
<b>Overall Survival</b>			
Median (months)	28.3	29.1	0.915 <sup>a</sup>
(95% CI)	(26.0, 35.5)	(25.4, 33.1)	(0.786, 1.065)
HR=hazard ratio; PFS=progression-free survival; <sup>a</sup> P value=0.245 (2-sided)			

**Figure 4 Kaplan-Meier Curve for progression-free survival by independent assessment for the overall population (VEG108844)**

Subgroup analyses of PFS were performed for 20 demographic and prognostic factors. The 95% confidence intervals for all subgroups include a hazard ratio of 1. In the three smallest of these 20 subgroups, the point estimate of the hazard ratio exceeded 1.25; i.e. in subjects with no prior nephrectomy (n=186, HR=1.403, 95% CI (0.955, 2.061)), baseline LDH >1.5 x ULN (n=68, HR=1.72, 95% CI (0.943, 3.139)), and MSKCC: poor risk (n=119, HR=1.472, 95% CI (0.937, 2.313)).

#### *Soft-tissue sarcoma (STS)*

The efficacy and safety of pazopanib in STS were evaluated in a pivotal Phase III randomised, double-blind, placebo-controlled multicentre study (VEG110727). A total of 369 patients with advanced STS were randomised to receive pazopanib 800 mg once daily or placebo. Importantly, only patients with selective histological subtypes of STS were allowed to participate to the study, therefore efficacy and safety of pazopanib can only be considered established for those subgroups of STS and treatment with pazopanib should be restricted to such STS subtypes.

The following tumour types were eligible:

Fibroblastic (adult fibrosarcoma, myxofibrosarcoma, sclerosing epithelioid fibrosarcoma, malignant solitary fibrous tumours), so-called fibrohistiocytic (pleomorphic malignant fibrous histiocytoma [MFH], giant cell MFH, inflammatory MFH), leiomyosarcoma, malignant glomus tumours, skeletal

muscles (pleomorphic and alveolar rhabdomyosarcoma), vascular (epithelioid hemangioendothelioma, angiosarcoma), uncertain differentiation (synovial, epithelioid, alveolar soft part, clear cell, desmoplastic small round cell, extra-renal rhabdoid, malignant mesenchymoma, PEComa, intimal sarcoma), malignant peripheral nerve sheath tumours, undifferentiated soft tissue sarcomas not otherwise specified (NOS) and other types of sarcoma (not listed as ineligible).

The following tumour types were not eligible:

Adipocytic sarcoma (all subtypes), all rhabdomyosarcoma that were not alveolar or pleomorphic, chondrosarcoma, osteosarcoma, Ewing tumours/primitive neuroectodermal tumours (PNET), GIST, dermatofibromatosis sarcoma protuberans, inflammatory myofibroblastic sarcoma, malignant mesothelioma and mixed mesodermal tumours of the uterus.

Of note, patients with adipocytic sarcoma were excluded from the pivotal Phase III study as in a preliminary Phase II study (VEG20002) activity (PFS at week 12) observed with pazopanib in adipocytic did not meet the prerequisite rate to allow further clinical testing.

Other key eligibility criteria of the VEG110727 study were: histological evidence of high or intermediate grade malignant STS and disease progression within 6 months of therapy for metastatic disease, or recurrence within 12 months of (neo) -/adjuvant therapy.

Ninety-eight percent (98%) of subjects received prior doxorubicin, 70% prior ifosfamide, and 65% of subjects had received at least three or more chemotherapeutic agents prior to study enrolment.

Patients were stratified by the factors of WHO performance status (WHO PS) (0 or 1) at baseline and the number of lines of prior systemic therapy for advanced disease (0 or 1 vs. 2+). In each treatment group, there was a slightly greater percentage of subjects in the 2+ lines of prior systemic therapy for advanced disease (58% and 55%, respectively, for placebo and pazopanib treatment arms) compared with 0 or 1 lines of prior systemic therapy (42% and 45%, respectively, for placebo and pazopanib treatment arms). The median duration of follow-up of subjects (defined as date of randomisation to date of last contact or death) was similar for both treatment arms (9.36 months for placebo [range 0.69 to 23.0 months] and 10.04 months for pazopanib [range 0.2 to 24.3 months]).

The primary objective of the study was progression-free survival (PFS assessed by independent radiological review); the secondary endpoints included overall survival (OS), overall response rate and duration of response.


**Table 6 Overall efficacy results in STS by independent assessment (VEG110727)**

<b>Endpoints / study population</b>	<b>Pazopanib</b>	<b>Placebo</b>	<b>HR (95% CI)</b>	<b>P value (two-sided)</b>
<b>PFS</b>				
Overall ITT	N=246	N=123		
Median (weeks)	20.0	7.0	0.35 (0.26, 0.48)	<0.001
Leiomyosarcoma	N=109	N=49		
Median (weeks)	20.1	8.1	0.37 (0.23, 0.60)	<0.001
Synovial sarcoma subgroups	N=25	N=13		
Median (weeks)	17.9	4.1	0.43 (0.19, 0.98)	0.005
'Other STS' subgroups	N=112	N=61		
Median (weeks)	20.1	4.3	0.39 (0.25, 0.60)	<0.001

Endpoints / study population	Pazopanib	Placebo	HR (95% CI)	P value (two-sided)
<b>OS</b>				
Overall ITT Median (months)	N=246 12.6	N=123 10.7	0.87 (0.67, 1.12)	0.256
Leiomyosarcoma* Median (months)	N=109 16.7	N=49 14.1	0.84 (0.56, 1.26)	0.363
Synovial sarcoma subgroups* Median (months)	N=25 8.7	N=13 21.6	1.62 (0.79, 3.33)	0.115
“Other STS” subgroups* Median (months)	N=112 10.3	N=61 9.5	0.84 (0.59, 1.21)	0.325
<b>Response rate (CR+PR)</b> % (95% CI)	4 (2.3, 7.9)	0 (0.0, 3.0)		
Duration of response Median (weeks) (95% CI)	38.9 (16.7, 40.0)			
HR=hazard ratio; ITT=intent to treat; PFS=progression-free survival; CR=complete response; PR=partial response. OS=overall survival * Overall survival for the respective STS histological subgroups (leiomyosarcoma, synovial sarcoma and “Other” STS) should be interpreted with caution due to the small number of subjects and wide confidence intervals				

A similar improvement in PFS based on investigator assessments was observed in the pazopanib arm compared with the placebo arm (in the overall ITT population HR: 0.39; 95% CI, 0.30 to 0.52,  $p < 0.001$ ).

**Figure 5 Kaplan-Meier Curve for Progression-Free Survival in STS by Independent Assessment for the Overall Population (VEG110727)**



No significant difference in OS was observed between the two treatment arms at the final OS analysis performed after 76% (280/369) of the events had occurred (HR 0.87, 95% CI 0.67, 1.12  $p=0.256$ ).



## 5.2 Pharmacokinetic properties

### Absorption

Upon oral administration of a single pazopanib 800 mg dose to patients with solid tumours, maximum plasma concentration ( $C_{\max}$ ) of approximately  $19 \pm 13$   $\mu\text{g/ml}$  was obtained after median 3.5 hours (range 1.0-11.9 hours) and an  $\text{AUC}_{0-\infty}$  of approximately  $650 \pm 500$   $\mu\text{g.h/ml}$  was obtained. Daily dosing results in 1.23- to 4-fold increase in  $\text{AUC}_{0-T}$ .

There was no consistent increase in AUC or  $C_{\max}$  at pazopanib doses above 800 mg.

Systemic exposure to pazopanib is increased when administered with food. Administration of pazopanib with a high-fat or low-fat meal results in an approximately 2-fold increase in AUC and  $C_{\max}$ . Therefore, pazopanib should be administered at least two hours after food or at least one hour before food (see section 4.2).

Administration of a pazopanib 400 mg crushed tablet increased  $\text{AUC}_{(0-72)}$  by 46% and  $C_{\max}$  by approximately 2 fold and decreased  $t_{\max}$  by approximately 2 hours compared to administration of the whole tablet. These results indicate that the bioavailability and the rate of pazopanib oral absorption are increased after administration of the crushed tablet relative to administration of the whole tablet (see section 4.2).

### Distribution

Binding of pazopanib to human plasma protein *in vivo* was greater than 99% with no concentration dependence over the range of 10-100  $\mu\text{g/ml}$ . *In vitro* studies suggest that pazopanib is a substrate for P-gp and BCRP.

### Biotransformation

Results from *in vitro* studies demonstrated that metabolism of pazopanib is mediated primarily by CYP3A4, with minor contributions from CYP1A2 and CYP2C8. The four principle pazopanib metabolites account for only 6% of the exposure in plasma. One of these metabolites inhibits the proliferation of VEGF-stimulated human umbilical vein endothelial cells with a similar potency to that of pazopanib, the others are 10- to 20-fold less active. Therefore, activity of pazopanib is mainly dependent on parent pazopanib exposure.

### Elimination

Pazopanib is eliminated slowly with a mean half-life of 30.9 hours after administration of the recommended dose of 800 mg. Elimination is primarily via faeces with renal elimination accounting for <4% of the administered dose.

### Special populations

#### *Renal impairment*

Results indicate that less than 4% of an orally administered pazopanib dose is excreted in the urine as pazopanib and metabolites. Results from population pharmacokinetic modelling (data from subjects with baseline CLCR values ranging from 30.8 ml/min to 150 ml/min) indicated that renal impairment is unlikely to have clinically relevant effect on pazopanib pharmacokinetics. No dose adjustment is required in patients with creatinine clearance above 30 ml/min. (see section 4.2).

#### *Hepatic impairment*

The median steady-state pazopanib  $C_{\max}$  and  $\text{AUC}_{(0-24)}$  in patients with mild hepatic impairment (defined as either normal bilirubin and any degree of ALT elevations or as an elevation of bilirubin up to 1.5 X ULN regardless of the ALT value) after a once daily dose of 800 mg/day (30.9 microgram/mL, range 12.5 to 47.3 and 841.8 microgram.hr/mL, range 600.4 to 1,078) are similar to the median in patients with no hepatic impairment (49.4 microgram/mL, range 17.1 to 85.7 and 888.2 microgram.hr/mL, range 345.5 to 1,482) (see section 4.2).

The maximally tolerated pazopanib dose (MTD) in patients with moderate hepatic impairment (defined as an elevation of bilirubin > 1.5 X to 3 X ULN regardless of the ALT values) was 200 mg

once daily. The median steady-state values of C<sub>max</sub> (22.4 microgram/mL, range 6.4 to 32.9) and AUC(0-24) (350.0 microgram.hr/mL, range 131.8 to 487.7) after administration of 200 mg pazopanib once daily in subjects with moderate hepatic impairment were approximately 45 % and 39 %, respectively, that of the corresponding median values after administration of 800 mg once daily in subjects with normal hepatic function (see section 4.2).

There are insufficient data in patients with severe hepatic impairment (total bilirubin > 3 X ULN regardless of the ALT value); therefore, use of pazopanib is not recommended in these patients.

### 5.3 Preclinical safety data

The preclinical safety profile of pazopanib was assessed in mice, rats, rabbits and monkeys. In repeat dose studies in rodents, effects in a variety of tissues (bone, teeth, nail beds, reproductive organs, haematological tissues, kidney and pancreas) appear related to the pharmacology of VEGFR inhibition and/or disruption of VEGF signalling pathways, with most effects occurring at plasma exposure levels below those observed in the clinic. Other observed effects include body weight loss, diarrhoea and/or morbidity that were either secondary to local gastrointestinal effects caused by high local mucosal medicinal product exposure (monkeys) or pharmacological effects (rodents). Proliferative hepatic lesions (eosinophilic foci and adenoma) were seen in female mice at exposures 2.5 times human exposure based on AUC.

In juvenile toxicity studies, when pre-weaning rats were dosed from day 9 post partum through to day 14 post partum, pazopanib caused mortalities and abnormal organ growth/maturation in kidney, lung, liver and heart, at a dose approximately 0.1 times the clinical exposure based on AUC in adult humans. When post-weaning rats were dosed from day 21 post partum to day 62 post partum, toxicological findings were similar to adult rats at comparable exposures. Human paediatric patients are at increased risk for bone and teeth effects as compared to adults, as these changes, including inhibition of growth (shortened limbs), fragile bones and remodelling of teeth, were present in juvenile rats at  $\geq 10$  mg/kg/day (equal to approximately 0.1-0.2 times the clinical exposure based on AUC in adult humans) (see section 4.4).

#### Reproductive, fertility and teratogenic effects

Pazopanib has been shown to be embryotoxic and teratogenic when administered to rats and rabbits at exposures more than 300-fold lower than the human exposure (based on AUC). Effects included reduced female fertility, increased pre- and post-implantation loss, early resorptions, embryo lethality, decreased foetal body weight and cardiovascular malformation. Decreased corpora lutea, increased cysts and ovarian atrophy have also been noted in rodents. In a rat male fertility study, there was no effect on mating or fertility, but decreased testicular and epididymal weights were noted with reductions in sperm production rates, sperm motility, and epididymal and testicular sperm concentrations observed at exposures 0.3 times human exposure based on AUC.

#### Genotoxicity

Pazopanib did not cause genetic damage when tested in genotoxicity assays (Ames assay, human peripheral lymphocyte chromosome aberration assay and rat *in vivo* micronucleus). A synthetic intermediate in manufacture of pazopanib, which is also present in the final drug substance in low amounts, was not mutagenic in the Ames assay but genotoxic in the mouse lymphoma assay and *in vivo* mouse micronucleus assay.

#### Carcinogenicity

In two-year carcinogenicity studies with pazopanib, there were increased numbers of liver adenomas noted in mice and duodenal adenocarcinomas noted in rats. Based on the rodent-specific pathogenesis and mechanism for these findings, they are not considered to represent an increased carcinogenic risk for patients taking pazopanib.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Cellulose, microcrystalline (E460)

Povidone (K-30)

Sodium starch glycolate (Type A)

Magnesium stearate (E470b)

#### Tablet coating

Titanium dioxide (E171)

Hypromellose

Macrogol 400

Polysorbate 80 (E433)

Iron oxide red (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

Please refer to outer carton.

### **6.4 Special precautions for storage**

Please refer to outer carton.

### **6.5 Nature and contents of container**

HDPE containers with child-resistant PP caps (with liner), or child-resistant HDPE/PP caps (with liner), and a desiccant cylinder, containing 30, 90 or 120 tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements.

## **7. PRODUCT OWNER**

LOTUS INTERNATIONAL PTE. LTD.

80 Robinson Road

#02-00

Singapore 068898

## **8. DATE OF REVISION OF THE TEXT**

10/2022