

Vitalipid™ N Adult

Interactions
Vitalipid N Adult contains vitamin K₁ which may interact with anticoagulants of the coumadin type.

Storage

Store below 25 °C.
Protect from light.
Do not freeze.

Further information
The manufacturer can be consulted for further information on complete and balanced intravenous nutrition regimens.

Package quantities
Ampoules 10x10 ml

Manufactured by

Fresenius Kabi AB, Uppsala, Sweden
Sept 2008

Indications
Vitalipid N Adult is indicated as a supplement in complete intravenous nutrition to meet the daily requirements of the fat-soluble vitamins A, D₂, E and K₁.

Contraindications

Known hypersensitivity to egg-, soya- or peanut protein or to any of the active substances or excipients.

Dosage

Must be diluted before use. Adults and children age 11 years and above.

One ampoule (10 ml) of Vitalipid N Adult is added to 500 ml Intralipid 10% or 20%. After mixing by gentle agitation the emulsion is infused as described for Intralipid. The daily maintenance dosage of the vitamins A, D₂, E and K₁ are thereby supplied.

Vitalipid N Adult should be added aseptically within one hour before the start of the infusion and should be used within 24 hours.

Vitalipid N Adult can be used to reconstitute Soluvit™ N. The contents of one vial of Soluvit N (the daily main-tenance dosages of water-soluble vitamins) are dissolved by the aseptic addition of 10 ml of Vitalipid N Adult and added to Intralipid.

Properties

The daily maintenance dosage of the vitamins A, D₂, E and K₁ are supplied during intravenous nutrition when 10 ml of Vitalipid N Adult are added to 500 ml Intralipid 10% or 20%.

Warnings and precautions

This medicinal product contains soya-bean oil and egg phospholipids which may rarely cause allergic reactions. Cross allergic reaction has been observed between soya-bean and peanut. Vitalipid N Adult must not be given undiluted.



331214

