

Berodual®



Composition

BERODUAL Solution for Inhalation

1 mL (= 20 drops) solution for inhalation contains:

(8*r*)-3*α*-hydroxy-8-isopropyl-1*α*H,5*α*H-tropanium bromide (±)-tropate monohydrate
(= ipratropium bromide) corresponding to 250 mcg ipratropium bromide anhydrous 261mcg

1-(3,5-dihydroxy-phenyl)-2-[[1-(4-hydroxy-benzyl)-ethyl]-amino]-ethanol
hydrobromide (= fenoterol hydrobromide) 500mcg

Excipients: benzalkonium chloride, disodium edetate dihydrate, sodium chloride, hydrochloric acid, water purified

BERODUAL N Metered Dose Inhaler

1 metered dose (puff) contains:

(8*r*)-3*α*-hydroxy-8-isopropyl-1*α*H,5*α*H-tropanium bromide (±)-tropate monohydrate 21mcg

(= ipratropium bromide) corresponding to 20 mcg ipratropium bromide anhydrous

1-(3,5-dihydroxy-phenyl)-2-[[1-(4-hydroxy-benzyl)-ethyl]-amino]-ethanol hydrobromide 50mcg
(= fenoterol hydrobromide)

Propellant: 1,1,1,2-Tetrafluoroethane (HFA 134a)

Other excipients: citric acid anhydrous, water purified, ethanol absolute, nitrogen (inert gas)

Description

BERODUAL Solution for Inhalation

Clear, colourless or almost colourless liquid, free from suspended particles

BERODUAL N Metered Dose Inhaler

Clear, colourless liquid, free from suspended particles

Pharmacological Properties

Pharmacotherapeutic group: Adrenergics in combination with anticholinergics for obstructive airway diseases

ATC code: R03AL01

Mode of Action

BERODUAL contains two active bronchodilating ingredients: ipratropium bromide, exhibiting an anticholinergic effect and fenoterol hydrobromide a beta-adrenergic agent.

Ipratropium bromide is a quaternary ammonium compound with anticholinergic (parasympatholytic) properties. In non-clinical studies, it inhibits vagally mediated reflexes by antagonising the action of acetylcholine, the transmitter agent released from the vagus nerve. Anticholinergics prevent the increase in intracellular concentration of Ca⁺⁺ which is caused by interaction of acetylcholine with the muscarinic receptor on bronchial smooth muscle. Ca⁺⁺ release is mediated by the second messenger system consisting of IP₃ (inositol triphosphate) and DAG (diacylglycerol).

The bronchodilatation following inhalation of ipratropium bromide is primarily a local, site-specific effect, not a systemic one.

Non-clinical and clinical evidence suggest no deleterious effect of ipratropium bromide on airway mucous secretion, mucociliary clearance or gas exchange.

Fenoterol hydrobromide is a direct acting sympathomimetic agent, selectively stimulating beta₂-receptors in the therapeutic dose range. The stimulation of beta₁-receptors comes into effect at a higher dose range. Occupation of beta₂-receptors activates adenylyl cyclase via a stimulatory G_s-protein.

The increase in cyclic AMP activates protein kinase A which then phosphorylates target proteins in smooth muscle cells. This in turn leads to the phosphorylation of myosin light chain kinase, inhibition of phosphoinositide hydrolysis, and the opening of large-conductance calcium-activated potassium channels.

Fenoterol hydrobromide relaxes bronchial and vascular smooth muscle and protects against bronchoconstricting stimuli such as histamine, methacholine, cold air, and allergen (early response). After acute administration the release of bronchoconstricting and pro-inflammatory mediators from mast cells is inhibited. Further, an increase in mucociliary clearance has been demonstrated after administration of doses of fenoterol (0.6 mg).

Higher plasma concentrations, which are more frequently achieved with oral, or even more so, with intravenous administration inhibit uterine motility. Also at higher doses, metabolic effects are observed: Lipolysis, glycogenolysis, hyperglycaemia and hypokalaemia, the latter caused by increased K⁺-uptake primarily into skeletal muscle. Beta-adrenergic effects on the heart such as increase in heart rate and contractility are caused by the vascular effects of fenoterol, cardiac beta₂-receptor stimulation, and at supratherapeutic doses, by beta₁-receptor stimulation. As with other beta-adrenergic agents, QTc prolongations have been reported. For fenoterol metered dose inhalers these were discrete and observed at doses higher than recommended. However, systemic exposure after administration with nebulisers (UDVs, solution for inhalation) might be higher than with recommended MDI doses. The clinical significance has not been established. Tremor is a more frequently observed effect of beta-agonists. Unlike the effects on the bronchial smooth muscle, the systemic effects on skeletal muscle of β-agonists are subject to the development of tolerance.

Concurrent use of these two active ingredients dilates the bronchi by affecting different pharmacological sites of action. The two active substances thus complement each other in their spasmolytic action on the bronchial muscles and allow a broad therapeutic use in the field of bronchopulmonary disorders associated with constriction of the respiratory tract. The complementary action is such that only a very low proportion of the β-adrenergic component is needed to obtain the desired effect, facilitating individual dosage suited to each patient with a minimum of adverse reactions.

Clinical efficacy and safety

In patients with asthma and COPD, better efficacy compared to its components ipratropium or fenoterol was demonstrated. Two studies (one with asthma patients, one with COPD patients) have shown that BERODUAL is as efficacious as double the dose of fenoterol administered without ipratropium but was better tolerated in cumulative dose response studies.

In acute bronchoconstriction BERODUAL is effective shortly after administration and is therefore also suitable for treating acute episodes of bronchospasm.

Pharmacokinetics

The therapeutic effect of the combination ipratropium bromide and fenoterol hydrobromide is produced by a local action in the airway. The pharmacodynamics of the bronchodilation are therefore not related to the pharmacokinetics of the active constituents of the preparation.

Following inhalation 10 to 30% of a dose is generally deposited in lungs, depending on the formulation, inhalation technique and device, while the remainder of the delivered dose is deposited in the mouthpiece, mouth and the upper part of the respiratory tract (oropharynx). A similar amount of the dose is deposited in the respiratory tract following inhalation by metered aerosol with HFA 134a propellant. In particular after inhalation of the aqueous solution via the RESPIMAT inhaler, a more than 2-fold higher lung deposition is experimentally observed as compared to the metered aerosol inhaler. The oropharyngeal deposition is correspondingly decreased and is significantly lower for the RESPIMAT inhaler as compared to the metered aerosol inhaler. The portion of the dose deposited in the lungs reaches the circulation rapidly (within minutes). The amount of the active substance deposited in the oropharynx is slowly swallowed and passes the gastrointestinal tract. Therefore the systemic exposure is a function of both oral and lung bioavailability.

There is no evidence that the pharmacokinetics of both ingredients in the combination differ from those of the mono-substance.

Fenoterol hydrobromide

Absorption

The absolute bioavailability following oral administration is low (approx. 1.5%).

The absolute bioavailability of fenoterol following inhalation is 18.7%. Absorption from the lung follows a biphasic course. 30% of the fenoterol hydrobromide dose is rapidly absorbed with a half-life of 11 minutes and 70% is slowly absorbed with a half-life of 120 minutes.

Distribution

Fenoterol distributes widely throughout the body. About 40 % of the drug are bound to plasma proteins. In this 3-compartment model the apparent volume of distribution of fenoterol at steady state (V_{dss}) is approximately 189 L (≈ 2.7 L/kg).

Non-clinical studies with rats revealed that fenoterol and its metabolites do not cross the blood-brain barrier.

Biotransformation

Fenoterol undergoes extensive metabolism by conjugation to glucuronides and sulphates in humans. Following oral administration, fenoterol is metabolised predominantly by sulphation. This metabolic inactivation of the parent compound starts already in the intestinal wall.

Elimination

After inhalation via BERODUAL metered dose inhaler approximately 1% of an inhaled dose is excreted as free fenoterol in the 24-hour urine. Based on these data, the total systemic bioavailability of inhaled doses of fenoterol hydrobromide is estimated at 7%. Fenoterol has a total clearance of 1.8 L/min and a renal clearance of 0.27 L/min.

Kinetic parameters describing the disposition of fenoterol were calculated from plasma concentrations after i.v. administration. Following intravenous administration, plasma concentration-time profiles can be described by a 3-compartment model, whereby the terminal half-life is approximately 3 hours.

Following oral administration, total radioactivity excreted in urine was approximately 39% of dose and total radioactivity excreted in faeces was 40.2% of dose within 48 hours.

Ipratropium bromide

Absorption

Cumulative renal excretion (0-24 hrs) of ipratropium (parent compound) is below 1% of an oral dose and approximately 3 to 13% of an inhaled dose via BERODUAL metered dose inhaler. Based on these data, the total systemic bioavailability of oral and inhaled doses of ipratropium bromide is estimated at 2% and 7 to 28% respectively. Taking this into account, swallowed dose portions of ipratropium bromide do not relevantly contribute to systemic exposure.

Distribution

Kinetic parameters describing the disposition of ipratropium were calculated from plasma concentrations after i.v. administration. A rapid biphasic decline in plasma concentrations is observed. The apparent volume of distribution at steady-state (V_{dss}) is approximately 176 L (≈ 2.4 L/kg). The drug is minimally (less than 20%) bound to plasma proteins. It is not known if the placental barrier is crossed. Non-clinical studies with rats and dogs, revealed that the quaternary amine ipratropium does not cross the blood-brain barrier.

Binding of the main urinary metabolites to the muscarinic receptor is negligible and the metabolites have to be regarded as ineffective.

Biotransformation

After intravenous administration approximately 60% of a dose is metabolised, the major portion probably in the liver by oxidation.

Elimination

The half-life of the terminal elimination phase is approximately 1.6 hours. Ipratropium has a total clearance of 2.3 L/min and a renal clearance of 0.9 L/min. In an excretion balance study cumulative renal excretion (6 days) of drug-related radioactivity (including parent compound and all metabolites) accounted for 9.3% after oral administration and 3.2% after inhalation. Total radioactivity excreted via the faeces was 88.5% following oral dosing and 69.4% after inhalation.

Indications

BERODUAL is a bronchodilator for the prevention and treatment of symptoms in chronic obstructive airway disorders with reversible airflow limitation such as bronchial asthma and especially chronic bronchitis with or without emphysema. Concomitant anti-inflammatory therapy should be considered for patients with bronchial asthma and steroid responsive chronic obstructive pulmonary disease (COPD).

Dosage and administration and Instruction for use

BERODUAL Solution for Inhalation

(1 mL contains 261 mcg ipratropium bromide + 500 mcg fenoterol hydrobromide)
(20 drops = 1 mL)

Dosage and administration

Treatment should be initiated and administered under medical supervision, e.g. in the hospital setting. Home based treatment can be recommended in patients when a low dose rapid acting beta-agonist bronchodilator such as BERODUAL metered dose inhaler has been insufficient in providing relief after consultation with an experienced physician. It can also be recommended in patients who are in need for nebuliser treatment for other reasons e.g. handling issues of metered dose inhaler or requirement of higher doses in experienced patients.

The treatment with the solution for inhalation should always be started with the lowest recommended dose. The dosage should be adapted to the individual requirements and tailored according to the severity of the acute episode. Administration should be stopped when sufficient symptom relief is achieved.

The following dosages are recommended:

Adults (including elderly) and adolescents >12 years of age:

Acute episodes of bronchospasm

Depending on the severity of the acute episode doses ranging between 261 mcg ipratropium bromide/500 mcg fenoterol hydrobromide (i.e. 1 mL = 20 drops) and 652.5 mcg ipratropium bromide/1250 mcg fenoterol hydrobromide (i.e. 2.5 mL = 50 drops) may be used. In exceptional particular severe cases doses up to 1044 mcg ipratropium bromide/2000 mcg fenoterol hydrobromide (i.e. 4 mL = 80 drops) may be used.

Children 6 - 12 years:

Acute asthma episodes

Depending on the severity of the acute episode and age doses ranging between 130.5 mcg ipratropium bromide/250 mcg fenoterol hydrobromide (i.e. 0.5 mL = 10 drops) and 522 mcg ipratropium bromide/1000 mcg fenoterol hydrobromide (i.e. 2 mL = 40 drops) may be used.

Children < 6 years (below 22 kg body weight):

Because there is limited information in this age group the following dose is recommended to be given under medical supervision only:

About 26.1 mcg ipratropium bromide/ 50 mcg fenoterol hydrobromide (i.e. 0.1 mL = 2 drops) per kg body weight = up to a maximum of 0.5 mL (=10 drops).

Instructions for use

The solution for inhalation is intended only for inhalation with suitable nebulising devices and must not be taken orally.

The recommended dose is to be diluted with physiological saline to a final volume of 3 - 4 mL and nebulised and inhaled until sufficient symptom relief is achieved.

BERODUAL solution for inhalation may, however, not be diluted with distilled water.

The solution should be freshly diluted each time before use; any residual diluted solution should be discarded.

The diluted solution should be inhaled directly after preparation of the solution.

The duration of inhalation can be controlled by the dilution volume.

BERODUAL solution for inhalation can be administered using a range of commercially available nebulising devices. The lung and systemic drug exposure is dependent on the nebuliser used and may be higher than with BERODUAL metered dose aerosol depending on the efficiency of the device.

Where wall oxygen is available the solution is best administered at a flow rate of 6 - 8 litres per minute.

The instructions provided by the manufacturer of the nebulising device for proper care, maintenance and cleaning of the equipment should be followed.

BERODUAL N Metered Dose Inhaler

Dosage and administration

The dosage should be adapted to the individual requirements. The following dosages are recommended for adults and children > 6 years:

Acute asthma episodes

21 mcg ipratropium bromide + 50 mcg fenoterol hydrobromide

2 actuations are sufficient for prompt symptom relief in many cases. In more severe cases, if breathing has not noticeably improved after 5 minutes, two further actuations may be taken.

If an attack has not been relieved by 4 actuations, further actuations may be required.

In these cases, patients should be advised to consult the doctor or the nearest hospital immediately.

Intermittent and long-term treatment (in asthma BERODUAL metered dose aerosol should be used only on an as-needed basis)

1 - 2 actuations for each administration, up to a maximum of 8 actuations per day (average 1 - 2 actuations 3 times daily).

In children BERODUAL metered dose aerosol should only be used on medical advice and under the supervision of an adult.

Patients should be instructed on the correct administration of the metered dose aerosol to ensure successful therapy (see Instructions for use).

Instruction for use:


Before first time use of the metered dose aerosol, the following rules should be observed:

Remove protective cap and depress the valve twice.

Before each use of the metered dose aerosol, the following rules should be observed:

1. Remove protective cap.
(If the inhaler has not been used for more than three days the valve has to be actuated once)
2. Breathe out deeply.

3. Hold the inhaler as shown in fig. 1, and close lips around the mouthpiece.
The arrow and the base of the container should be pointing upwards.



(Fig. 1)


4. Breathe in as deeply as possible, pressing the base of the canister firmly at the same time, this releases one metered dose. Hold the breath for a few seconds, then remove the mouthpiece from the mouth and breathe out. The same action should be repeated for a second inhalation.
5. Replace the protective cap after use.

The container is not transparent. It is therefore not possible to see when it is empty. The inhaler will deliver 200* puffs. When the labelled number of doses have been used the canister may still appear to contain a small amount of fluid. The inhaler should, however, be replaced so that you can be certain that you are getting the right amount of your medicine in each actuation.

Clean your mouthpiece at least once a week.


It is important to keep the mouthpiece of your inhaler clean to ensure that medicine does not build up and block the spray.

For cleaning, first take off the dust cap and remove the canister from the mouthpiece. Rinse with warm water through the mouthpiece until no medication build-up and/or dirt is visible.



(fig.2)

After cleaning shake out the mouthpiece and let it air-dry **without** using any heating system. Once the mouthpiece is dry, replace the canister and the dust cap.



(fig.3)

*300 [when the inhaler delivers 300 puffs]

WARNING:

The plastic mouthpiece has been specially designed for use with BERODUAL metered dose aerosol to ensure that you always get the right amount of the medicine. The mouthpiece must never be used with any other metered aerosol nor must the BERODUAL metered dose aerosol be used with any mouthpiece other than the one supplied with the product.

The container is under pressure and should by no account be opened by force or exposed to temperatures above 50°C.

Contraindications

BERODUAL is contraindicated in patients with known hypersensitivity to fenoterol hydrobromide or atropine-like substances or to any of the excipients of the product. BERODUAL is also contraindicated in patients with hypertrophic obstructive cardiomyopathy and tachyarrhythmia.

Special warnings and precautions

Hypersensitivity

Immediate hypersensitivity reactions may occur after administration of BERODUAL, as demonstrated by rare cases of urticaria, angio-oedema, rash, bronchospasm, oropharyngeal oedema and anaphylaxis.

Paradoxical bronchospasm

As with other inhaled medicines, BERODUAL may result in paradoxical bronchospasm that may be life-threatening. If paradoxical bronchospasm occurs BERODUAL should be discontinued immediately and substituted with an alternative therapy.

Ocular complications

BERODUAL should be used with caution in patients predisposed to narrow-angle glaucoma.

There have been isolated reports of ocular complications (i.e. mydriasis, increased intraocular pressure, narrow-angle glaucoma, eye pain) when aerosolised ipratropium bromide either alone or in combination with an adrenergic beta₂-agonist, has come in contact with the eyes.

Eye pain or discomfort, blurred vision, visual halos or coloured images in association with red eyes from conjunctival congestion and corneal oedema may be signs of acute narrow-angle glaucoma. Should any combination of these symptoms develop, treatment with miotic drops should be initiated and specialist advice sought immediately.

Thus patients must be instructed in the correct administration of BERODUAL. Care must be taken not to allow the product to enter the eyes.

For BERODUAL solution for inhalation, it is recommended that the nebulised solution is administered via a mouth piece. If this is not available and a nebuliser mask is used, it must fit properly. Patients who may be predisposed to glaucoma should be warned specifically to protect their eyes.

Systemic effects

In the following conditions BERODUAL should only be used after careful risk/benefit assessment, especially when doses higher than recommended are used:

Insufficiently controlled diabetes mellitus, recent myocardial infarction, severe organic heart or vascular disorders, hyperthyroidism, phaeochromocytoma, or with pre-existing urinary outflow tract obstruction (e.g. prostatic hyperplasia or bladder-neck obstruction).

Cardiovascular effects

Cardiovascular effects may be seen with sympathomimetic drugs, including BERODUAL. There is some evidence from post-marketing data and published literature of rare occurrences of myocardial ischaemia associated with beta-agonists. Patients with underlying severe heart disease (e.g. ischaemic heart disease, arrhythmia or severe heart failure) who are receiving BERODUAL, should be warned to seek medical advice if they experience chest pain or other symptoms of worsening heart disease. Attention should be paid to assessment of symptoms such as dyspnoea and chest pain, as they may be of either respiratory or cardiac origin.

Hypokalaemia

Potentially serious hypokalaemia may result from beta₂-agonist therapy (see also section Overdose).

Gastro-intestinal motility disturbances

Patients with cystic fibrosis may be more prone to gastro-intestinal motility disturbances.

Dyspnoea

In the case of acute, rapidly worsening dyspnoea, a physician should be consulted immediately.

Prolonged use:

- In patients with bronchial asthma BERODUAL should be used only on an as-needed basis. In patients with mild COPD on demand (symptom-oriented) treatment may be preferable to regular use.
- The addition or the increase of anti-inflammatory therapy to control airway inflammation and to prevent deterioration of disease control should be considered for patients with bronchial asthma and with steroid-responsive COPD.

The use of increasing amounts of beta₂-agonists containing products such as BERODUAL on a regular basis to control symptoms of bronchial obstruction may suggest declining disease control. If bronchial obstruction deteriorates it is inappropriate and possibly hazardous to simply increase the use of beta₂-agonist containing products such as BERODUAL beyond the recommended dose over extended periods of time. In this situation, the patient's therapy plan, and in particular the adequacy of anti-inflammatory therapy with inhaled corticosteroids, should be reviewed to prevent potentially life-threatening deterioration of disease control.

Concomitant use with other sympathomimetic bronchodilators

Other sympathomimetic bronchodilators should only be used with BERODUAL under medical supervision (see section Interactions).

Doping warning

The use of BERODUAL may lead to positive results with regard to fenoterol in tests for non-clinical substance abuse, e.g. in the context of athletic performance enhancement (doping).

Excipients

BERODUAL Solution for Inhalation

This product contains 0.1mg benzalkonium chloride in each mL. Benzalkonium chloride may cause wheezing and breathing difficulties. Patients with asthma are at an increased risk for these adverse events.

BERODUAL N Metered Dose Inhaler

This medicine contains about 13 mg of alcohol (ethanol) in each actuation. The amount in each actuation of this medicine is equivalent to less than 1 ml beer or 1 ml wine.

Interactions

The chronic co-administration of BERODUAL with other anticholinergic drugs has not been studied. Therefore, the chronic co-administration of BERODUAL with other anticholinergic drugs is not recommended.

Other beta-adrenergics, anticholinergics and xanthine derivatives (such as theophylline) may enhance the bronchodilatory effect. The concurrent administration of other beta-mimetics, systemically available anticholinergics and xanthine derivatives (e.g. theophylline) may increase the adverse reactions.

A potentially serious reduction in bronchodilatation may occur during concurrent administration of beta-blockers.

Hypokalaemia induced by beta₂-agonists may be increased by concomitant treatment with xanthine derivatives, corticosteroids, and diuretics. This should be taken into account particularly in patients with severe airway obstruction.

Hypokalaemia may result in an increased susceptibility to arrhythmias in patients receiving digoxin. Additionally, hypoxia may aggravate the effects of hypokalaemia on cardiac rhythm. It is recommended that serum potassium levels are monitored in such situations.

Beta₂-agonist containing medicinal products should be administered with caution to patients being treated with monoamine oxidase inhibitors or tricyclic antidepressants, since the action of beta-adrenergic agonists may be enhanced.

Inhalation of halogenated hydrocarbon anaesthetics such as halothane, trichloroethylene and enflurane may increase the susceptibility on the cardiovascular effects of beta-agonists.

Fertility, pregnancy and lactation

Pregnancy

Non-clinical data, combined with available experience in humans have shown no evidence of adverse effects in pregnancy of fenoterol or ipratropium. Nonetheless, the usual precautions regarding the use of drugs during pregnancy, especially during the first trimester, should be exercised.

The inhibitory effect of fenoterol on uterine contraction should be taken into account.

Lactation

Non-clinical studies have shown that fenoterol hydrobromide, is excreted into breast milk. It is unknown whether ipratropium is excreted into breast milk. But it is unlikely that ipratropium would reach the infant to an important extent, especially when taken by aerosol. However, caution should be exercised when BERODUAL is administered to a nursing woman.

Fertility

Clinical data on fertility are neither available for the combination of ipratropium bromide and fenoterol hydrobromide nor for each of the two components of the combination. Non-clinical studies performed with the individual components ipratropium bromide and fenoterol hydrobromide showed no adverse effect on fertility.

Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

However, patients should be advised that they may experience undesirable effects such as dizziness, tremor, accommodation disorder, mydriasis and blurred vision during treatment with BERODUAL. Therefore, caution should be recommended when driving a car or operating machinery.

Side effects

Many of the listed undesirable effects can be assigned to the anticholinergic and beta-adrenergic properties of BERODUAL. As with all inhalation therapy BERODUAL may show symptoms of local irritation.

The most frequent side effects reported in clinical trials were cough, dry mouth, headache, tremor, pharyngitis, nausea, dizziness, dysphonia, tachycardia, palpitations, vomiting, blood pressure systolic increased and nervousness.

The following adverse reactions have been reported during the use of BERODUAL in clinical trials and during the post-marketing experience.

Immune system disorders

- anaphylactic reaction
- hypersensitivity

Metabolism and nutritional disorders

- hypokalemia

Psychiatric disorders

- nervousness
- agitation
- mental disorder

Nervous system disorders

- headache
- tremor
- dizziness

Eye disorders

- glaucoma
- intraocular pressure increased
- accommodation disorder
- mydriasis
- vision blurred
- eye pain
- corneal oedema
- conjunctival hyperaemia
- halo vision

Cardiac disorders

- tachycardia, heart rate increased
- palpitations
- arrhythmia
- atrial fibrillation
- supraventricular tachycardia
- myocardial ischaemia

Respiratory, thoracic and mediastinal disorders

- cough
- pharyngitis
- dysphonia
- bronchospasm
- throat irritation
- pharyngeal oedema
- laryngospasm
- bronchospasm paradoxical
- dry throat

Gastrointestinal disorders

- vomiting
- nausea
- dry mouth
- stomatitis
- glossitis
- gastrointestinal motility disorder
- diarrhoea
- constipation
- oedema mouth

Skin and subcutaneous tissue disorders

- urticaria
- rash
- pruritus
- angioedema
- hyperhidrosis

Musculoskeletal and connective tissue disorders

- muscular weakness
- muscle spasms

- myalgia

Renal and urinary disorders

- urinary retention

Investigations

- blood pressure systolic increased
- blood pressure diastolic decreased

Overdose

Symptoms

The effects of overdose are expected to be primarily related to fenoterol. The expected symptoms with overdose are those of excessive beta-adrenergic-stimulation, the most prominent being tachycardia, palpitation, tremor, hypertension, hypotension, widening of the pulse pressure, anginal pain, arrhythmias, and flushing. Metabolic acidosis and hypokalaemia have also been observed with fenoterol when applied in doses higher than recommended for the approved indications of BERODUAL.

Expected symptoms of overdose with ipratropium bromide (such as dry mouth, visual accommodation disorder) are mild because the systemic availability of inhaled ipratropium is very low.

Therapy

Treatment with BERODUAL should be discontinued. Acid base and electrolyte monitoring should be considered. Administration of sedatives and in severe cases intensive care treatment may be needed.

Beta-receptor blockers, preferably beta₁-selective, are suitable as specific antidotes; however, a possible increase in bronchial obstruction must be taken into account and the dose should be adjusted carefully in patients suffering from bronchial asthma or COPD because of the risk of precipitating severe bronchospasm, which may be fatal.

Availability

BERODUAL Solution for Inhalation

Bottle of 20ml

BERODUAL N Metered Dose Inhaler

200 metered doses (10ml)

Store below 30°C.

Please refer to the packaging for information on shelf-life.

Manufactured by

BERODUAL Solution for Inhalation

Manufactured by

Istituto De Angeli S.r.L.

Florence, Italy

for

Boehringer Ingelheim International GmbH

Ingelheim am Rhein

Germany

BERODUAL N Metered Dose Inhaler

Manufactured by

Boehringer Ingelheim Pharma GmbH & Co. KG

Ingelheim am Rhein, Germany

for

Boehringer Ingelheim International GmbH
Ingelheim am Rhein
Germany

Date of Revision: 18 April 2023