

# DHACOPAN

PHARMACODE  
AREA

## PRODUCT DESCRIPTION:

### Tablet:

Orange film-coated tablets, round and deep convex, 8mm diameter without score-line and logo.

### Syrup:

Clear yellow syrup, sweet taste with banana flavor.

quaternary ammonium

anticholinergic agent with both central and peripheral actions.

The peripheral effects include decreased production of saliva, sweat, bronchial, lacrimal, nasal, gastric and intestinal secretions, decreased intestinal motility and inhibition of micturition.

## COMPOSITION:

- 1) Each tablet contains:  
Hyoscine Butylbromide  
BP 10mg
- 2) Each 5ml contains:  
Hyoscine Butylbromide  
BP 5mg/5ml

## INDICATION:

For relief of gastro-intestinal and colonic spasm, acute spasm in renal or biliary colic; dysmenorrhoea.

## Excipients:

Dhacopan Tablet 10mg:  
Corn Starch, Lactose mesh 200, Polyvidone 30, Magnesium stearate, Titanium Dioxide, Methocel E15, Sunset Yellow No.6 Alum lake, Propylene Glycol.

## RECOMMENDED DOSAGE:

### Tablets:

Adults and Children more than 12 years: 1 to 2 tablets, 3 or 4 times a day.

### Syrup:

Children: 6 to 12 years:  
2 teaspoonfuls 3 times a day.

Dhacopan Syrup 5mg/5ml:  
Granulated Fine White Sugar, Sodium Saccharin, Propyl Hydroxybenzoate, Methyl Hydroxybenzoate, Propylene Glycol, Sodium Citrate Dihydrate, Citric Acid Monohydrate, FDC Yellow No. 6 Water Soluble (40-44%), Quinoline Yellow E104 Water Soluble 70-72%, Banana Flavour PCA 5639, Alcohol 96%, Purified Water.

## SIDE EFFECTS:

Dry mouth with difficulty in swallowing and thirst, increased intraocular pressure, flushing, dry skin, difficulty in micturition, constipation, palpitation and arrhythmias.

## PHARMACOLOGY:

Dhacopan contains Hyoscine Butylbromide which is a

## CONTRA-INDICATIONS:

Contra-indicated in patients with glaucoma; with gastric or duodenal ulcer, with urinary retention; prostatic enlargement; paralytic ileus; pyloric stenosis.

**CAUTION:**

Tachycardia and cardiac insufficiency. May cause drowsiness, if affected do not drive or operate machinery. Avoid alcohol.

**USE IN PREGNANCY AND LACTATION:**

There are no adequate studies in pregnant and lactating women. Therefore, Hyoscine Butylbromide should not be used in pregnant and lactating women unless the potential advantage justifies the possible risk to the foetus.

**DRUG INTERACTIONS:**

Hyoscine Butylbromide and Metoclopramide have opposing effects on the gastro-intestinal activity.

**INCOMPATIBILITY:**

No known report.

**OVERDOSAGE AND TREATMENT:**

Anticholinergic symptoms may occur e.g. urinary retention, dry mouth and tachycardia. Oral poisoning may be treated by gastric lavage with medicinal charcoal followed by magnesium sulphate (15%). Symptoms of overdose can be countered with parasympathomimetics.

**AVAILABILITY:**

Tablets:  
10x10x10's in blister pack

Syrup:  
3.8 litre in high density polyethylene plastic bottle.

**STORAGE CONDITION:**

- 1) Tablets:  
Keep below 30°C. Protect from light.
- 2) Syrup:  
Keep below 25°C. Protect from light.

**EXPIRY PERIOD:**

Do not use after the expiry date stated on the label.

**DATE OF REVISION:**

September 2021

For further information, please consult your physician or pharmacist.

Manufactured By:  
PT Actavis Indonesia  
Jl. Raya Bogor Km 28  
Jakarta Timur 13710  
Indonesia