For the use only of a Registered

Medical Practitioner or a Hospital

Biocon Biologics

Biocon Biologics

1 vial with 10 mL of solution

COMPOSITION

1 ml solution contains 100 units (equivalent to 3.5 mg) insulin aspart*.

Kirsty 100 units/ml solution for injection in vial Each vial contains 10 ml equivalent to 1,000 units.

*Insulin aspart is produced in Pichia pastoris by recombinant DNA technology.

For the full list of excipients, see "List of Excipients" section.

PHARMACEUTICAL FORM

Solution for injection (injection).

The solution is clear, colourless and aqueous. PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Pharmacotherapeutic group - Drugs used in diabetes. Insulins and analogues for injection, fast acting. ATC code: A10AB05.

Kirsty is a biosimilar medicinal product.

Mechanism of action and pharmacodynamic effects The blood glucose lowering effect of insulin aspart is due to the facilitated uptake of glucose following binding of insulin to receptors on muscle and

fat cells and to the simultaneous inhibition of glucose output from the

Insulin aspart produces a more rapid onset of action compared to soluble human insulin, together with a lower glucose concentration, as assessed within the first four hours after a meal. Insulin aspart has a shorter duration of action compared to soluble human insulin after subcutaneous injection.




Fig. 1. Blood glucose concentrations following a single pre meal dose of insulin aspart injected immediately before a meal (solid curve) or soluble human insulin administered 30 minutes before a meal (hatched curve) in patients with type 1 diabetes mellitus.

When insulin aspart is injected subcutaneously, the onset of action will occur within 10 to 20 minutes of injection. The maximum effect is exerted between 1 and 3 hours after injection. The duration of action is 3 to 5

Clinical efficacy and safety

Clinical trials in patients with type 1 diabetes have demonstrated a lower postprandial blood glucose with insulin aspart compared to soluble human insulin (Fig. I). In two long term open label trials in patients with type 1 diabetes comprising 1070 and 884 patients, respectively, insulin aspart reduced glycated haemoglobin by 0.12 [95% C.I. 0.03; 0.22] percentage points and by 0.15 [95% C.I. 0.05; 0.26] percentage points compared to human insulin; a difference of limited clinical significance.

Clinical trials in patients with type 1 diabetes have demonstrated a reduced risk of nocturnal hypoglycaemia with insulin aspart compared with soluble human insulin. The risk of daytime hypoglycaemia was not significantly increased.

Insulin aspart is equipotent to soluble human insulin on a molar basis.

Special populations Elderly (≥ 65 years old)

A randomised, double blind cross over PK/PD trial comparing insulin aspart with soluble human insulin was performed in elderly patients with type 2 diabetes (19 patients aged 65 83 years, mean age 70 years). The relative differences in the pharmacodynamic properties (GIRmax, AUCGIR, 0 120 min) between insulin aspart and soluble human insulin in the elderly were similar to those seen in healthy subjects and in younger patients with diabetes.

Paediatric population

A clinical trial comparing preprandial soluble human insulin with postprandial insulin aspart was performed in small children (20 patients aged 2 to less than 6 years, studied for 12 weeks, among those four were younger than 4 years old) and a single dose PK/PD trial was performed in

children (6 12 years) and adolescents (13 17 years). The pharmacodynamic profile of insulin aspart in children was similar to that seen in adults.

Size: 750 (W) x 440 (H) mm; Colour: PANTONE Black C

The efficacy and safety of insulin aspart given as bolus insulin in

combination with either insulin detemir or insulin degludec as basal insulin has been studied for up to 12 months, in two randomised controlled clinical trials in adolescents and children aged 1 to less than 18 years (n=712). The trials included 167 children aged 1 5 years, 260 aged 6 11 and 285 aged 12 17. The observed improvements in HbA1c and the safety profiles were comparable between all age groups.

A clinical trial comparing safety and efficacy of insulin aspart vs. human insulin in the treatment of pregnant women with type 1 diabetes (322 exposed pregnancies (insulin aspart: 157; human insulin: 165)) did not indicate any adverse effect of insulin aspart on pregnancy or on the health of the foetus/newborn.

In addition, the data from a clinical trial including 27 women with gestational diabetes randomised to treatment with insulin aspart vs. human insulin (insulin aspart: 14; human insulin: 13) showed similar safety profiles between treatments.

Pharmacokinetic properties

Absorption, distribution and elimination

In Kirsty substitution of amino acid proline with aspartic acid at position B28 reduces the tendency to form hexamers as observed with soluble human insulin. Kirsty is therefore more rapidly absorbed from the subcutaneous layer compared to soluble human insulin.

The time to maximum concentration is, on average, half of that for soluble human insulin. A mean maximum plasma concentration of 492±256 pmol/I was reached 40 (interquartile range: 30 40) minutes after a subcutaneous dose of 0.15 unit/kg bodyweight in type 1 diabetic patients. The insulin concentrations returned to baseline about 4 to 6 hours after dose. The absorption rate was somewhat slower in type 2 diabetic patients, resulting in a lower Cmax (352±240 pmol/l) and later tmax (60 (interquartile range: 50 90) minutes). The intra individual variability in time to maximum concentration is significantly less for Kirsty

than for soluble human insulin, whereas the intra individual variability in

Cmax for Kirsty is larger. Special populations Elderly (≥ 65 years old)

The relative differences in pharmacokinetic properties between insulin aspart and soluble human insulin in elderly patients (65 83 years, mean age 70 years) with type 2 diabetes were similar to those observed in healthy subjects and in younger patients with diabetes. A decreased absorption rate was observed in elderly patients, resulting in a later tmax (82 (interquartile range: 60 1 20) minutes), whereas Cmax was similar to that observed in younger patients with type 2 diabetes and slightly lower

than in patients with type 1 diabetes.

A single dose pharmacokinetic study of insulin aspart was performed in 24 subjects with hepatic function ranging from normal to severely impaired. In patients with hepatic impairment, absorption rate was decreased and more variable, resulting in delayed tmax from about 50 min in subjects with normal hepatic function to about 85 min in patients with moderate and severe hepatic impairment. AUC, Cmax and CL/F were similar in patients with reduced hepatic function compared with subjects with normal hepatic function.

Renal impairment

A single dose pharmacokinetic study of insulin aspart in 18 subjects with renal function ranging from normal to severely impaired was performed. No apparent effect of creatinine clearance values on AUC, Cmax, CL/F and tmax of insulin aspart was found. Data were limited in patients with moderate and severe renal impairment. Patients with renal failure

necessitating dialysis treatment were not investigated.

Paediatric population The pharmacokinetic and pharmacodynamic properties of insulin aspart were investigated in children (6 12 years) and adolescents (13 17 years) with type 1 diabetes. Insulin aspart was rapidly absorbed in both age groups, with similar tmax as in adults. However, Cmax differed between the age groups, stressing the importance of the individual titration of

Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity and toxicity to reproduction and development.

In in vitro tests, including binding to insulin and IGF 1 receptor sites and effects on cell growth, insulin aspart behaved in a manner that closely resembled human insulin. Studies also demonstrate that the dissociation of binding to the insulin receptor of insulin aspart is equivalent to human

CLINICAL PARTICULARS

Therapeutic indications

Kirsty is indicated for treatment of diabetes mellitus in adults, adolescents and children aged 1 year and above.

Posology and method of administration

infusion (CSII) in pump systems

Folding Size: 43.33mm-9 folds, 30mm-1 fold, 20mm-1 fold x 150 (H) mm with pasting

Folding & Pharmacode position to be followed as per Drawing No. 70M1806504

Note: Booklet to be sealed with 2 stickers; Paper: 60 gsm Maplitho / SMPO Paper

The potency of insulin analogues, including insulin aspart, is expressed in units, whereas the potency of human insulin is expressed in international

Kirsty dosing is individual and determined in accordance with the needs of the patient. It should normally be used in combination with intermediate acting or long acting insulin. Moreover, Kirsty vial can be used for continuous subcutaneous insulin

Kirsty vial can also be used if intravenous administration of insulin aspart, by physicians or other healthcare staff, is applicable. Blood glucose monitoring and insulin dose adjustments are

recommended to achieve optimal glycaemic control. The individual insulin requirement in adults and children is usually between 0.5 and 1.0 unit/kg/day. In a basal bolus treatment regimen 50

70% of this requirement may be provided by Kirsty and the remainder by intermediate acting or long acting insulin. Adjustment of dose may be necessary if patients undertake increased physical activity, change their usual diet or during concomitant illness. **Special populations**

Elderly (≥ 65 years old) Kirsty can be used in elderly patients.

In elderly patients, glucose monitoring should be intensified, and the insulin aspart dose adjusted on an individual basis. Renal impairment

Renal impairment may reduce the patient's insulin requirements. In patients with renal impairment, glucose monitoring should be intensified, and the insulin aspart dose adjusted on an individual basis. Hepatic impairment

Hepatic impairment may reduce the patient's insulin requirements. In patients with hepatic impairment, glucose monitoring should be intensified, and the insulin aspart dose adjusted on an individual basis.

Kirsty can be used in children and adolescents aged 1 year and above in preference to soluble human insulin when a rapid onset of action might be beneficial, for example, in the timing of the injections in relation to meals (see Pharmacodynamic Properties and Pharmacokinetics

The safety and efficacy of Kirsty in children below 1 year of age have not No data are available

Transfer from other insulin medicinal products

When transferring from other insulin medicinal products, adjustment of the Kirsty dose and the dose of the basal insulin may be necessary. Kirsty has a faster onset and a shorter duration of action than soluble human insulin. When injected subcutaneously into the abdominal wall, the onset of action will occur within 10 20 minutes of injection. The maximum effect is exerted between 1 and 3 hours after the injection. The duration of

Close glucose monitoring is recommended during the transfer and in the initial weeks thereafter (see Special warnings and precautions for use

Method of administration

Insulin aspart is a rapid acting insulin analogue.

Kirsty is administered subcutaneously by injection in the abdominal wall, the thigh, the upper arm, the deltoid region or the gluteal region. Injection sites should always be rotated within the same region in order to reduce the risk of lipodystrophy and cutaneous amyloidosis (see Special warnings and precautions for use and Undesirable effects sections).

Subcutaneous injection in the abdominal wall ensures a faster absorption than other injection sites. Compared to soluble human insulin the faster onset of action of Kirsty is maintained regardless of the injection site. The duration of action will vary according to the dose, injection site, blood

flow, temperature and level of physical activity. Due to the faster onset of action, Kirsty should generally be given immediately before a meal. When necessary Kirsty can be given soon after a meal.

Kirsty 100 units/ml solution for injection in vial Continuous subcutaneous insulin infusion (CSII)

Kirsty may be used for CSII in pump systems suitable for insulin infusion. CSII should be administered in the abdominal wall. Infusion sites should

When used with an insulin infusion pump, Kirsty should not be mixed with any other insulin medicinal products. Patients using CSII should be comprehensively instructed in the use of

the pump system and use the correct reservoir and tubing for the pump (see Special precautions for disposal and other handling section). The infusion set (tubing and cannula) should be changed in accordance with the instructions in the product information supplied with the infusion set. Patients administering Kirsty by CSII must have an alternative insulin delivery method available in case of pump system failure.

If necessary, Kirsty can be administered intravenously which should be carried out by physicians or other healthcare professionals. For intravenous use, infusion systems with Kirsty 100 units/ml at concentrations from 0.05 unit/ml to 1.0 unit/ml insulin aspart in the solutions for infusion sodium chloride 9 mg/ml (0.9%), 5% glucose or

10% glucose including 40 mmol/l potassium chloride using polypropylene infusion bags, are stable at room temperature for 24 hours. Although stable over time, a certain amount of insulin will be initially adsorbed to the material of the infusion bag. Monitoring of blood glucose is necessary during insulin infusion.

Administration with a syringe

Kirsty vials are for use with insulin syringes with the corresponding unit scale. See also Special precautions for disposal and other handling

For detailed user instructions, please refer to the package leaflet. Hypersensitivity to the active substance or to any of the excipients listed in List of excipients section.

Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered medicinal product should be clearly recorded.

Before travelling between different time zones, the patient should seek the doctor's advice since this may mean that the patient has to use the insulin and meals at different times.

Hyperglycaemia

Inadequate dosing or discontinuation of treatment, especially in type 1 diabetes, may lead to hyperglycaemia and diabetic ketoacidosis. Usually the first symptoms of hyperglycaemia develop gradually over a period of hours or days. They include thirst, increased frequency of urination, nausea, vomiting, drowsiness, flushed dry skin, dry mouth, loss of appetite as well as acetone odour of breath. In type 1 diabetes, untreated hyperglycaemic events eventually lead to diabetic ketoacidosis, which is potentially lethal.

Hypoglycaemia Omission of a meal or unplanned, strenuous physical exercise may lead

to hypoglycaemia. Especially in children, care should be taken to match insulin doses (especially in basal bolus regimens) with food intake, physical activities and current blood glucose level in order to minimise the risk of

Hypoglycaemia may occur if the insulin dose is too high in relation to the insulin requirement. In case of hypoglycaemia or if hypoglycaemia is suspected Kirsty must not be injected. After stabilization of patient's blood glucose adjustment of the dose should be considered (see **Undesirable effects** and **Overdose** sections).

Patients whose blood glucose control is greatly improved, e.g. by intensified insulin therapy, may experience a change in their usual warning symptoms of hypoglycaemia, and should be advised accordingly. Usual warning symptoms may disappear in patients with longstanding diabetes.

A consequence of the pharmacodynamics of rapid acting insulin analogues is that if hypoglycaemia occurs, it may occur earlier after an injection when compared with soluble human insulin. Since Kirsty should be administered in immediate relation to a meal, the rapid onset of action should be considered in patients with concomitant

diseases or treatment where a delayed absorption of food might be Concomitant illness, especially infections and feverish conditions, usually increases the patient's insulin requirements. Concomitant diseases in the

kidney, liver or affecting the adrenal, pituitary or thyroid gland can require

changes in the insulin dose. When patients are transferred between different types of insulin medicinal products, the early warning symptoms of hypoglycaemia may change or become less pronounced than those experienced with their previous

Transfer from other insulin medicinal products

Transferring a patient to another type or brand of insulin should be done under strict medical supervision. Changes in strength, brand (manufacturer), type, origin (animal, human insulin or human insulin analogue) and/or method of manufacture (recombinant DNA versus animal source insulin) may result in the need for a change in dose. Patients transferred to Kirsty from another type of insulin may require an increased number of daily injections or a change in dose from that used with their usual insulin medicinal products. If an adjustment is needed, it may occur with the first dose or during the first few weeks or months.

<u>Injection site reactions</u>

As with any insulin therapy, injection site reactions may occur and include pain, redness, hives, inflammation, bruising, swelling and itching. Continuous rotation of the injection site within a given area reduces the risk of developing these reactions. Reactions usually resolve in a few days to a few weeks. On rare occasions, injection site reactions may require discontinuation of Kirsty.

Skin and subcutaneous tissue disorders

Patients must be instructed to perform continuous rotation of the injection site to reduce the risk of developing lipodystrophy and cutaneous amyloidosis. There is a potential risk of delayed insulin absorption and worsened glycaemic control following insulin injections at sites with these reactions. A sudden change in the injection site to an unaffected area has been reported to result in hypoglycaemia. Blood

glucose monitoring is recommended after the change in the injection site from an affected to an unaffected area, and dose adjustment of antidiabetic medications may be considered.

Combination of insulin aspart with pioglitazone

Cases of cardiac failure have been reported when pioglitazone was used in combination with insulin, especially in patients with risk factors for development of cardiac heart failure. This should be kept in mind if treatment with the combination of pioglitazone and insulin aspart is considered. If the combination is used, patients should be observed for signs and symptoms of heart failure, weight gain and oedema. Pioglitazone should be discontinued if any deterioration in cardiac symptoms occurs.

Avoidance of accidental mix ups/medication errors

Patients must be instructed to always check the insulin label before each injection to avoid accidental mix ups between insulin aspart and other insulin products.

<u>Insulin antibodies</u>

Insulin administration may cause insulin antibodies to form. In rare cases the presence of such insulin antibodies may necessitate adjustment of the insulin dose in order to correct a tendency to hyper or hypoglycaemia.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium free'.

Interaction with other medicinal products and other forms of interaction

A number of medicinal products are known to interact with the glucose

The following substances may reduce the patient's insulin requirements: Oral antidiabetic medicinal products, monoamine oxidase inhibitors (MAOI), beta blockers, angiotensin converting enzyme (ACE) inhibitors, salicylates, anabolic steroids and sulphonamides.

The following substances may increase the patient's insulin requirements: Oral contraceptives, thiazides, glucocorticoids, thyroid hormones, sympathomimetics, growth hormone and danazol.

Octreotide/lanreotide may either increase or decrease the insulin Alcohol may intensify or reduce the hypoglycaemic effect of insulin.

Beta blockers may mask the symptoms of hypoglycaemia.

Fertility, pregnancy, and lactation

Kirsty can be used in pregnancy. Data from two randomized controlled clinical trials (322 and 27 exposed pregnancies) do not indicate any adverse effect of insulin aspart on pregnancy or on the health of the foetus/newborn when compared to human insulin (see

Pharmacodynamic properties section). Intensified blood glucose control and monitoring of pregnant women with diabetes (type 1 diabetes, type 2 diabetes or gestational diabetes) are recommended throughout pregnancy and when contemplating pregnancy. Insulin requirements usually fall in the first trimester and

increase subsequently during the second and third trimester. After

delivery, insulin requirements normally return rapidly to pre pregnancy

There are no restrictions on treatment with Kirsty during breast feeding Insulin treatment of the nursing mother presents no risk to the baby.

Animal reproduction studies have not revealed any differences between insulin aspart and human insulin regarding fertility.

Effects on ability to drive and use machines

However, the Kirsty dose may need to be adjusted.

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating

Patients should be advised to take precautions to avoid hypoglycaemia while driving. This is particularly important in those who have reduced or absent awareness of the warning signs of hypoglycaemia or have frequent episodes of hypoglycaemia. The advisability of driving should be considered in these circumstances.

Undesirable effects

Summary of the safety profile Adverse reactions observed in patients using insulin aspart are mainly due to the pharmacologic effect of insulin.

The most frequently reported adverse reaction during treatment is hypoglycaemia. The frequencies of hypoglycaemia vary with patient population, dose regimens and level of glycaemic control (see **Undesirable effects** section, Description of selected adverse reactions). At the beginning of the insulin treatment, refraction anomalies, oedema and injection site reactions (pain, redness, hives, inflammation, bruising, swelling and itching at the injection site) may occur. These reactions are usually of transitory nature. Fast improvement in blood glucose control may be associated with acute painful neuropathy, which is usually

reversible. Intensification of insulin therapy with abrupt improvement in

glycaemic control may be associated with temporary worsening of

diabetic retinopathy, while long term improved glycaemic control decreases the risk of progression of diabetic retinopathy.

<u>Tabulated list of adverse reactions</u> Adverse reactions listed below are based on clinical trial data and classified according to MedDRA frequency and System Organ Class. Frequency categories are defined according to the following convention: Very common (\geq 1/10); common (\geq 1/100 to < 1/10); uncommon (\geq 1/1,000 to < 1/100); rare ($\ge 1/10,000 \text{ to} < 1/1,000$); very rare (< 1/10,000);

not known (cannot be estimated from the available data).

MedDRA system organ class	Very Common	Uncommon	Rare	Very Rare	Not Known
Immune system disorders		Urticaria, rash, eruptions		Anaphylactic reactions*	
Metabolism and nutrition's disorders	Hypoglycaemia*				
Nervous system disorders			Peripheral neuropathy (painful neuropathy)		
Eye disorders		Refraction disorders, diabetic retinopathy			
Skin and subcutaneous tissue disorders		Lipodystrophy*			Cutaneous amyloidosis*†
General disorders and administration site conditions		Injection site reactions, oedema			

*see **Undesirable effects** section, Description of selected adverse reactions.

† adverse drug reaction (ADR) from postmarketing sources.

The occurrence of generalised hypersensitivity reactions (including generalized skin rash, itching, sweating, gastrointestinal upset, angioneurotic oedema, difficulties in breathing, palpitation and reduction

<u>Hypoglycaemia</u> The most frequently reported adverse reaction is hypoglycaemia. It may occur if the insulin dose is too high in relation to the insulin requirement Severe hypoglycaemia may lead to unconsciousness and/or convulsions and may result in temporary or permanent impairment of brain function or

even death. The symptoms of hypoglycaemia usually occur suddenly. They may include cold sweats, cool pale skin, fatigue, nervousness or tremor, anxiousness, unusual tiredness or weakness, confusion, difficulty in concentration, drowsiness, excessive hunger, vision changes,

In clinical trials, the frequency of hypoglycaemia varied with patient population, dose regimens and level of glycaemic control. During clinical trials the overall rates of hypoglycaemia did not differ between patients

Lipodystrophy (including lipohypertrophy, lipoatrophy) and cutaneous amyloidosis may occur at the injection site and delay local insulin absorption. Continuous rotation of the injection site within the given injection area may help to reduce or prevent these reactions (see Special warnings and precautions for use section).

Paediatric population

Based on post marketing sources and clinical trials, the frequency, type and severity of adverse reactions observed in the paediatric population do not indicate any differences to the broader experience in the general

Based on post marketing sources and clinical trials, the frequency, type and severity of adverse reactions observed in the elderly patients and in patients with renal or hepatic impairment do not indicate any differences to the broader experience in the general population.

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

A specific overdose for insulin cannot be defined, however, hypoglycaemia may develop over sequential stages if too high doses

relative to the patient's requirement are administered

Description of selected adverse reactions Anaphylactic reactions

in blood pressure) is very rare but can potentially be life threatening.

headache, nausea and palpitation. treated with insulin aspart compared to human insulin.

Skin and subcutaneous tissue disorders

population.

Other special populations

Reporting of suspected adverse reactions

McDermott Laboratories T/A Mylan Dublin Biologics Newenham Court Northern Cross Malahide Road Dublin, 17, Ireland

Johor, 79200, Iskandar Puteri (Formerly Nusajaya), Malaysia



Chemical and physical in use stability has been demonstrated for 31 days

at 30°C and 5°C. From a microbiological point of view, once opened, the

Keep the medicinal product in the outer carton in order to protect from light.

For storage conditions of the medicinal product after first opening, see

10 ml solution in vial (type 1 glass) closed with a bromobutyl rubber

Pack sizes of 1, or 5 vials or a multipack containing 5 (5 packs of 1) vials.

Do not use this medicinal product if you notice that the solution is not

The patient should be advised to discard the needle after each injection.

Kirsty may be used in an infusion pump system (CSII) as described in

Posology and method of administration section. Tubings in which the

Name and address of the manufacturer(s) of the biological active

Name and address of the manufacturer responsible for batch release

inner surface materials are made of polyethylene or polyolefin have been

Any unused medicinal product or waste material should be disposed of in

medicinal product may be stored for a maximum of 28 days at 30°C.

Other in use storage times are the responsibility of the user.

Keep the vial in the outer carton in order to protect from light.

Kirsty 100 units/ml solution for injection in vial

Store in a refrigerator (2°C 8°C). Do not freeze.

Kirsty 100 units/ml solution for injection in vial

Special precautions for disposal and other handling

Needles, syringes, and pre-filled pens must not be shared.

Kirsty which has been frozen must not be used.

Kirsty 100 units/ml solution for injection in vial

evaluated and found compatible with pump use.

Kawasan Perindusterian SiLC, 79200 Iskandar Puteri,

No. 1, Jalan Bioteknologi 1, Kawasan perindustrian SiLC,

Special precautions for storage

Nature and contents of container

stopper and aluminium flip off seal.

clear, colourless and aqueous.

accordance with local requirements.

Shelf life section.

substance(s)

Biocon Sdn. Bhd.,

Johor, Malaysia

Marketed by

Biocon SDN BHD;

No. 1, Jalan Bioteknologi 1,

Store below 30°C. Do not refrigerate. Do not freeze.



BM/XXXX/XX 1 vial with 10 mL of solution Mild hypoglycaemic episodes can be treated by oral administration of

Insulin Aspart for Injection (r-DNA origin)

Solution for Subcutaneous Injection

glucose or sugary products. It is therefore recommended that the diabetic patient always carries sugar containing products. • Severe hypoglycaemic episodes, where the patient has become Biocon House - Semicon Park unconscious, can be treated with glucagon (0.5 to 1 mg) given Electronic city Phase II intramuscularly or subcutaneously by a trained person, or with glucose Bangalore – 560100 given intravenously by physicians or other healthcare staff. Glucose must be given intravenously, if the patient does not respond to glucagon within

in Vial

relapse. PHARMACEUTICAL PARTICULARS

10 to 15 minutes. Upon regaining consciousness, administration of oral

carbohydrates is recommended for the patient in order to prevent a

List of excipients Phenol Metacresol

Disodium phosphate dihydrate Sodium chloride

Sodium hydroxide (for pH adjustment) Water for injections Incompatibilities

products. Shelf life Before opening 30 months

After first opening

28 days

Medical Advisor / Medical Affairs **Biocon Biologics Ltd**

Note: Unless otherwise stated and claimed, the data related to the studies, tests, treatment and application contained herein are from the published databases.

Kirsty® is registered by Biocon Biologics Limited.

To report adverse events and/or product complaints visit our website www.biocon.com or e-mail us at DrugSafety@biocon.com

Leaflet generated August 2023

⋘ Biocon Biologics

¬

750 mm

Note: Booklet to be sealed with 2 numbers of transparent perforated stickers.

Size:13x30mm plain perforated tape WITH MATT VARNISH Material: UPM 6L/RP37/01 PP CLEAR TC50/RP37/WG65

Z:\EM\3rd SET\Singapore\Kirsty

Hydrochloric acid (for pH adjustment)

This medicinal product must not be diluted or mixed with other medicinal