NUROFEN COATED TABLETS 200mg

(Ibuprofen 200mg)

PLEASE READ THIS ENTIRE LEAFLET CAREFULLY BEFORE YOU START TO TAKE YOUR MEDICINE.

Keep this leaflet. You may want to read it again.

WHAT IS IN THIS PACK?

Each tablet of Nurofen Coated Tablets 200mg contains the

active ingredient Ibuprofen 200mg.

Also contains: Tablet core: Croscarmellose sodium, sodium laurilsulfate, sodium citrate, stearic acid, colloidal anhydrous slitica. Sugar coat: Carmellose sodium, french chalk for tablets (falc), accaic spray dried, sucrose, titanium dioxide, macrogal 6000. Printing: Black printing ink (contains shellac, iron oxide black (E172) and propylene glycol).

Manufactured by:
Reckilt Benckiser Healthcare International Limited
Thane Road, Nottingham, Nottinghamshire NG90 2DB,
United Kingdom.

Reckitt Benckiser (Singapore) Pte. Ltd.
12 Marina Boulevard #19 – 01,
Marina Bay Financial Centre,
Tower 3, Singapore 018982

Product licence No.: ISIN09488P

HOW DOES NUROFEN WORK?

Nurofen, a non-steroidal anti-inflammatory drug (NSAID), provides relief by changing the body's response to pain.

WHAT IS IT FOR?

Nurofen relieves pain such as headache, migraine pain, backache, period pain, dental pain, rheumatic pain and muscular pain. Nurofen relieves fever such as fever associated with cold and flu.

WHILST TAKING NUROFEN:

CHECK BEFORE YOU TAKE THIS PRODUCT:

As with some pain relievers, do not take if you:

• experience symptoms of dengue fever (e.g. high fever, severe headache, muscle and joint pains)

- have or have ever had a stomach ulcer, perforation or
- have or nave ever had a stomach ucer, performing of bleeding
 are allergic to ibuprofen, to any of the ingredients, or to aspirin or other pain killers
 are taking other NSAID pain killers (e.g. ibuprofen, naproxen) or aspirin with a daily dose above 75mg.
 have severe kidney, liver or heart problems.
- ¹The use of ibuprofen during pregnancy should, if possible,

Please seek your doctor's advice before taking Nurofen

- Irlease seek your acctor's advice before taking Nuroten

 (Coated Tablets 200mg if you:

 have heart problems including heart failure, angina (chest
 pain), or if you have had a heart attack, bypass surgery,
 peripheral artery disease (poor circulation in the legs of feet
 due to narrow or blocked arteries), or any kind of stoke

 (including 'mini-stroke' or transient ischaemic attack "TIA")

- have or have had asthma, diabetes, high cholesterol, high blood pressure, liver, kidney or bowel problems or are dehydrated.
 have or have had history of gastrointestinal disease (e.g. ulcerative collits, Crohn's disease).
 have or have had Systemic Lupus Erythematosus (SLE) and mixed connective tissue disease.

 - are a smoker.
- blood pressure (e.g. water retention), heart disease or anticoagulants (blood thinners).

lbuprofen may be associated with a small increased risk of heart attack or stroke. Any risk is more likely with high doses (2400mg daily or 12 tablets) or prolonged treatment. Do not exceed the recommended dose and duration.

For oral administration and short-term use only.

Adults and children over 12 years: Swallow 1 or 2 tablets with water, up to three times a day as required.

Leave at least 4 hours between doses.

Do not exceed 6 tablets in 24 hours. Do not give to children under 12 years.

Problems in overdose are rare, but if you accidentally take more than the recommended dose, contact your doctor.

Side effects are rare but may include the following stomach pain, nausea, indigestion and occasionally stomach ulcer and black! tarry stools. Also worsening of asthma, itchy skin, 'nettle rash' and very rarely skin peeling and bruising.

Other side effects which have been very rarely reported are impaired kidney and liver function, headache, ringing in the ears and dizziness.

Side effects may be reduced bytaking the lowest effective dose

symptoms, stop the medication and contact your doctor or I pharmacist.

Use by the date shown on the label or carton. Store below 30°C.

KEEP ALL MEDICINE OUT OF REACH OF CHILDREN.

Taking NSAIDs at about 20 weeks of pregnancy or later may harm your unborn baby.

Date of Revision: Aug 2021

NUROFEN TABLETS AND NUROFEN CAPLETS

NUROFEN TABLETS IBUPROFEN 200mg



NUROFEN CAPLETS IBUPROFEN 200mg







