

YUCOMY SHAMPOO 2% W/W

PATIENT INFORMATION LEAFLET

What is Yucomy Shampoo 2% w/w

Yucomy Shampoo 2% w/w is a shampoo preparation containing 20mg/ml of ketoconazole and inactive ingredients sodium lauryl ether sulfate 70%, cocamidopropyl betaine, cocamide dea, isostearamidopropyl ML, acrylamide/dadmac copolymer, DM DM hydantoin, cocodimonium hydroxypropyl hydrolyzed wheat protein, disodium edetate, I – menthol powder, citric acid, select SW 10254, FD&C Red No 40, triethanolamine and purified water. It is available in plastic bottle of 120ml.

What is Yucomy Shampoo 2% w/w used for

Yucomy Shampoo 2% is used for the treatment and prophylaxis of infections in which the yeast *Pityrosporum* is involved for instances, pityriasis versicolor (localized), seborrhoeic dermatitis and pityriasis capitis (dandruff).

How to use Yucomy Shampoo 2% w/w

The affected areas of the skin or the hair should be washed with Yucomy Shampoo 2% w/w, which should be left on the skin for 3 to 5 minutes before rinsing.

For treatment:

Pityriasis versicolor: Once daily for 5 days.

Seborrhoeic dermatitis and Pityriasis capitis: Twice weekly for 3 to 4 weeks.

For prophylaxis:

Pityriasis versicolor: Once daily for 3 days during a single treatment course before exposure to sunshine.

Seborrhoeic dermatitis and Pityriasis capitis: Once every 1 to 2 weeks.

When not to use Yucomy Shampoo 2% w/w

Yucomy Shampoo 2% should be avoided if you have a known history of allergy to ketoconazole or any other listed ingredients in the product.

What are the side effects

Topical treatment with Yucomy Shampoo 2% is generally well-tolerated. As with other local shampoos, a local burning sensation, itching or contact dermatitis (due to irritation or allergy) may occur on exposed areas. Oiliness or dryness of the hair and scalp has been reported rarely.

What are the precautions

- To prevent a rebound effect after stopping a prolonged treatment with topical corticosteroids, it is recommended to continue applying the topical corticosteroid together with Yucomy Shampoo 2% and to subsequently and gradually withdraw the steroid therapy over a period of 2-3 weeks.
- Avoid contact with the eyes. If the shampoo should get into the eyes, they should be bathed with cold water.
- Prolonged use of topical ketoconazole may rarely lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical or systemic use of the medication.
- Pityriasis versicolor may cause hyper or hypopigmented patches on the body. Treatment of the infection may not result in normalization of the pigment for several months.

Usage during pregnancy and in nursing mothers

Since ketoconazole is not absorbed through the skin after topical application, pregnancy and lactation are not a contraindication for the use of Yucomy Shampoo 2%.

What are the symptoms and treatment of overdose

Not expected as Yucomy Shampoo 2% is intended for external use only. Ingestion may result in nausea and vomiting due to the detergent. In the event of accidental ingestion, only supportive measures should be carried out. In order to avoid aspiration, gastric lavage or induced emesis should not be performed.

How to keep Yucomy Shampoo 2% w/w

Keep out of the reach and sight of children.

Keep in a tight container.

Store at temperature below 30°C.

Protect from light and moisture.

The information contained in this leaflet is limited. Further information can be obtained from your doctor or pharmacist.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.

Product Registrant and Importer:
YUNG SHIN PHARMACEUTICAL (S) PTE. LTD.
10 Ubi Crescent, #06-57/58 Ubi Techpark,
Singapore 408564.

xxxxxx printing code xxxxxx

Item Code	: 204864 - 01
PM Code	: Ins.P YCL2 (S)
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: R 176 - R180 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC

L
260mm

W
110mm