

DAVMORID FILM COATED TABLET


PATIENT INFORMATION LEAFLET

What is Davmorid Film Coated Tablet

Davmorid Film Coated Tablet is a tablet containing 450mg of Diosmin and 50mg of Hesperidin and inactive ingredients Potato Starch, Sodium Starch Glycolate, Colloidal Silicon Dioxide, Ethanol 96%, Microcrystalline Cellulose, Magnesium Stearate, Purified Water, Iron Oxide Black, Lecithin, Talc, Macrogol, Titanium Dioxide, Iron Oxide Yellow, Polyvinyl Alcohol, Iron Oxide Red.

It is a vascular protector (it increases resistance in small blood vessels) and venous tonic (it increases venous tone).

Davmorid Film Coated Tablet is a light brown color film coated elliptical tablet.

“”.

It is available in blister packing of 10's x 3.

What is Davmorid Film Coated Tablet used for

Davmorid Film Coated Tablet is used for the treatment of organic and functional chronic venous insufficiency of the lower limbs with the following symptoms: swollen legs, pain and nocturnal cramps. Treatment of chronic hemorrhoids and acute hemorrhoidal attacks.

How to take Davmorid Film Coated Tablet

Usual dose in chronic conditions: 2 tablets daily in two divided doses morning and evening at mealtimes.

In acute hemorrhoidal attacks: 6 tablets daily (in 3 divided doses) for 3 to 4 days, then 4 tablets daily (in 2 divided doses).

What are the side effects

Like all medicines, Davmorid Film Coated Tablet can cause side effects, although not everybody gets them.

They may include:

Common (affects 1 to 10 users out of 100): gastrointestinal side effects (diarrhoea, dyspepsia, nausea, vomiting).

Uncommon (affects 1 to 10 users out of 1,000): Colitis (inflammation of colon).

Rare (affects 1 to 10 users out of 10,000): dizziness, headache, malaise, hypersensitivity of the skin (rash, pruritis, urticaria)

Frequency not known (cannot be estimated from available data): abdominal pain, isolated face, lip, eyelid oedema (swelling). Exceptionally Quincke's oedema (rapid swelling of tissues such as the face, lips, mouth, tongue or throat that may result in breathing difficulty).

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

What are the precautions

Acute hemorrhoidal attack:

If the hemorrhoid symptoms do not disappear within 15 days, you should ask your doctor for advice.

Venous circulation disorders:

The most effective way of giving this treatment is in combination with a healthy lifestyle. Avoid exposure to the sun, heat, excessive standing and being overweight. Walking and wearing special support stockings stimulate blood circulation.

What are the drug interactions

Tell your doctor if you are taking any other medicine or have recently taken any other medicines, including medicines obtained without a prescription.

What are the contraindications

Do not take Davmorid Film Coated Tablet if you are allergic (hypersensitive) to micronized purified flavonoid fraction or any of the other ingredients.

Usage during pregnancy and breast-feeding

As a precautionary measure, it is preferable to avoid the use of Davmorid Film Coated Tablet during pregnancy. Breast-feeding is not recommended for the duration of the treatment, due to the absence of data on the excretion of the medicine into breast milk. In general, if you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, you should always ask your doctor or pharmacist for advice before using a medication.

What are the symptoms and treatment of overdose

If you have taken more Davmorid Film Coated Tablet than you should, contact your doctor or pharmacist immediately.

The experience of overdoses with Davmorid Film Coated Tablet is limited but reported symptoms include diarrhoea, nausea, abdominal pain, pruritus and rash.

How to keep Davmorid Film Coated Tablet

Store at temperature below 30°C.

Protect from light and moisture.

Keep medicine out of reach of children.

The information contained in this leaflet is limited. Further information can be obtained from your doctor or pharmacist.

**Y. S. P.**

Manufactured by:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan,
Malaysia.

Imported by:

YUNG SHIN PHARMACEUTICAL (S) PTE. LTD.

10, Ubi Crescent, #06-57/58 Ubi Techpark,
Singapore, 408564.