

ACTAL™

Acid Neutralization

Mint-flavoured chewable tablet

DESCRIPTION

A round white mint flavoured tablets with 'ACTAL' markings on both sides.
Free from foreign matter, no chipping and capping.

COMPOSITION

Each tablet contains 360mg sodium polyhydroxyaluminium monocarbonate hexitol complex (equivalent to 216 mg of aluminium hydroxide).

EXCIPIENTS

Corn Starch, Mannitol 60, Talcum Purified Powder, Magnesium Stearate, Sodium Saccharin, Peppermint Oil, Purified Water.

INDICATIONS

For the relief of stomach discomfort such as indigestion, heartburn due to gastric hyperacidity, peptic ulcer and heartburn of pregnancy.

DOSAGE

Adult (tablet must be sucked and chewed)

Peptic ulcer: Suck and chew 2 to 4 tablets.

Gastric hyperacidity: Suck and chew 1 to 2 tablets, one-half to one hour after each meal

Heartburn of pregnancy: Suck and chew 1 to 2 tablets before breakfast and one half-hour after each meal.

ADVERSE REACTIONS

Overdose or prolonged use may on occasion cause constipation.

PRECAUTIONS

This product should not be taken concomitantly with an antibiotic containing any form of tetracycline because the antibiotic absorption may be impaired by the antacid.

WARNING

If symptoms persist please consult your doctor.

For further information please consult your doctor and pharmacist.

NATURE AND CONTENTS OF CONTAINER

The medicinal product is packaged in PTP Aluminium Foil-PVC Rigid blister packs. Each blister contains 10 tablets, packaged into cardboard boxes.

Available in boxes of 20 and 120 chewable tablets.

Not all pack sizes will be marketed.

STORAGE CONDITION

Do not store above 30°C

EXPIRY DATE

Do not use beyond expiry date.

KEEP THIS MEDICINE OUT OF THE SIGHT AND REACH OF CHILDREN.

PRODUCT REGISTRANT

A. Menarini Singapore Pte. Ltd.

30 Pasir Panjang Road, #08-32 Mapletree Business City, Singapore 117440

DATE OF LAST REVISION

07 Feb 2022

ACTAL™

酸中和

薄荷味咀嚼片

说明

圆形白色薄荷味片剂，两面均标有“ACTAL”标记。
无掺杂，无碎屑和成帽反应。

成分

每片含360毫克单羧钠聚羟铝环己醇（相当于216毫克氢氧化铝）

辅料

玉米淀粉，甘露醇60，滑石粉，硬脂酸镁，糖精钠，薄荷油，纯净水。

适应症

用于缓解消化不良、胃酸过多引起的胃灼热、消化性溃疡及妊娠期胃灼热等胃部不适。

用量

成人（必须吸吮并咀嚼药片）

消化性溃疡：吸吮和咀嚼2-4片。

胃酸过多：每餐后半小时到1小时，吸吮和咀嚼1-2片。

烧心或妊娠胃灼热：早餐前及每餐后半小时，吸吮和咀嚼1-2片。

不良反应

过量服用或长期服用可能会导致便秘。

预防措施

本品不应与含有任何形式四环素的抗生素同时服用，因为抗酸剂可能会影响抗生素的吸收。

警告

如症状持续，请咨询医生。

如需进一步信息，请咨询您的医生或药剂师。

包装性质和内容

药品采用PTP铝箔 – PVC硬质吸塑包装。每板10片，装入纸盒。

每盒有20片或120片咀嚼片。

并非所有包装类型均上市销售。

储藏条件

储藏温度不得超过30°C。

到期日

超过到期日后请勿服用。

请勿将药品放在儿童可视和可及之处。

产品注册人

A. Menarini Singapore Pte. Ltd.

30 Pasir Panjang Road, #08-32 Mapletree Business City, Singapore 117440

最后修订日期

2022 年 2 月 7 日