



**What is Covee?**

Covee Cream contains Clotrimazole 1% w/w.

It is a white, smooth, homogeneous cream. Available in pack size of 15g.

**What is this medication used for?**

Covee is used for fungal skin infections such as ringworm, athlete's foot, jock itch, white spots, fungal infections of skinfolds and erythrasma.

Covee Cream can also be used for female fungal infections of the vaginal area. It can also be used in males where inflammation is present in the penis caused by yeast fungi (*candida vulvitis and candida balanitis*)

Covee contains clotrimazole, an antifungal agent belongs to the group of medicine called imidazoles. It works by inhibiting the biosynthesis of ergosterol, a component for cell wall production; hence altering the permeability of cell membrane of fungi.

**How to use this medication?**

For topical use only.

Apply thin and evenly to the affected area two or three times daily.

To ensure complete healing, and depending on the indications, the treatment should be continued for about two weeks after the disappearance of the symptoms.

Usual treatment duration:

White spots: 1-3 weeks

Erythrasma: 2-4 weeks

Candida vulvitis and candida balanitis: 1-2 weeks

Other fungal infections: 3-4 weeks

To prevent reoccurrence of fungal infections of the feet, treatment should be continued for about 2 weeks beyond the disappearance of all symptoms.

After washing, the feet should be thoroughly dried (spaces between the toes) before application of the cream.

You should notify your physician if there is no improvement after the recommended duration of treatment.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

**When should you not take this medicine?**

Do not use Covee if you have known hypersensitivity to any ingredients in this medicine.

**Care that should be taken when taking this medicine?**

Tell your doctor or pharmacist if:

- you are pregnant or plan to get pregnant.
- you are breast-feeding

when used for treatment of fungal infection.

Covee may reduce the effectiveness and safety of latex products such as condoms and diaphragms when applied on the genital area. The effect is temporary and occurs only during treatment.

Keep medicine out of the sight and reach of children.

Avoid contact with eyes. Do not swallow.

Cetostearyl alcohol may cause local skin reactions (e.g. contact dermatitis).

**Interactions with other medications**

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. Medicine interactions can change the way a medicine works. This may decrease or increase the effects of the medicine and it may also lead to other harmful effects.

Do inform your doctor and/or pharmacist if you are taking, have recently taken or plan to take other medications, nutritional supplements or herbal products.

**Side effects**

Along with its desired effects, a medicine may cause some unwanted effects. Tell your pharmacist or doctor as soon as possible if you do not feel well while you are taking Covee.

Common side effects are irritation, burning sensation, itching, redness.

Like all medicines, this product can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

**When to seek further medical advice**

Inform your doctor and/or pharmacist immediately if you experience any allergic symptoms or symptoms of side effects.

Consult your doctor if the symptoms do not get better, worsen while taking this medicine.

**What should you do if you miss a dose?**

If you miss a dose, take it as soon as possible. However, if it is almost time for next dose, skip the missed dose and back to your regular dosing schedule. Do not take double dose to make up for the dose that you have missed.

**What to do when you have taken more than the recommended dosage?**

Risk of overdose symptoms is unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favorable to absorption) or accidental oral ingestion. In the case of overdose, there is no specific antidote.

**How should this medication be kept?**

Keep this medicine in its original container until use.

Store below 30°C.

**List of excipients:** Macrogol Cetostearyl Ether, Cetostearyl Alcohol, Liquid Paraffin, Propylene Glycol, Sodium Citrate, Citric Acid Anhydrous, Methyl Hydroxybenzoate, Propyl Hydroxybenzoate and Purified Water.

Revision date: 30-Sep-2021

Manufactured by

**Xepa-Soul Pattinson (Malaysia) Sdn Bhd**

1-5 Cheng Industrial Estate, 75250, Melaka, Malaysia.

Product Registrant

**Apex Pharma Marketing Pte Ltd**