

# FLUCUR Nebuliser

## LOCAL APPLICATION FOR NASAL CONGESTION AND DISCHARGE CAUSED BY ALLERGIC RHINITIS OR OTHER NASAL SYMPTOMS

In many cases, rhinorrhea and/or rhinostenosis are the symptoms which reflect swelling or hyperemia of the nasal mucosa attributable to common cold virus, pollen or house dust. FLUCUR Nebuliser is available as a spray-type preparation. Spraying this drug into the nasal cavity allows its active ingredient to spread uniformly, thereby alleviating uncomfortable symptoms and leading to smooth nasal respiration.

### [COMPOSITION]

100ml of FLUCUR Nebuliser contains:

Naphazoline Hydrochloride .....50mg      Boric Acid (Buffer) .....1700mg  
Chlorpheniramine Maleate .....300mg      Sodium Borate (pH adjusting agent) .....appropriate amount

### [DOSAGE AND ADMINISTRATION]

Adults and children aged 7 years & above:

Spray into each nasal cavity once or twice each time. Application should be up to 5 times per day at intervals of at least 3 hours.

(How to Apply FLUCUR Nebuliser)

1. Remove the cap and hold the nebuliser vertically. Insert the tip of nebuliser slightly into the nostril.

2. Press the body of nebuliser so that the solution is sprayed into the nostril.

3. Take out the tip of nebuliser from the nostril and release the nebuliser.

4. Wipe the tip of nebuliser with sterilized gauze after use to keep the nebuliser clean.

\* There is an appropriate air space left in the nebuliser for the effective operation, however, the quantified 14ml of the solution is accurately filled in the nebuliser.

### [INDICATIONS]

Alleviation of symptoms caused by acute rhinitis, allergic rhinitis or paranasal sinusitis (sneezing, runny nose, nasal congestion)

### [PRECAUTIONS]

1. The following persons should not use FLUCUR Nebuliser.

1) Children aged below 7 years.

2. The following persons should consult the physician or pharmacist before use.

1) Patients with hypertension, cardiac disease, diabetes, hyperthyroidism, glaucoma etc.

2) Patients with a history of allergic reaction to nose drops (such as eruption, redness, swelling, feeling of irritation etc.).

3) Pregnant women or women who may be pregnant.

4) Patients under treatment by a physician.

3. The following cautions should be exercised in usage.

1) Since the excessive use of FLUCUR Nebuliser may induce the rebound of nasal congestion or secondary hyperemia, the indicated dosage and administration should be strictly adhered.

2) Children should be allowed to use FLUCUR Nebuliser only under the instruction and supervision of guardian.

3) FLUCUR Nebuliser is only for application to nasal cavity, and should not be used to eyes or ears.

4. The following cautions should be exercised during and after use.

1) Discontinue the application when symptoms such as eruption, redness, swelling, feeling of irritation etc. appear, and immediately consult the physician or pharmacist.

2) In case no improvement of symptoms is shown after administration for about 3 days, discontinue the application and consult the physician or pharmacist.

3) FLUCUR Nebuliser should not be used continuously for long term.

4) FLUCUR Nebuliser should not be used in common in order to avoid contamination.

5. Precautions in storage and handling

1) Keep out of reach of children.

2) Store away from direct sunlight, in a cool place, tightly capped.

3) To avoid misuse and maintain the quality, FLUCUR Nebuliser should not be transferred to another nebuliser.

4) FLUCUR Nebuliser should not be used after the indicated expiry date.

*Sole Agents*

### LUEN WAH MEDICAL CO., (S)PTE. LTD.

40, North Canal Road SINGAPORE 059296, Phone:65-5339141-6 Fax:65-5332238

### LUEN WAH MEDICAL CO. SDN. BHD.

No. 37G, 37-1F, 39-1F, The Highway Centre, Jalan 51/205, 46050 Petaling Jaya, Selangor Darul Ehsan, MALAYSIA.  
Phone:603-7955510 Fax:603-7955520

### KIN FUNG WEISEN-U CO., LTD.

Unit 715, 7/F, Block B, Seaview Estate, 2-8 Watson Road, North Point, HONG KONG

Phone:852-25444182-5 Fax:852-28158500

Manufactured by

### SHIGAKEN PHARM. IND. CO., LTD.

879 Taki, Koka-cho, Koka-gun, Shiga, JAPAN

SP

在很多种情况下，感冒病毒、花粉或室内尘埃，都会使鼻腔中的黏膜红肿充血或充满鼻液，导致流涕及鼻道狭窄等症。『呼佳』喷务剂即以喷雾方式制成，用此药物向鼻腔喷射可将其主要成份均匀散开，减轻上述病症，并使鼻腔呼吸恢复顺畅。

每百毫升(CU)之「呼佳」含有

盐酸茶碱	50 毫克
马来酸氯苯那敏·扑尔敏	300 毫克
明胶(缓冲剂)	1700 毫克
硬脂酸钠(平日调节剂)	适量

成人或七岁以上之儿童均可使用。每隔三小时在鼻腔内喷射一至二次, 每天最多使用五次。


一、转开瓶盖(参阅插图一),将头部略向前俯,把瓶端伸入鼻腔(插图二)。

- 二、用手挤压眼眶中部使药液喷向鼻腔。  
三、将喷雾剂自鼻腔中喷出,手指放松。  
四、用药棉将眼眶抹干,并将眼眶套套紧,保持清洁。
- (注:眼眶内保留部分空间,挤压时可增强喷射力。药瓶注明容量十四毫升乃药液之实际容量。)

会迅速减轻鼻涕直流，鼻腔敏感，或其他并发鼻腔病征。（如打喷嚏，流鼻涕，鼻腔阻塞等）。

一、下列人士不宜使用

- [illegible]



滋賀県製薬株式会社製造

日本・滋賀县甲賀郡甲賀町 八七九

香港总代理 建丰胃仙U有限公司

香港北角屈臣道二至八号海景大厦 B 座七楼七一五室

电话: 八五二二一五四四四一八二一五  
传真: 八五二二一五八五〇〇

圖 10-2-10 半圓形

新加坡联华药行(私人)有限公司

新加坡〇五九二九六郵區怒干那律四〇號

電話: 六五二五三三九一 傳真: 六五二五三三三九

马来西亚联华药行有限公司

马来西亚雪兰莪州八打灵再也西六〇五〇邮区五一—一二〇五路  
大道中心门牌三七G、三七—F及三九—F

电话:六〇三-七九五五五二〇 传真:六〇三-七九五五五二〇