

PATIENT INFORMATION LEAFLET

IFIMOL*500

Film-coated Tablet 500mg

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or ask your doctor or pharmacist.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

Description of Product

Each Ifimol*500 film coated pink coloured capsule shaped film coated tablets having score line on both the sides of the tablet contains:

Paracetamol BP.....500mg

The tablet can be divided into equal halves.

Excipient: Maize Starch, Talc, Magnesium Stearate, Colloidal Anhydrous Silica, Sodium Starch Glycolate, and Colour Opadry pink II 85 G 84925 (Lecithin Soya, Polyvinyl Alcohol part hydrolyzed, Ponceau 4R Aluminium Lake, Talc, Titanium Dioxide, Macrogol/PEG).

INDICATIONS

IFIMOL*500 has analgesic and antipyretic effect and can be used for relief of fever and painful disorders such as headache, migraine, backache, joint and muscle pain, period pain (dysmenorrhea), toothache and sore throat.

DOSAGE AND ADMINISTRATION

For oral administration only. Do not exceed the stated dose. Always use the lowest effective dose to relieve your symptoms. Do not take more frequently than every 4 hours. Do not use this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu.

Adults (including the elderly) and children over 12 years: 1 - 2 tablets, taken every 4 to 6 hours as required. Do not take more than 8 tablets in one day.

Children aged 6 to 12 years: Half to one tablet, taken every 4 to 6 hours as required. Do not take more than four doses in one day. Do not take for more than 3 days without asking your doctors.

Children under 6 years: Not recommended for children under the age of 6 years.

CONTRAINDICATION

Do not use if you are allergic to Paracetamol or any of the other ingredients in the product

ADVERSE REACTIONS

Stop taking this medicine and tell your doctor immediately if:

- You experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- You experience any skin rash or reaction. Symptoms of serious skin reactions may include skin reddening, rash, blisters, and separation of the upper surface of the skin from the lower layers. These serious skin reactions are rare, but can occur even if you have taken this medicine in the past without any problems.
- You have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product.
- You experience nausea, sudden weight loss, loss of appetite and yellowing of the eyes and skin.
- You experience unexplained bruising or bleeding.
- You experience mouth ulcers.

These reactions are rare.

DRUG INTERACTIONS

Before taking this medicine, make sure you consult your doctor if you are taking Warfarin or similar medicines used to thin the blood.

WARNINGS AND PRECAUTIONS

Taking too much paracetamol can cause serious harm to your liver. Always read and follow the label.

Check with your doctor before use if:

- You have liver or kidney problems.
- You are underweight or malnourished.
- You regularly drink alcohol.

You may need to avoid using this product altogether or limit the amount of paracetamol that you take.

- You have a severe infection as this may increase the risk of metabolic acidosis.

Signs of metabolic acidosis include:

- deep, rapid, difficult breathing
- feeling sick (nausea), being sick (vomiting)
- loss of appetite

Contact a doctor immediately if you get a combination of these symptoms. If you have had a serious skin reaction with this medicine, do not take it or any products containing paracetamol or acetaminophen again. Doing so could cause you to have another serious skin reaction.

Avoid taking alcohol with this medicine. Please see your doctor if your symptoms do not improve. Keep out of sight and reach of children.

OVERDOSE

If you take more of the medicine than you should: seek medical advice immediately even if you feel well because of the risk of liver failure.

For further information, please refer to your doctor or pharmacist.

PACKS

Box of 10 strips x 10's and box of 100 strips x 10's

STORAGE CONDITION

Store below 30°C. Protect from light.

EXPIRY

Do not use the medication beyond expiry date.

Name of manufacturer

Unique Pharmaceutical Laboratories, India
(A Division of J. B. Chemicals & Pharmaceuticals Limited)