

652348
the first time include headache, nausea, vomiting, tingling/ burning sensations, inflammation or pain in your mouth or lips, changes in taste perception, changes in saliva production, dryness of the mouth, indigestion and hiccups. These effects may be due to incorrect use of NICORETTE® QuickMist like using too much, or swallowing too much nicotine.

Other commonly reported side-effects include dizziness, vomiting, sore mouth, throat irritation, dry mouth, taste disturbance, tingling or numbness of hands or feet, stomach pain or discomfort, diarrhoea, wind, cough, burning sensation, fatigue, hypersensitivity.

Uncommonly reported side-effects include chest palpitations, fast heartbeat, flushing, high blood pressure, redness or itching of the skin, hives, wheezing, shallow breathing, difficulty speaking, nasal congestion, sneezing, tight throat, burping, red tongue, mouth blisters, mouth tingling or numbness, excessive sweating, limb weakness, chest discomfort or pain, general feeling of being unwell, abnormal dreams.

Very rarely, abnormal beating of the heart, swelling of the face, lips, mouth or tongue, difficulty swallowing, blurred vision, watery eyes, dry throat lip pain, retching, muscle tightness and seizure have been reported.

When you stop smoking you may also develop mouth ulcers. The reason why this happens is unknown.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**5 HOW TO STORE
NICORETTE® QUICKMIST**

Keep out of sight and reach of children. In the event of overdose, get medical help right away.

Do not use Nicorette® QuickMist after the expiry date which is stated on the dispenser and on the blister card. The expiry date refers to the last day of that month unless stated otherwise.

Store at temperatures not exceeding 30°C

Medicines should not be disposed of via waste-water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**6 FURTHER
INFORMATION**

What Nicorette® QuickMist contains

- The active substance is nicotine. One spray dose contains 1 mg of nicotine.
- The other ingredients are: propylene glycol, anhydrous ethanol, trometamol, poloxamer 407, butylated hydroxytoluene (BHT), glycerol, sodium hydrogen carbonate, levomenthol, mint flavour, cooling flavour, sucralose, acesulfame potassium, hydrochloric acid and purified water.

What Nicorette® QuickMist looks like and contents of the pack

Nicorette® QuickMist consists of a plastic bottle of solution held in a dispenser with a mechanical spray pump. The dispenser has a child resistant feature.

Each bottle contains 13.2 ml of solution which provides approximately 150 sprays, each containing 1mg nicotine.

Nicorette® QuickMist is supplied in packs of either 1 or 2 dispensers.

Not all pack sizes may be marketed.

Product Registration Holder:
Johnson & Johnson (Thailand) Ltd.
Bangkok, Thailand.
Reg.No 1C 15008/66 (N)
Tel.1800-333-666 (Toll free except hand phone)

Johnson & Johnson Pte. Ltd.
2 Science Park Drive #07-13
Ascent, Singapore Science Park 1
Singapore 118222
SIN16567P.
Singapore consumer hotline: 800-6011-460

Manufacturer:
McNeil AB, P.O. Box 941, Norrbroplatsen 2,
SE-251 09 Helsingborg, Sweden