140 mm

140 mm

Difflam® FORTE ANTI-INFLAMMATORY THROAT SPRAY White metered dose pump with a green lid containing 15mL of clear solution What Difflam® Forte Anti-inflammatory Throat Spray contains Active ingredient Each mL of Difflam® Forte Anti-inflammatory Throat Spray contains benzydamine hydrochloride 3.0mg. Each spray delivers approximately 0.5mg of benzydamine hydrochloride It also contains ethanol, saccharin sodium, methylparahydroxybenzoate, glycerol, PEG-40 hydrogenated castor oil, purified water and mint flavouring. It does not contain gluten, lactose, sugar or colour. How does Difflam® Forte Anti-inflammatory Throat Spray work Difflam® Forte Anti-inflammatory Throat Spray contains benzydamine hydrochloride. It is an anti-inflammatory and analgesic agent. It helps to reduce swelling and pain. What Difflam® Forte Anti-inflammatory Throat Spray is used for Difflam® Forte Anti-inflammatory Throat Spray is used for the temporary relief of painful conditions of the mouth and throat including tonsillitis, sore throat, radiation mucositis, aphthous ulcers, pharyngitis, swelling, redness, inflammatory conditions, post orosurgical and periodontal procedures. How to use Difflam® Forte Anti-inflammatory Throat Spray Adults and children over 12 years: 2-4 sprays (1-2 mg) directly onto the sore/inflamed area and swallow gently. Repeat every 1½ to 3 hours as necessary 2 sprays (1 mg) directly onto sore/inflamed area and swallow gently. Repeat every 11/2 to 3 hours as Children under 6 years: Uninterrupted treatment should not exceed seven days, unless under medical supervision After use, wipe the nozzle with a clean tissue to prevent blockage. Contraindications Patients with known hypersensitivity to benzydamine hydrochloride or to any other ingredients present in Difflam® Forte Anti-inflammatory Throat Spray. If a sore throat is either caused or complicated by a bacterial infection, appropriate anti-bacterial therapy should be considered in addition to the use of Difflam® Forte Anti-inflammatory Throat Spray. Before using Difflam® Forte Anti-inflammatory Throat Spray Tell your doctor or pharmacist if you have any of the following conditions liver problems kidney problems

Difflam® Forte Anti-inflammatory Throat Spray is generally well tolerated and side effects are minor.

The most commonly reported reaction is oral numbness and has been reported in 2.6% of treated

____ TEXT AREA (116 mm x 186 mm)

TEXT AREA (116 mm x 186 mm) cases. Occasional burning or stinging sensation may occur (1.4%). Other local adverse effects are less common and included dryness or thirst (0.2%), tingling (0.2%), warm feeling in mouth and altered sense of taste (0.1%).

Systemic side effects are rare. These include nausea, vomiting, retching, gastro-intestinal disorders (0.4%), dizziness (0.1%), headache and drowsiness (0.1%). Hypersensitivity reactions occur very rarely but may be associated with rash, urticaria, photodermatitis and occasionally laryngospasm or bronchespasm.

Use in pregnancy

The safety of benzydamine hydrochloride has not been established in pregnant patients. Seek your doctor's advice if you are pregnant or likely to be pregnant during your course of medication.

Use during breastfeeding

Use of benzydamine hydrochloride during lactation has not been established in humans. Seek your doctor's advice if you are breastfeeding or likely to breastfeed during your course of medication.

Use in children

Because of the lack of sufficient clinical experience, Difflam® Forte Anti-inflammatory Throat Spray is not recommended in children under 6 years of age.

Interactions with other medicines

There are no known drug interactions with Difflam® Forte Anti-inflammatory Throat Spray.

Overdose

There are no known cases of overdosage with Difilam® Forte Anti-inflammatory Throat Spray. Adverse CNS effects have been reported following overdosage with high doses of benzydamine hydrochloride in solution form. There is no specific antidote for benzydamine hydrochloride and should excessive quantities be ingested, the treatment should be symptomatic.

Storage

Protect from light and store below 30°C.

inova

Revision Date: October 2019

A2969



RE-TYPESET ARTWORK. Due to the nature of the files received, this artwork has been re-typeset.

