

UK

Lonquex pre-filled syringe

6mg/0.6ml

Lipegfilgrastim

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

- What is in this leaflet
- What Lonquex is and what it is used for
 - What you need to know before you use Lonquex
 - How to use Lonquex
 - Possible side effects
 - How to store Lonquex
 - Contents of the pack and other information
 - Information for injecting yourself

1 WHAT LONQUEX IS AND WHAT IT IS USED FOR

What Lonquex is

Lonquex contains the active substance lipegfilgrastim. Lipegfilgrastim is a long-acting modified protein produced by biotechnology in bacteria called *Escherichia coli*. It belongs to a group of proteins called cytokines and is similar to a natural protein (granulocyte-colony stimulating factor [G-CSF]) produced by your own body.

What Lonquex is used for

Your doctor has prescribed Lonquex for you in order to reduce the duration of a condition called neutropenia (low white blood cell count) and the occurrence of febrile neutropenia (low white blood cell count with a fever). These can be caused by the use of cytotoxic chemotherapy (medicines that destroy rapidly-growing cells).

How Lonquex works

Lipegfilgrastim stimulates the bone marrow (the tissue where new blood cells are made) to produce more white blood cells. White blood cells are important as they help your body fight infection. These cells are very sensitive to the effects of chemotherapy which can cause the number of these cells in your body to decrease. If white blood cells fall to a low level, there may not be enough left in the body to fight bacteria and you may have an increased risk of infection.

2 WHAT YOU NEED TO KNOW BEFORE YOU USE LONQUEX

Do not use Lonquex:

- if you are allergic to lipegfilgrastim or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before using Lonquex

- if you get left upper abdominal pain or pain at the tip of your shoulder. It could be a consequence of a spleen disorder (see section "4. Possible side effects").
- if you have a cough, fever and difficulty breathing. It could be a consequence of a pulmonary disorder (see section "4. Possible side effects").
- if you have sickle cell anaemia, which is an inherited disease characterised by sickle-shaped red blood cells.
- if you have previously experienced allergic reactions to other medicines like this one (e.g. filgrastim, lenograstim or pegfilgrastim of the group of G-CSFs). There could be a risk of reacting to Lonquex too. Your doctor will carry out regular blood tests in order to monitor various blood components and their levels.

Children and adolescents

Do not give this medicine to children and adolescents under 18 years of age because there are no data that show that this medicine is safe and works in this age group.

Other medicines and Lonquex

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines.

You will normally inject your dose of Lonquex approximately 24 hours after your last dose of chemotherapy at the end of each chemotherapy cycle.

Pregnancy and breast-feeding

Lonquex has not been tested in pregnant women. It is important to tell your doctor if you are pregnant, think you may be pregnant or are planning to have a baby, as the doctor may decide that you should not use this medicine.

It is unknown whether the active substance in this medicine passes into the breast milk. You should therefore interrupt breast feeding during treatment.

Driving and using machines

Lonquex has no or negligible influence on your ability to drive and use machines.

Lonquex contains sorbitol and sodium

This medicine contains sorbitol. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before using this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per pre-filled syringe, i.e. essentially 'sodium-free'.

3 HOW TO USE LONQUEX

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

What the recommended dose is

The recommended dose is one pre-filled syringe (6 mg lipegfilgrastim) **once per chemotherapy cycle**.

When to use Lonquex

This medicine should be given approximately 24 hours after your last dose of chemotherapy at the end of each chemotherapy cycle.

How are the injections given?

This medicine is given as an injection using a pre-filled syringe. The injection is given into the tissue just under the skin (subcutaneous injection).

Your doctor may suggest that you learn how to inject yourself with this medicine. Your doctor or nurse will give you instructions on how to do this. Do not attempt to give Lonquex to yourself without this training. Information required for using the pre-filled syringe can be found at the end of this package leaflet (see section "7. Information for injecting yourself"). Proper treatment of your disease, however, requires close and constant co-operation with your doctor.

If you use more Lonquex than you should

If you use more Lonquex than you should, talk to your doctor.

If you forget to use Lonquex

If you have missed an injection, contact your doctor to discuss when you should inject the next dose.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4 POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Most serious side effects

- Allergic reactions such as skin rash, raised itchy areas of skin and serious allergic reactions with weakness, drop in blood pressure, difficulty breathing and swelling of the face have been reported uncommonly (may affect up to 1 in 100 people). If you think you are having this type of reaction, you must stop your Lonquex injection and get medical help immediately.
- Increased spleen size and cases of spleen ruptures have been reported with other medicines similar to Lonquex. Some cases of splenic rupture were fatal. It is important to contact your doctor immediately if you experience **pain in the upper left side of the abdomen or left shoulder pain** since this may relate to a problem with your spleen.

- Cough, fever and difficult or painful breathing can be signs of uncommon (may affect up to 1 in 100 people) serious pulmonary side effects, such as pneumonia and acute respiratory distress syndrome, which may be fatal. If you have a fever or any of these symptoms, it is important to contact your doctor immediately.

Other side effects

Very common (may affect more than 1 in 10 people)

- Musculoskeletal pains such as bone pain and pain in the joints, muscles, limbs, chest, neck or back. Tell your doctor if you experience severe musculoskeletal pain.
- Nausea.

Common (may affect up to 1 in 10 people)

- Reduction in blood platelets, which increases risk of bleeding or bruising.
- Headache.
- Skin reactions, such as redness or rash.
- Low blood levels of potassium, which can cause muscle weakness, twitching or abnormal heart rhythm.
- Chest pain.

Uncommon (may affect up to 1 in 100 people)

- Rise in white blood cells.
- Local reactions at the injection site, such as pain or hardening.
- Some changes may occur in your blood, but these will be detected by routine blood tests.

Side effects that have been seen with similar medicines, but not yet with Lonquex

- Sickle cell crises in patients with sickle cell anaemia.
- Plum-coloured raised painful sores on the limbs and sometimes the face and neck with fever (Sweet's syndrome).
- Inflammation of the blood vessels in the skin.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

By reporting side effects you can help provide more information on the safety of this medicine.

5 HOW TO STORE LONQUEX

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the outer carton and on the label of the pre-filled syringe after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2 °C – 8 °C).

Do not freeze.

Keep the pre-filled syringe in the outer carton, in order to protect from light.

Lonquex may be removed from the refrigerator and stored below 25 °C for a maximum single period of up to 3 days. Once removed from the refrigerator, the medicine must be used within this period or disposed of. Do not use this medicine if you notice that it is cloudy or there are particles in it.

Dispose of this medicine as instructed by your doctor, nurse or pharmacist.

6 CONTENTS OF THE PACK AND OTHER INFORMATION

What Lonquex contains

- The active substance is lipegfilgrastim. Each pre-filled syringe contains 6 mg lipegfilgrastim. Each ml solution contains 10 mg lipegfilgrastim.
- The other ingredients (excipients) are glacial acetic acid, sodium hydroxide, sorbitol (E420), polysorbate 20 and water for injections.

What Lonquex looks like and contents of the pack

Lonquex is a solution for injection (injection) in pre-filled syringe with a fixed injection needle in a blister. Lonquex is a clear and colourless solution. Each pre-filled syringe contains 0.6 ml solution.

Each pack contains 1 pre filled syringe with or without safety device.

Not all pack sizes may be marketed.

Manufacturer

TEVA Pharmaceutical Industries, Ltd
18, Eli Hurvitz St., Industrial Zone
Kfar Saba, 4410202, Israel

7 INFORMATION FOR INJECTING YOURSELF

This section contains information on how to give yourself an injection of Lonquex under the skin. It is important that you do not try to give yourself the injection unless you have received special training from your doctor or nurse. If you are not sure about giving yourself the injection or you have any questions, please ask your doctor or nurse for help.

How Lonquex is used

You will need to give yourself the injection into the tissue just under the skin. This is known as a subcutaneous injection.

Equipment that you need

To give yourself an injection into the tissue under the skin you will need:

- a pre-filled syringe of Lonquex,
- an alcohol wipe,
- a piece of gauze bandage or a sterile gauze swab,
- a puncture-proof container (plastic container provided by the hospital or pharmacy) so you can dispose of used syringes safely.

Teva Artwork Department | Harlow

Job Number 63252

PRODUCT

Lonquex 6mg Sol for Inj pre-filled syringe 1 (P) Sp SG (Ulm)

Component Leaflet

Dimensions 640x215mm

SAP No. N/A

COLOURS

Cutterguide ☒

Black ☒

FONTS

Neo Sans for Teva

Body Size 11pt

Planner Kerry

SOFTWARE USED

Adobe InDesign CC

ARTWORK

FIRST DRAFT

Artworker SC

Date 9-6-20

AMENDS


Draft 3.0

Artworker IG


Date 16-10-20

APPROVAL

ARTWORKER


Iain Greengrass
9:07 am, 16-10-2020

PLANNER


10:33 am, 16-10-2020

Supplier Instructions:

Artwork, text and content must NOT be altered. The only exceptions to this are: bleeds, chokes, spreads or other adjustments required for print reproduction purposes only. If you have any difficulties please contact the Teva Artwork Team. We must receive a copy of the 3rd Party Vendors Proof before final approval can be made.

Viewing Instructions:


To ensure this file views correctly, please set overprint preview to 'ALWAYS' in your Adobe Acrobat preferences.
Acrobat → Preferences → Page Display → Use Overprint Preview → Always
When overprint preview is set up correctly there should be no 'X' appearing in any box marked A-D. Please follow the instructions above if you see an 'X'.

JB V 12.indd

How to prepare for your injection

Before you give yourself an Lonquex injection, you must do the following:

1. Hold the syringe and gently remove the cover from the needle without twisting. Pull straight as shown in pictures 1 and 2. Do not touch the needle or push the plunger.
2. You may notice small air bubbles in the pre-filled syringe. If there are air bubbles present, gently tap the syringe with your fingers until the air bubbles rise to the top of the syringe. With the syringe pointing upwards, expel all air from the syringe by pushing the plunger slowly upwards.
3. You can now use the pre-filled syringe.




Where your injection should be given

The most suitable places to inject yourself are:


- the top of your thighs,
- the abdomen (see grey areas in picture 3) avoiding the skin directly surrounding the navel.

If someone else is injecting you, they can also use the back and side of your upper arms (see grey areas in picture 4 and 5).



How you should inject yourself

1. Disinfect the injection site on the skin by using an alcohol wipe and pinch the skin between your thumb and forefinger, without squeezing it (see picture 6).
2. Put the needle fully into the skin as shown by your nurse or doctor. The angle between the syringe and skin should not be too narrow (at least 45°, see picture 7).
3. Pull slightly on the plunger to check that a blood vessel has not been punctured. If you see blood in the syringe, remove the needle and re-insert it in another place.
4. Inject the liquid into the tissue slowly and evenly, always keeping your skin pinched.
5. After injecting the liquid, remove the needle and let go of your skin.
6. Press the injection site with a piece of gauze bandage or a sterile gauze swab for several seconds.
7. Only use each syringe for one injection. Do not use any Lonquex that is left in the syringe.



Remember

If you have any problems, please ask your doctor or nurse for help and advice.

Disposing of used syringes

- Do not put the cover back on used needles.
- Put used syringes into the puncture-proof container and keep this container out of the sight and reach of children.
- Dispose of the full puncture-proof container as instructed by your doctor, nurse or pharmacist.
- Never put the syringes that you have used into your normal household rubbish bin.

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

This leaflet was last revised in October 2020.