

98 mm

# VERAGEL-DMS TABLET

## Indications:

For the management and treatment of hyperacidity, gastritis and peptic ulcer.

## Method of administration:

Oral administration.

## Dosage/ Times of administration:

1 to 2 tablets four times a day, to be taken 1 hour after meals and at bedtime or, as recommended by a doctor.

## Precautions:

Aluminum hydroxide may cause constipation. This preparation should not be given when antibiotic is presently being taken.

## Side Effects:

Constipation may result due to the Aluminum hydroxide or at times diarrhea due to the magnesium carbonate component. In either case, temporarily discontinue the drug for a couple of days then resume by taking half of the recommended dose.

## Storage and disposal:

Medicine should not be exposed to sunlight and should be stored at temperature not exceeding 30°C.

## Formulation:

Each tablet contains

Aluminum Hydroxide-Magnesium	- 325 mg
Carbonate Co-precipitate	- 10 mg
Dimethylpolysiloxane	- 10 mg
Dicyclomine HCl	- 2.5 mg

There are no potentially sensitizing inactive ingredients in the formulation.

**This leaflet contains limited information on the preparation. Please consult a doctor or a pharmacist for further information.**

Manufactured by  
**UNILAB, Inc.**

No. 66 United Street, Mandaluyong City,  
Metro Manila, Philippines

Imported by:

Goldplus Universal Pte Ltd  
103 Kallang Avenue #06-02, Singapore 339504.



P300000033030

140 mm