

Reckitt
Benckiser
 Manufacturer: Reckitt Benckiser Healthcare International Limited
 Thane Road, Nottingham, NG90 2DB, United Kingdom
 MAL10021806X HK-596774
 For/Imported by: Reckitt Benckiser (M) Sdn. Bhd. (175141-H), Menara UIC, Mutiara Damansara, Selangor, 47800, Malaysia.
 Reckitt Benckiser (S) Pte. Ltd. (1397702628K), Guthrie House, Singapore 266802.
 Reckitt Benckiser (HK) Ltd. (072739660), Hong Kong. 032310 02752575

Drug Facts

Active ingredient (in each extended-release tablet)
Guaifenesin 600 mg.....Expectorant

Uses

- helps loosen phlegm and thin bronchial secretions
- symptomatic relief of deep chesty coughs
- expectorant for productive cough

有效化解濃痰及支氣管分泌物
舒緩咳嗽不適症狀
幫助咳出痰沫

Warnings

Warnings
Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
In case of overdose, get medical help or contact a Poison Control Center right away.

Possible side effects

- Possible side effects**
 ■ nausea ■ abdominal discomfort ■ vomiting
 Tell your doctor if you have any side effects
 after taking this medicine, including any not
 listed here. ►

Drug Facts (continued)

Directions

- do not crush, chew, or break tablet
 - swallow tablet whole with a full glass of water
 - adults and children 12 years of age and over:
 - 1 or 2 tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
 - children under 12 years of age: do not use
 - if allergic to any of the ingredients: do not use
- 服用方法:**
 用水吞服。請勿壓碎，咀嚼或碾破藥片。
 只供口服。

劑量：
十二歲或以上青少年及成人，每十二小時一至二片。二十四小時內，不可超過四片。
十二歲以下兒童不宜服用。

Other information

- tamper evident: do not use if carton is open or if printed seal on blister is broken or missing
- do not store above 30°C

請勿儲存於30°C以上。

**Keep out of reach of children.
Jauhi dari kanak-kanak.**

Inactive ingredients

Inactive ingredients: carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

MucinexTM
Extended-release tablets 600 mg
Guaifenesin 600 mg
EXPECTORANT

Unique Bi-Layer Tablet

Tablet shown actual size

Immediate-Release Layer

Extended-Release Layer

A diagram of a bi-layer tablet. The tablet is shown in a perspective view, revealing two distinct layers. The top layer is white and labeled 'Immediate-Release Layer' with a downward arrow. The bottom layer is blue and labeled 'Extended-Release Layer' with an upward arrow. The entire tablet is labeled 'Unique Bi-Layer Tablet' at the top and 'Tablet shown actual size' to its left. A small 'TM' symbol is visible on the bottom right of the blue layer.

Mucinex™
Extended-release tablets 600 mg
Guaifenesin 600 mg
6 EXTENDED-RELEASE TABLETS

Mucinex™
美清痰™

**Extended-release tablets 600 mg
Guaifenesin 600 mg**

EXPECTORANT 化痰藥片



- ✓ **Thins And Loosens Mucus** ✓ **有效化痰**
- ✓ **Relieves Chest Congestion** ✓ **舒緩積痰咳嗽**

6 EXTENDED-RELEASE TABLETS



Reckitt Benckiser Manufacturer: Reckitt Benckiser Healthcare International Limited
Thane Road, Nottingham, NG90 2DB, United Kingdom
MAL10021806X HK-59774
For Import by: Reckitt Benckiser (M) Sdn. Bhd. (175141-H), Menara UAC,
Mutiara Damansara, Selangor 47800, Malaysia
Reckitt Benckiser (S) Pte. Ltd. (197202628X) Guthrie House, Singapore 268802.
032310 0273787
Reckitt Benckiser (HK) Ltd. (07279860), Hong Kong.

Drug Facts

Active ingredient (in each extended-release tablet)
Guaifenesin 600 mg.....Expectorant

Uses

- helps loosen phlegm and thin bronchial secretions
- symptomatic relief of deep chesty coughs
- expectorant for productive cough

有效化解濃痰及支氣管分泌物

紓緩咳嗽不適症狀

幫助咳出痰沫

Warnings

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
In case of overdose, get medical help or contact a Poison Control Center right away.

Possible side effects

- nausea
 - abdominal discomfort
 - vomiting
- Tell your doctor if you have any side effects after taking this medicine, including any not listed here.

Drug Facts (continued)

Directions

- do not crush, chew, or break tablet
 - swallow tablet whole with a full glass of water
 - adults and children 12 years of age and over: 1 or 2 tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
 - children under 12 years of age: do not use
 - if allergic to any of the ingredients: do not use
- 服用方法:**
用水吞服。請勿壓碎，咀嚼或砸破藥片。只供口服。

劑量:

十二歲或以上青少年及成人，每十二小時一至二片。二十四小時內，不可超過四片。十二歲以下兒童不宜服用。


Other information

- tamper evident: do not use if carton is open or if printed seal on blister is broken or missing
- do not store above 30°C

請勿儲存於 30°C 以上。

Keep out of reach of children.
Jauh dari kanak-kanak.

Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF



Mucinex™
Extended-release tablets 600 mg
Guaifenesin 600 mg
20 EXTENDED-RELEASE TABLETS

Mucinex™
美清痰™
Extended-release tablets 600 mg
Guaifenesin 600 mg
EXPECTORANT 化痰藥片

✓ Thins And Loosens Mucus
✓ Relieves Chest Congestion
20 EXTENDED-RELEASE TABLETS

✓ 有效化痰
✓ 紓緩積痰咳嗽



Mucinex™
Extended-release tablets 600 mg
Guaifenesin 600 mg

EXPECTORANT

40 EXTENDED-RELEASE TABLETS

Mucinex™
美清痰™
Extended-release tablets 600 mg
Guaifenesin 600 mg

EXPECTORANT 化痰藥片

12
HOUR

✓ Thins And Loosens Mucus ✓ 有效化痰
✓ Relieves Chest Congestion ✓ 舒緩積痰咳嗽

40 EXTENDED-RELEASE TABLETS



9550 9723

MAL10021806X
HK-59774

032310 0273788

Reckitt
Benckiser

Manufacturer: Reckitt Benckiser Healthcare International Limited
Thane Road, Nottingham, NG90 2DB, United Kingdom
For/Imported by: Reckitt Benckiser (M) Sdn. Bhd. (175141-H), Menara UAC,
Mutiara Damansara, Selangor, 47800, Malaysia.
Reckitt Benckiser (S) Pte. Ltd. (197702628K), Guthrie House, Singapore 268802.
Reckitt Benckiser (HK) Ltd. (07279860), Hong Kong.

Drug Facts

Active ingredient (in each extended-release tablet)
Guaifenesin 600 mg.....Expectorant

Uses
■ helps loosen phlegm and thin bronchial secretions
■ symptomatic relief of deep chesty coughs
■ expectorant for productive cough
有效化解濃痰及支氣管分泌物
舒緩咳嗽不適症狀
幫助咳出痰沫

Warnings
Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough accompanied by too much phlegm (mucus)
Stop use and ask a doctor if
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.
If pregnant or breast-feeding, ask a health professional before use.
In case of overdose, get medical help or contact a Poison Control Center right away.
Possible side effects
■ nausea ■ abdominal discomfort ■ vomiting
Tell your doctor if you have any side effects after taking this medicine, including any not listed here.

Drug Facts (continued)

Directions
■ do not crush, chew, or break tablet
■ swallow tablet whole with a full glass of water
■ adults and children 12 years of age and over: 1 or 2 tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
■ children under 12 years of age: do not use
■ if allergic to any of the ingredients: do not use
服用方法:
用水吞服。請勿壓碎，咀嚼或碾破藥片。只供口服。
劑量:
十二歲或以上青少年及成人，每十二小時一至二片。二十四小時內，不可超過四片。十二歲以下兒童不宜服用。

Other information
■ tamper evident: do not use if carton is open or if printed seal on blister is broken or missing
■ do not store above 30°C
請勿儲存於30°C以上。
Keep out of reach of children.
Jauhi dari kanak-kanak.

Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

EXPECTORANT

Mucinex™
Extended-release tablets 600 mg
Guaifenesin 600 mg

Unique
Bi-Layer
Tablet
actual size

Extended-Release Layer

Immediate-Release Layer