



Hirudoid®

**Percutaneous anti-thrombotic,
anti-coagulant and anti inflammatory remedy.**

Hirudoid contains active ingredient Mucopolysaccharide Polysulphate (MPS) as an anti-coagulant substance activating fibrinolysis. This special formulation allows MPS to pass through the skin so that its valuable properties can evolve their full activity.

Hirudoid brings about a marked anti-inflammatory effect activates the blood circulation, encourages the absorption of haematomata and swellings, alleviates pain and relieves tension in the affected area promotes the regeneration of the connective tissue does not harm the tissue and is well tolerated.

Hirudoid does not irritate the skin reduces the duration of the disease.

Composition

Mucopolysaccharide polysulphate (MPS)
(Organo-Heparinoid)
Excip.

0.3 g
ad 100 g

Indications

- **superficial phlebitis and thrombosis of the veins**
- **inflammation of varicose veins**
- **after treatment of scleroses varices**
- **general treatment of the veins** (prevention of phlebitis)

- **haematomata**
- **swellings**
- **softening of hard scars**
- **to improve the formation of scars** (due to injuries, burns, operation, etc.)

Application

unless the doctor prescribes otherwise, a ribbon of Hirudoid about 3 to 5 cm in length (if necessary even more) is applied to the affected area 2 to 3 times daily and made to enter the skin by slight friction till the cream has disappeared.

Hirudoid Cream should only be used on unbroken skin and should not be brought into contact with open wounds, the eyes or mucous membranes.

In very painful inflammations and thrombosis, the affected area and its vicinity must carefully spread over with Hirudoid and covered with a piece of linen soaked with it. To intensify its effect, Hirudoid should be rubbed onto the affected and surrounding area.

To loosen hard scars, the cream is massaged vigorously into them.

Side effects

In single cases local skin irritation or allergies can occur.

Precaution

None.

Further information available through doctors and pharmacists.

Storage condition

Do not freeze but store below 25°C.

Commercial forms (Not all presentations are available.)

Packages containing 5 g, 14 g and 40 g of cream.

Manufacturers:

OLJC (Thailand) Limited
166 Moo 16 Bangpa-In Industrial Estate,
Udomsarak Road, Bangpa-In District,
Ayutthaya Province, Thailand
For : Medinova AG, Switzerland

 **medinova**
SWITZERLAND

