

## Package Leaflet: Information for the user

### Simponi® 50 mg solution for injection in pre-filled pen golimumab

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Simponi® is and what it is used for
2. What you need to know before you use Simponi®
3. How to use Simponi®
4. Possible side effects
5. How to store Simponi®
6. Contents of the pack and other information

#### **1. What Simponi® is and what it is used for**

Simponi® contains the active substance called golimumab.

Simponi® belongs to a group of medicines called 'TNF blockers'. It is used **in adults** for the treatment of the following inflammatory diseases:

- Rheumatoid arthritis
- Psoriatic arthritis
- Axial spondyloarthritis, including ankylosing spondylitis and non-radiographic axial spondyloarthritis
- Ulcerative colitis

**In children** weighing at least 40 kg, Simponi® is used for the treatment of polyarticular juvenile idiopathic arthritis.

Simponi® works by blocking the action of a protein called 'tumour necrosis factor alpha' (TNF- $\alpha$ ). This protein is involved in inflammatory processes of the body, and blocking it can reduce the inflammation in your body.

#### **Rheumatoid Arthritis**

Rheumatoid arthritis is an inflammatory disease of the joints. If you have active rheumatoid arthritis you will first be given other medicines. If you do not respond well enough to these medicines, you may be given Simponi® which you will take in combination with another medicine called methotrexate to:

- Reduce the signs and symptoms of your disease.
- Slow down the damage to your bones and joints.
- Improve your physical function.

#### **Psoriatic Arthritis**

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis, an inflammatory disease of the skin. If you have active psoriatic arthritis you will first be given other medicines. If you do not respond well enough to these medicines, you may be given Simponi® to:

- Reduce the signs and symptoms of your disease.
  - Slow down the damage to your bones and joints.
  - Improve your physical function.
-

## **Ankylosing Spondylitis and non-radiographic axial spondyloarthritis**

Ankylosing spondylitis and non-radiographic axial spondyloarthritis are inflammatory diseases of the spine. If you have ankylosing spondylitis or non-radiographic axial spondyloarthritis you will first be given other medicines. If you do not respond well enough to these medicines, you may be given Simponi<sup>®</sup> to:

- Reduce the signs and symptoms of your disease.
- Improve your physical function.

## **Ulcerative Colitis**

Ulcerative colitis is an inflammatory disease of the bowel. If you have ulcerative colitis you will first be given other medicines. If you do not respond well enough to these medicines, you will be given Simponi<sup>®</sup> to treat your disease.

## **Polyarticular juvenile idiopathic arthritis**

Polyarticular juvenile idiopathic arthritis is an inflammatory disease that causes joint pain and swelling in children. If your child has polyarticular juvenile idiopathic arthritis your child will first be given other medicines. If your child does not respond well enough to these medicines, your child will be given Simponi<sup>®</sup> in combination with methotrexate to treat the disease.

## **2. What you need to know before you use Simponi<sup>®</sup>**

Do not use Simponi<sup>®</sup>:

- If you are allergic (hypersensitive) to golimumab or any of the other ingredients of this medicine (listed in Section 6).
- If you have tuberculosis (TB) or any other severe infection.
- If you have moderate or severe heart failure.

If you are not sure if any of the above applies to you, talk to your doctor, pharmacist or nurse before using Simponi<sup>®</sup>.

## **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before using Simponi<sup>®</sup>.

### Infections

Tell your doctor straight away if you already have or get any symptoms of infection, during or after your treatment with Simponi<sup>®</sup>. Symptoms of infection include fever, cough, shortness of breath, flu-like symptoms, diarrhoea, wounds, dental problems or a burning feeling when urinating.

- You may get infections more easily while using Simponi<sup>®</sup>.
- Infections may progress more rapidly and may be more severe. In addition, some previous infections may reappear.

### *Tuberculosis (TB)*

Tell your doctor straight away if symptoms of TB appear during or after your treatment.

Symptoms of TB include persistent cough, weight loss, tiredness, fever or night sweats.

- Cases of TB have been reported in patients treated with Simponi<sup>®</sup>, in rare occasions even in patients who have been treated with medicines for TB. Your doctor will test you to see if you have TB.
- It is very important that you tell your doctor if you have ever had TB, or if you have been in close contact with someone who has had or has TB.
- If your doctor feels that you are at risk of TB, you may be treated with medicines for TB before you begin using Simponi<sup>®</sup>.

### *Hepatitis B virus (HBV)*

- Tell your doctor if you are a carrier or if you have or have had HBV before you are given Simponi<sup>®</sup>.
  - Tell your doctor if you think you might be at risk of contracting HBV.
  - Your doctor should test you for HBV.
  - Treatment with TNF blockers such as Simponi<sup>®</sup> may result in reactivation of HBV in patients who carry this virus, which can be life-threatening in some cases.
-

### *Invasive fungal infections*

If you have lived in or travelled to an area where infections caused by specific type of fungi that can affect the lungs or other parts of the body (called histoplasmosis, coccidioidomycosis, or blastomycosis), are common, tell your doctor straight away. Ask your doctor if you don't know if these fungal infections are common in the area in which you have lived or travelled.

### Cancer and lymphoma

Tell your doctor if you have ever been diagnosed with lymphoma (a type of blood cancer) or any other cancer before you use Simponi®.

- If you use Simponi® or other TNF blockers, your risk for developing lymphoma or another cancer may increase.
- Patients with severe rheumatoid arthritis and other inflammatory diseases, who have had the disease for a long time, may be at higher than average risk of developing lymphoma.
- There have been cases of cancers, including unusual types, in children and teenage patients taking TNF-blocking agents, which sometimes resulted in death.
- On rare occasions, a specific and severe type of lymphoma called hepatosplenic T-cell lymphoma has been observed in patients taking other TNF-blockers. Most of these patients were adolescent or young adult males. This type of cancer has usually resulted in death. Almost all of these patients had also received medicines known as azathioprine or 6-mercaptopurine.
- Tell your doctor if you are taking azathioprine or 6-mercaptopurine with Simponi®.
- Patients with severe persistent asthma, chronic obstructive pulmonary disease (COPD), or are heavy smokers may be at increased risk for cancer with Simponi® treatment. If you have severe persistent asthma, COPD or are a heavy smoker, you should discuss with your doctor whether treatment with a TNF blocker is appropriate for you.
- Some patients treated with golimumab have developed certain kinds of skin cancer. If any changes in the appearance of the skin or growths on the skin occur during or after therapy, tell your doctor.

### Heart failure

Tell your doctor straight away if you get new or worsening symptoms of heart failure. Symptoms of heart failure include shortness of breath or swelling of your feet.

- New and worsening congestive heart failure has been reported with TNF blockers, including Simponi®. Some of these patients died.
- If you have mild heart failure and you are being treated with Simponi®, you must be closely monitored by your doctor.

### Nervous system disease

Tell your doctor straight away if you have ever been diagnosed with or develop symptoms of a demyelinating disease such as multiple sclerosis. Symptoms may include changes in your vision, weakness in your arms or legs or numbness or tingling in any part of your body. Your doctor will decide if you should receive Simponi®.

### Operations or dental procedures

- Talk to your doctor if you are going to have any operations or dental procedures.
- Tell your surgeon or dentist performing the procedure that you are having treatment with Simponi®.

### Autoimmune disease

Tell your doctor if you develop symptoms of a disease called lupus. Symptoms include persistent rash, fever, joint pain and tiredness.

- On rare occasions, people treated with TNF blockers have developed lupus.

### Blood disease

In some patients the body may fail to produce enough of the blood cells that help your body fight infections or help you to stop bleeding. If you develop a fever that does not go away, bruise or bleed very easily or look very pale, call your doctor right away. Your doctor may decide to stop treatment.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before using Simponi®.

---

### Vaccinations

Talk to your doctor if you have had, or are due to have a vaccine.

- You should not receive certain (live) vaccines while using Simponi®.
- Certain vaccinations may cause infections. If you received Simponi® while you were pregnant, your baby may be at higher risk for getting such an infection for up to approximately six months after the last dose you received during pregnancy. It is important that you tell your baby's doctors and other health care professionals about your Simponi® use so they can decide when your baby should receive any vaccine.

Talk to your child's doctor regarding vaccinations for your child. If possible, your child should be up to date with all vaccinations before using Simponi®.

### Therapeutic infectious agents

Talk to your doctor if you have recently received or are scheduled to receive treatment with a therapeutic infectious agent (such as BCG instillation used for the treatment of cancer).

### Allergic reactions

Tell your doctor straight away if you develop symptoms of an allergic reaction after your treatment with Simponi®. Symptoms of an allergic reaction may include swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing, skin rash, hives, swelling of the hands, feet or ankles.

- Some of these reactions may be serious or, rarely, life-threatening.
- Some of these reactions occurred after the first administration of Simponi®.

### **Children and adolescents**

Simponi® is not recommended for children weighing less than 40 kg with polyarticular juvenile idiopathic arthritis or in children and adolescents younger than 18 years for any other condition.

### **Other medicines and Simponi®**

- Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines, including any other medicines to treat rheumatoid arthritis, polyarticular juvenile idiopathic arthritis, psoriatic arthritis, ankylosing spondylitis, non-radiographic axial spondyloarthritis or ulcerative colitis.
- You should not take Simponi® with medicines containing the active substance anakinra or abatacept. These medicines are used for the treatment of rheumatic diseases.
- Tell your doctor or pharmacist if you are taking any other medicines that affect your immune system.
- You should not receive certain (live) vaccines while using Simponi®.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before using Simponi®.

### **Pregnancy and breast-feeding**

Talk to your doctor before using Simponi® if:

- You are pregnant or are planning to become pregnant while using Simponi®. The effects of this medicine in pregnant women are not known. The use of Simponi® in pregnant women is not recommended. If you are being treated with Simponi®, you must avoid becoming pregnant by using adequate contraception during your treatment and for at least 6 months after the last Simponi® injection.
- Before starting breast-feeding, your last treatment with Simponi® must be at least 6 months ago. You must stop breast-feeding if you are to be given Simponi®.
- If you received Simponi® during your pregnancy, your baby may have a higher risk for getting an infection. It is important that you tell your baby's doctors and other health care professionals about your Simponi® use before the baby receives any vaccine (for more information see section on vaccination).

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

Simponi® may have a minor influence on your ability to drive and use tools or machines. Dizziness may occur after you take Simponi®. If this happens, do not drive or use any tools or machines.

---

## **Simponi® contains latex and sorbitol**

### Latex sensitivity

A part of the pre-filled pen, the needle cover, contains latex. Because latex may cause severe allergic reactions, talk to your doctor before using Simponi® if you or your carer are allergic to latex.

### Sorbitol intolerance

Simponi® contains sorbitol (E420). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

## **3. How to use Simponi®**

Always use this medicine exactly as your doctor or pharmacist has told you. You should check with your doctor or pharmacist if you are not sure.

### **How much Simponi® is given**

Rheumatoid arthritis, psoriatic arthritis, and axial spondyloarthritis, including ankylosing spondylitis and non-radiographic axial spondyloarthritis:

- The recommended dose is 50 mg (the content of one pre-filled pen) given once a month, on the same date each month.
- Talk to your doctor before taking your fourth dose. Your doctor will determine if you should continue Simponi® treatment.
  - If you weigh more than 100 kg, the dose might be increased to 100 mg (the content of 2 pre-filled pens) given once a month, on the same date each month.

Polyarticular juvenile idiopathic arthritis:

- The recommended dose is 50 mg given once a month, on the same date each month.
- Talk to your child's doctor before your child takes their fourth dose. Your child's doctor will determine if they should continue Simponi® treatment.

Ulcerative colitis

- The table below shows how you will usually use this medicine.

Initial Treatment	A starting dose of 200mg (the contents of 4 pre-filled pens) followed by 100mg (the contents of 2 pre-filled pens) 2 weeks later.
Maintenance Treatment	<ul style="list-style-type: none"><li>• In patients weighing less than 80kg, 50mg (the contents of 1 pre-filled pen) 4 weeks after your last treatment, then every 4 weeks thereafter. Your doctor may decide to prescribe 100 mg (the contents of 2 pre-filled pens), depending on how well Simponi® works for you.</li><li>• In patients weighing 80kg or more, 100mg (the contents of 2 pre-filled pens) 4 weeks after your last treatment, then every 4 weeks thereafter.</li></ul>

### **How Simponi® is given**

- Simponi® is given by injection under the skin (subcutaneously).
- At the start, your doctor or nurse may inject Simponi®. However, you and your doctor may decide that you may inject Simponi® yourself. In this case you will get training on how to inject Simponi® yourself.

Talk to your doctor if you have any questions about giving yourself an injection. You will find detailed "Instructions for administration" at the end of this leaflet.

### **If you use more Simponi® than you should**

If you have used or been given too much Simponi® (either by injecting too much on a single occasion, or by using it too often), talk to your doctor or pharmacist straight away. Always take the outer carton and this leaflet with you, even if it is empty.

### **If you forget to use Simponi®**

If you forget to use Simponi® on your planned date, inject the forgotten dose as soon as you remember. Do not use a double dose to make up for a forgotten dose.

---

When to inject your next dose:

- If you are less than 2 weeks late, inject the forgotten dose as soon as you remember and stay on your original schedule.
- If you are more than 2 weeks late, inject the forgotten dose as soon as you remember and talk to your doctor or pharmacist to ask when you need to take the next dose.

If you are not sure what to do, talk to your doctor or pharmacist.

### **If you stop using Simponi®**

If you are considering stopping Simponi®, talk to your doctor or pharmacist first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some patients may experience serious side effects and may require treatment. The risk of certain side effects is greater with the 100 mg dose compared with the 50 mg dose. Side effects may appear up to several months after the last injection.

Tell your doctor straight away if you notice any of the following serious side effects of Simponi® which include:

- **allergic reactions which may be serious, or rarely, life-threatening (rare).** Symptoms of an allergic reaction may include swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing, skin rash, hives, swelling of the hands, feet or ankles. Some of these reactions occurred after the first administration of Simponi®.
- **serious infections (including TB, bacterial infections including serious blood infections and pneumonia, severe fungal infections and other opportunistic infections) (common).** Symptoms of an infection can include fever, tiredness, (persistent) cough, shortness of breath, flu-like symptoms, weight loss, night sweats, diarrhoea, wounds, dental problems and a burning feeling when urinating.
- **reactivation of hepatitis B virus if you are a carrier or have had hepatitis B before (rare).** Symptoms can include yellowing of the skin and eyes, dark brown-coloured urine, right-sided abdominal pain, fever, feeling sick, being sick, and feeling very tired.
- **nervous system disease such as multiple sclerosis (rare).** Symptoms of nervous system disease can include changes in your vision, weakness in your arms or legs, numbness or tingling in any part of your body.
- **cancer of the lymph nodes (lymphoma) (rare).** Symptoms of lymphoma can include swelling of the lymph nodes, weight loss, or fever.
- **heart failure (rare).** Symptoms of heart failure can include shortness of breath or swelling of your feet.
- **signs of immune system disorders called:**
  - **lupus (rare).** Symptoms can include joint pain or a rash on cheeks or arms that is sensitive to the sun.
  - **sarcoidosis (rare).** Symptoms can include a persistent cough, being short of breath, chest pain, fever, swelling of your lymph nodes, weight loss, skin rashes, and blurred vision.
- **swelling of small blood vessels (vasculitis) (rare).** Symptoms can include fever, headache, weight loss, night sweats, rash, and nerve problems such as numbness and tingling.
- **skin cancer (uncommon).** Symptoms of skin cancer can include changes in the appearance of your skin or growths on your skin.
- **blood disease (common).** Symptoms of blood disease can include a fever that does not go away, bruising or bleeding very easily or looking very pale.
- **blood cancer (leukaemia) (rare).** Symptoms of leukaemia can include fever, feeling tired, frequent infections, easy bruising, and night sweats.

Tell your doctor straight away if you notice any of the above symptoms.

### **The following additional side effects have been observed with Simponi®:**

Very common side effects (may affect more than 1 in 10 people):

---

- Upper respiratory tract infections, sore throat or hoarseness, runny nose

Common side effects (may affect up to 1 in 10 people):

- Abnormal liver tests (increased liver enzymes) found during blood tests done by your doctor
- Feeling dizzy
- Headache
- Feeling numb or having a tingling feeling
- Superficial fungal infections
- Abscess
- Bacterial infections (such as cellulitis)
- Low red blood cell counts
- Low white blood cell counts
- Positive blood lupus test
- Allergic reactions
- Indigestion
- Stomach pain
- Feeling sick (nausea)
- Flu
- Bronchitis
- Sinus infection
- Cold sores
- High blood pressure
- Fever
- Asthma, shortness of breath, wheezing
- Stomach and bowel disorders which include inflammation of the stomach lining and colon which may cause fever
- Pain and ulcers in the mouth
- Injection site reactions (including redness, hardness, pain, bruising, itching, tingling and irritation)
- Hair loss
- Rash and itching of the skin
- Difficulty sleeping
- Depression
- Feeling weak
- Bone fractures
- Chest discomfort

Uncommon side effects (may affect up to 1 in 100 people):

- Kidney infection
  - Cancers, including skin cancer and non-cancerous growths or lumps, including skin moles
  - Skin blisters
  - Psoriasis (including on the palms of your hand and/or the soles of your feet and/or in the form of skin blisters)
  - Low platelet count
  - Combined low platelet, red, and white blood cell count
  - Thyroid disorders
  - Increase in blood sugar levels
  - Increase in blood cholesterol levels
  - Balance disorders
  - Vision disturbances
  - Sensation of heart beating irregularly
  - Narrowing of the blood vessels in the heart
  - Blood clots
  - Flushing
  - Constipation
  - Chronic inflammatory condition of the lungs
  - Acid reflux
  - Gall stones
  - Liver disorders
  - Breast disorders
  - Menstrual disorders
-

Rare side effects (may affect up to 1 in 1,000 people):

- Failure of the bone marrow to produce blood cells
- Severely decreased number of white blood cells
- Infection of the joints or the tissue around them
- Impaired healing
- Inflammation of blood vessels in internal organs
- Leukaemia
- Melanoma (a type of skin cancer)
- Merkel cell carcinoma (a type of skin cancer)
- Lichenoid reactions (itchy reddish-purple skin rash and/or threadlike white-grey lines on mucous membranes)
- Scaly, peeling skin
- Immune disorders that could affect the lungs, skin and lymph nodes (most commonly presenting as sarcoidosis)
- Pain and discoloration in the fingers or toes
- Taste disturbances
- Bladder disorders
- Kidney disorders
- Inflammation of the blood vessels in your skin which results in rash

Side effects of which the frequency is not known:

- A rare blood cancer affecting mostly young people (hepatosplenic T-cell lymphoma)

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Simponi®**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the label and the carton after “EXP”. The expiry date refers to the last day of that month.
- Store in a refrigerator (2°C-8°C). Do not freeze.
- Keep the pre-filled syringe/pen in the outer carton in order to protect it from light.
- This medicine can also be stored out of the refrigerator at temperatures up to a maximum of 25°C for a single period of up to 30 days, but not beyond the original expiry date printed on the carton. Write the new expiry date on the carton including day/month/year (no more than 30 days after the medicine is removed from the refrigerator). Do not return this medicine to refrigerator if it has reached room temperature. Discard this medicine if not used by the new expiry date or the expiry date printed on the carton, whichever is earlier.
- Do not use this medicine if you notice that the liquid is not a clear to light yellow colour, cloudy, or contains foreign particles.
- Do not throw away any medicines via wastewater or household waste. Ask your doctor or pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

**6. Contents of the pack and other information**

**What Simponi® contains**

The active substance is golimumab. One 0.5 mL pre-filled pen contains 50 mg of golimumab.

The other ingredients are sorbitol (E420), L-histidine, L-histidine monohydrochloride monohydrate, polysorbate 80 and water for injections.

**What Simponi® looks like and contents of the pack**

Simponi® is supplied as solution for injection in a single-use pre-filled pen. Simponi® is available in packs containing 1 pre-filled pen and multipacks containing 3 (3 packs of 1) pre-filled pens. Not all pack sizes may be marketed.

---



The solution is clear to slightly opalescent (having a pearl-like shine), colourless to light yellow and may contain a few small translucent or white particles of protein. Do not use Simponi® if the solution is discoloured, cloudy or you can see foreign particles in it.

**Batch Releaser**

Cilag AG  
Hochstrasse 201  
CH-8200 Schaffhausen  
Switzerland

**Product Registrant**

Johnson & Johnson International (Singapore) Pte. Ltd.  
2 Science Park Drive  
#07-13, Ascent  
Singapore Science Park 1  
Singapore 118222

**Last Revision of Text**

02 December 2022 (CPPI 20 September 2022)

**INSTRUCTIONS FOR ADMINISTRATION**

**If you would like to self-inject Simponi®, you must be trained by a healthcare professional to prepare an injection and give it to yourself. If you have not been trained, please contact your doctor, nurse or pharmacist to schedule a training session.**

In these instructions:

1. Preparing for use of the pre-filled pen
2. Choosing and preparing the injection site
3. Injecting the medicine
4. After the injection

The diagram below (see figure 1) shows what the “SmartJect” pre-filled pen looks like:




Figure 1

## 1. Preparing for use of the pre-filled pen

- Do **NOT** shake the pre-filled pen at any time.
- Do **NOT** remove the cap from the pre-filled pen until immediately before the injection.

### Check the number of pre-filled pens

Check the pre-filled pens to make sure

- the number of pre-filled pens and strength is correct.
  - If your dose is 50 mg, you will get one 50 mg pre-filled pen.
  - If your dose is 100 mg, you will get two 50 mg pre-filled pens and you will need to give yourself two injections. Choose two different sites for these injections (e.g. one injection in the right thigh and the other injection in the left thigh), and give the injections one right after the other.
  - If your dose is 200 mg, you will get four 50 mg pre-filled pens and you will need to give yourself four injections. Choose different sites for these injections and give the injections one right after the other.

### Check expiry date

- Check the expiration date printed or written on the carton.
- Check the expiration date (as indicated as “EXP”) on the pre-filled pen.
- **Do not** use the pre-filled pen if the expiration date has passed. The printed expiration refers to the last day of the month. Please contact your doctor or pharmacist for assistance.

### Check security seal

- Check the security seal around the cap of the pre-filled pen.
- **Do not** use the pre-filled pen if the seal is broken. Please contact your doctor or pharmacist.

### Wait 30 minutes to allow pre-filled pen to reach room temperature

- To ensure proper injection, allow the pre-filled pen to sit at room temperature outside the box for 30 minutes out of the reach of children.
- Do **NOT** warm the pre-filled pen in any other way (for example, do **NOT** warm it in a microwave or in hot water).
- Do **NOT** remove the pre-filled pen cap while allowing it to reach room temperature.

### Get the rest of your equipment ready

- While you are waiting you can get the rest of your equipment ready, including an alcohol swab, a cotton ball or gauze, and a sharps container.

### Check the liquid in the pre-filled pen

- Look through the viewing window to make sure that the liquid in the pre-filled pen is clear to slightly opalescent (having a pearl-like shine) and colourless to light yellow. The solution can be used if it contains a few small translucent or white particles of protein.
- You will also notice an air bubble, which is normal.
- Do **NOT** use the pre-filled pen if the liquid is the wrong colour, cloudy or contains larger particles. If this happens, talk to your doctor or pharmacist.

## 2. Choosing and preparing the injection site (see figure 2)

- You can inject the medicine into the front of the middle thighs.
  - You can also use the stomach (abdomen) below the belly button, except for approximately the 5 cm area directly underneath the belly button.
  - Do **NOT** inject into the arm.
  - Do **NOT** inject into areas where the skin is tender, bruised, red, scaly, hard or has scars or stretch marks.
  - If multiple injections are required for a single administration, the injections should be administered at different sites on the body.
-



Figure 2

### Preparing injection site

- Wash your hands thoroughly with soap and warm water.
- Wipe the injection site with an alcohol swab.
- Allow the skin to dry before injecting. Do **NOT** fan or blow on the clean area.
- Do **NOT** touch this area again before giving the injection.

### 3. Injecting the medicine

- The cap should **NOT** be removed until you are ready to inject the medicine.
- The medicine should be injected within 5 minutes after the cap has been removed.

### Remove the cap (see figure 3)

- When you are ready to inject, twist the cap slightly to break the security seal.
- Pull the cap off and throw it away after your injection.
- Do **NOT** put the cap back on because it may damage the needle inside the pre-filled pen.
- Do not use pre-filled pen if it is dropped without the cap in place. If this happens, please contact your doctor or pharmacist for assistance.




Figure 3

### Push the pre-filled pen firmly against the skin (see figures 4 and 5)

- Hold the pre-filled pen comfortably in your hand. Do **NOT** press the button at this time.
- Push the open end of the pre-filled pen firmly against the skin at a 90-degree angle.
- Do **NOT** pinch the skin.
- Do **NOT** press the button until **after** the pre-filled pen is pushed firmly against the skin and the Safety Sleeve slides fully into the Clear Cover.




Figure 4

- Inject without pinching the skin.




Figure 5

**Press button to inject (see figure 6)**

- Continue to push the pre-filled pen firmly against the skin and press the front raised part of the button with your fingers or thumb. You will not be able to press in the button unless the pre-filled pen is pushed firmly against your skin and the Safety Sleeve slides into the Clear Cover.
- Once the button is pressed, it will remain pressed in so you do not need to keep pressure on it.




Figure 6

- **You will hear a loud 'click' sound – don't be alarmed.** The first loud 'click' indicates that the needle has been inserted and the injection has started. You may or may not feel a needle prick at this time.

**DO NOT lift the pre-filled pen away from your skin. If you pull the pre-filled pen away from the skin, you may not get your full dose of medicine.**

**Continue to hold until the second 'click' (see figure 7)**

- **Continue to hold the pre-filled pen down firmly against the skin until you hear the second 'click'. This usually takes about 3-6 seconds, but it may take up to 15 seconds for you to hear the second 'click' sound.**
- The second 'click' means that the injection is finished and the needle has gone back into the pre-filled pen. **Note:** If you have a hearing problem and do not hear the second 'click', count 15

seconds from the time you first press the button and then lift the pre-filled pen from the injection site.

- Lift the pre-filled pen from the injection site.




Figure 7

#### 4. After the injection

##### Use a cotton ball or gauze

- There may be a small amount of blood or liquid at the injection site. This is normal.
- You can press a cotton ball or gauze over the injection site for 10 seconds.
- You may cover the injection site with a small adhesive bandage, if necessary.
- Do **NOT** rub your skin.

##### Check the window – a yellow indicator confirms proper administration (see figure 8)

- The yellow indicator is connected to the plunger of the pre-filled pen. If the yellow indicator is not shown in the window, the plunger has not advanced adequately, and the injection has not occurred.
- The yellow indicator will fill about half of the viewing window. This is normal.
- Talk to your doctor or pharmacist if the yellow indicator is not visible in the window or if you suspect that you may not have received a complete dose. Do **NOT** administer a second dose without speaking to your doctor.




Figure 8

**Throw the pre-filled pen away (see figure 9)**

- Place your pen in a sharps container straight away. Make sure you dispose of the bin as instructed by your doctor or nurse when the container is full.

If you feel that something has gone wrong with the injection or if you are not sure, talk to your doctor or pharmacist.



Figure 9